

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

75313


DRAFT FINAL PRINTED LABELING

IPRATROPIUM BROMIDE INHALATION SOLUTION 0.02%

PRESCRIBING INFORMATION

Rx only

DESCRIPTION: The active ingredient in Ipratropium Bromide Inhalation Solution is ipratropium bromide monohydrate. It is an anticholinergic bronchodilator chemically described as 8-azoniabicyclo[3.2.1]octane, 3-(3-hydroxy-1-oxo-2-phenylpropoxy)-8-methyl-8-(1-methylethyl)-, bromide, monohydrate (*endo, syn*)-, (±); a synthetic quaternary ammonium compound, chemically related to atropine.



Ipratropium bromide is a white crystalline substance, freely soluble in water and lower alcohols. It is a quaternary ammonium compound and thus exists in an ionized state in aqueous solutions. It is relatively insoluble in non-polar media.

Ipratropium Bromide Inhalation Solution is administered by oral inhalation with the aid of a nebulizer. It contains ipratropium bromide 0.02% (anhydrous basis) in a sterile, preservative free, isotonic saline solution, pH-adjusted to 3.4 (3 to 4) with hydrochloric acid.

CLINICAL PHARMACOLOGY: Ipratropium bromide is an anticholinergic (parasympatholytic) agent that, based on animal studies, appears to inhibit vagally mediated reflexes by antagonizing the action of acetylcholine, the transmitter agent released from the vagus nerve.

Anticholinergics prevent the increases in intracellular concentration of cyclic guanosine monophosphate (cyclic GMP) that are caused by interaction of acetylcholine with the muscarinic receptor on bronchial smooth muscle.

The bronchodilation following inhalation of ipratropium bromide is primarily a local, site-specific effect, not a systemic one. Much of an administered dose is swallowed but not absorbed, as shown by fecal excretion studies. Following nebulization of a 2 mg dose, a mean 7% of the dose was absorbed into the systemic circulation either from the surface of the lung or from the gastrointestinal tract. The half-life of elimination is about 1.6 hours after intravenous administration. Ipratropium bromide is minimally (0 to 9% *in vitro*) bound to plasma albumin and α_1 -acid glycoproteins. It is partially metabolized. Autoradiographic studies in rats have shown that ipratropium bromide does not penetrate the blood-brain barrier. Ipratropium bromide has not been studied in patients with hepatic or renal insufficiency. It should be used with caution in those patient populations.

In controlled 12-week studies in patients with bronchospasm associated with chronic obstructive pulmonary disease (chronic bronchitis and emphysema) significant improvements in pulmonary function (FEV₁ increases of 15% or more) occurred within 15 to 30 minutes, reached a peak in 1-2 hours, and persisted for periods of 4-5 hours in the majority of patients, with about 25%-38% of the patients demonstrating increases of 15% or more for at least 7-8 hours. Continued effectiveness of Ipratropium Bromide Inhalation Solution was demonstrated throughout the 12-week period. In addition, significant increases in forced vital capacity (FVC) have been demonstrated. However, ipratropium bromide did not consistently produce significant improvement in subjective symptom scores nor in quality of life scores over the 12-week duration of study.

Additional controlled 12-week studies were conducted to evaluate the safety and effectiveness of Ipratropium Bromide Inhalation Solution administered concomitantly with the beta adrenergic bronchodilator solutions metaproterenol and albuterol compared with the administration of each of the beta agonists alone. Combined therapy produced significant additional improvement in FEV₁ and FVC.

On combined therapy, the median duration of 15% improvement in FEV₁ was 5-7 hours, compared with 3-4 hours in patients receiving a beta agonist alone.

INDICATIONS AND USAGE: Ipratropium Bromide Inhalation Solution administered either alone or with other bronchodilators, especially beta adrenergics, is indicated as a bronchodilator for maintenance treatment of bronchospasm associated with chronic obstructive pulmonary disease including chronic bronchitis and emphysema.

CONTRAINDICATIONS: Ipratropium bromide is contraindicated in known or suspected cases of hypersensitivity to ipratropium bromide, or to atropine and its derivatives.

WARNINGS: The use of Ipratropium Bromide Inhalation Solution as a single agent for the relief of bronchospasm in acute COPD exacerbation has not been adequately studied. Drugs with faster onset of action may be preferable as initial therapy in this situation. Combination of ipratropium bromide and beta agonists has not been shown to be more effective than either drug alone in reversing the bronchospasm associated with acute COPD exacerbation.

Immediate hypersensitivity reactions may occur after administration of ipratropium bromide as demonstrated by rare cases of urticaria, angioedema, rash, bronchospasm and oropharyngeal edema.

PRECAUTIONS: General: Ipratropium bromide should be used with caution in patients with narrow angle glaucoma, prostatic hypertrophy or bladder-neck obstruction.

Information for Patients: Patients should be advised that temporary blurring of vision, precipitation or worsening of narrow-angle glaucoma or eye pain may result if the solution comes into direct contact with the eyes. Use of a nebulizer with mouthpiece rather than face mask may be preferable, to reduce the likelihood of the nebulizer solution reaching the eyes. Patients should be advised that Ipratropium Bromide Inhalation Solution can be mixed in the nebulizer with albuterol or metaproterenol if used within one hour. Drug stability and safety of Ipratropium Bromide Inhalation Solution when mixed with other drugs in a nebulizer have not been established. Patients should be reminded that Ipratropium Bromide Inhalation Solution should be used consistently as prescribed throughout the course of therapy.

Drug Interactions: Ipratropium bromide has been shown to be a safe and effective bronchodilator when used in conjunction with beta adrenergic bronchodilators. Ipratropium bromide has also been used with other pulmonary medications, including methylxanthines and corticosteroids without adverse drug interactions.

Carcinogenesis, Mutagenesis, Impairment of Fertility: Two-year oral carcinogenicity studies in rats and mice have revealed no carcinogenic potential at dietary doses up to 6 mg/kg/day of ipratropium bromide.

Results of various mutagenicity studies (Ames test, mouse dominant lethal test, mouse micronucleus test and chromosome aberration of bone marrow in Chinese hamsters) were negative.






Fertility of male or female rats at oral doses up to 50 mg/kg/day was unaffected by ipratropium bromide administration. At doses above 90 mg/kg, increased resorption and decreased conception rates were observed.

Pregnancy: TERATOGENIC EFFECTS

Pregnancy Category B. Oral reproduction studies performed in mice, rats and rabbits at doses of 10, 100 and 125 mg/kg respectively, and inhalation reproduction studies in rats and rabbits at doses of 1.5 and 1.8 mg/kg (or approximately 38 and 45 times the recommended human daily dose) respectively, have demonstrated no evidence of teratogenic effects as a result of ipratropium bromide. However, no adequate or well-controlled studies have been conducted in pregnant women. Because animal reproduction studies are not always predictive of human response, ipratropium bromide should be used during pregnancy only if clearly needed.

Patient's Instructions for Use

IPRATROPIUM BROMIDE Inhalation Solution Not for Injection For Oral Inhalation Use Only



6. Clean the nebulizer (see manufacturer's instructions).
Instructions continued overleaf.

2

Nursing Mothers: It is not known whether ipratropium bromide is excreted in human milk. Although lipid-insoluble quaternary bases pass into breast milk, it is unlikely that ipratropium bromide would reach the infant to a significant extent, especially when taken by inhalation since ipratropium bromide is not well absorbed systemically after inhalation or oral administration. However, because many drugs are excreted in human milk, caution should be exercised when ipratropium bromide is administered to a nursing woman.

Pediatric Use: Safety and effectiveness in pediatric patients below the age of 12 have not been established.

ADVERSE REACTIONS: Adverse reaction information concerning Ipratropium Bromide Inhalation Solution is derived from 12-week active controlled clinical trials. Additional information is derived from the foreign postmarketing experience and the published literature.

All adverse events, regardless of drug relationship, reported by three percent or more patients in the 12-week controlled clinical trials appear in the table below.

Additional adverse reactions reported in less than three percent of the patients treated with ipratropium bromide include tachycardia, palpitations, eye pain, urinary retention, urinary tract infection and urticaria. Cases of precipitation or worsening of narrow-angle glaucoma and acute eye pain have been reported.

Lower respiratory adverse reactions (bronchitis, dyspnea and bronchospasm) were the most common events leading to discontinuation of ipratropium bromide therapy in the 12-week trials. Headache, mouth dryness and aggravation of COPD symptoms are more common when the total daily dose of ipratropium bromide equals or exceeds 2,000 mcg.

Allergic-type reactions such as skin rash, angioedema of tongue, lips and face, urticaria, laryngospasm and anaphylactic reaction have been reported. Many of the patients had a history of allergies to other drugs and/or foods.

OVERDOSAGE: Acute systemic overdosage by inhalation is unlikely since ipratropium bromide is not well absorbed after inhalation at up to four-fold the recommended dose, or after oral administration at up to forty-fold the recommended dose. The oral LD₅₀ of ipratropium bromide ranged between 1001 and 2010 mg/kg in mice; between 1667 and 4000 mg/kg in rats and between 400 and 1300 mg/kg in dogs.

DOSAGE AND ADMINISTRATION: The usual dosage of Ipratropium Bromide Inhalation Solution is 500 mcg (1 Unit-Dose Vial) administered three to four times a day by oral nebulization, with doses 6 to 8 hours apart. Ipratropium Bromide Inhalation Solution Unit-Dose Vials contain 500 mcg ipratropium bromide anhydrous in 2.5 mL normal saline. Ipratropium Bromide Inhalation Solution can be mixed in the nebulizer with albuterol or metaproterenol if used within one hour.

Drug stability and safety of Ipratropium Bromide Inhalation Solution when mixed with other drugs in a nebulizer have not been established.

HOW SUPPLIED: Ipratropium Bromide Inhalation Solution Unit Dose Vial is supplied as a 0.02% clear, colorless solution containing 2.5 mL with 5 vials per foil pouch supplied in cartons as listed below.

25's pack - 0172-6407-44
 60's pack - 0172-6407-49

Each vial is made from a low density polyethylene (LDPE) resin.
 Store between 15° and 30° C (59° and 86° F). Protect from light.
 Retain in foil pouch until time of use.

Manufactured by: Steripak Limited Runcorn, Cheshire WA7 1QF England
 Manufactured for: Zenith Goldline Pharmaceuticals, Inc. Miami FL 33137

0172
 11/99
 DI

	PERCENT OF PATIENTS				
	Ipratropium Bromide (500 mcg t.i.d.) n= 219	Metaproterenol (15 mg t.i.d.) n=212	Ipratropium Bromide/ Metaproterenol (500 mcg t.i.d./15 mg t.i.d.) n= 108	Albuterol (2.5 mg t.i.d.) n=205	Ipratropium Bromide/ Albuterol (500 mcg t.i.d./2.5 mg t.i.d.) n=100
Body as a Whole-General Disorders					
Headache	6.4	5.2	6.5	6.3	9.0
Pain	4.1	3.5	0.9	2.9	5.0
Influenza-like symptoms	3.7	4.7	6.5	0.5	1.0
Back Pain	3.2	1.9	1.9	2.4	0.0
Chest Pain	3.2	4.2	5.6	2.0	1.0
Cardiovascular Disorders					
Hypertension/Hypertension Aggravated	0.9	1.9	0.9	1.5	4.0
Central and Peripheral Nervous System					
Dizziness	2.3	3.3	1.9	3.9	4.0
Insomnia	0.9	0.5	4.6	1.0	1.0
Tremor	0.9	7.1	8.3	1.0	0.0
Nervousness	0.5	4.7	6.5	1.0	1.0
Gastrointestinal System Disorders					
Mouth Dryness	3.2	0.0	1.9	2.0	3.0
Nausea	4.1	3.8	1.9	2.9	2.0
Constipation	0.9	0.0	3.7	1.0	1.0
Musculo-Skeletal System Disorders					
Arthritis	0.9	1.4	0.9	0.5	3.0
Respiratory System Disorders (Lower)					
Coughing	4.6	8.0	6.5	5.4	6.0
Dyspnea	9.8	13.2	16.7	12.7	9.0
Bronchitis	14.6	24.5	15.7	16.6	20.0
Bronchospasm	2.3	2.8	4.6	5.4	5.0
Sputum Increased	1.4	1.4	4.6	3.4	0.0
Respiratory Disorder	0.0	6.1	6.5	2.0	4.0
Respiratory System Disorders (Upper)					
Upper Respiratory Tract Infection	13.2	11.3	9.3	12.2	16.0
Pharyngitis	3.7	4.2	5.6	2.9	4.0
Rhinitis	2.3	4.2	1.9	2.4	0.0
Sinusitis	2.3	2.8	0.9	5.4	4.0

* All adverse events, regardless of drug relationship, reported by three or more patients in the 12-week controlled clinical trials

Note: Use only as directed by your physician. More frequent administration or higher doses are not recommended. Ipratropium Bromide Inhalation Solution can be mixed in the nebulizer with albuterol or metaproterenol if used within one hour, but not with other drugs.

Drug stability and safety of Ipratropium Bromide Inhalation Solution when mixed with other drugs in a nebulizer have not been established.

Store between 15° and 30° C

(59° and 86° F).

Protect from light.

Retain in foil pouch until time of use.

Additional instructions:

Blank lined area for additional instructions.



©1999

1199 J



Manufactured for:
ZENITH GOLDLINE PHARMACEUTICALS, INC.
MIAMI, FL 33137
By: Stangek Limited
Runcorn, Cheshire WA7 1GF England
Star-Neb™ is a trademark of Zenith Goldline Pharmaceuticals, Inc.

KEEP OUT OF THE REACH OF CHILDREN.

Retain in foil pouch until time of use.

Store between 15° and 30° C (59° and 86° F). Protect from light.

Adjusted to pH 3.4 (3 to 4) with hydrochloric acid.

Each Low Density Polyethylene Vial Contains: 2.5 mL Ipratropium Bromide 0.02% preservative-free, sterile, isotonic aqueous solution containing sodium chloride.

package insert and dispense with solution.

ATTENTION PHARMACIST: Detach "Patient's Instructions For Use" from

Do not exceed recommended dosage.

Use only as directed by your physician.

USUAL DOSAGE: Read accompanying insert.

PC0756
PC0756
PC0756

Zenith

NDC 0172-6407-49

IPRATROPIUM BROMIDE INHALATION SOLUTION

FEB

0.02%
0.5 mg/2.5 mL

Rx Only
Star-Neb™
brand of

STERILE UNIT DOSE VIALS • NOT FOR INJECTION
FOR ORAL INHALATION USE ONLY

unmarked Area
x print

PC0756

Colours Cyan (not for print) PMS Reflex Blue PMS 226 PMS 873	Template reference Electronic artwork or Colour separated bromide artwork need not be returned. If it is retained, you are responsible for ensuring that it is securely stored.	Packaging Technology approval Signature Date / /
Job no 843/4	Other information PC 0756	Regulatory Affairs approval Signature Date / /
Date 00/00/99		
Designer 3rd Party		
Draft 05		
Revision TKS		
Rev Date 10/11/99		
Software Adobe Illustrator 8.0		

Medica	PROOF AT 10.11.99	W/T 58519
SIGNED	DATE	

10.11.99 - PROFILE CHANGED TO 196 x 92 x 130mm

Zenith

NDC 0172-6407-49

IPRATROPIUM BROMIDE
INHALATION SOLUTION

0.02%
0.5 mg/2.5 mL

Rx Only

Steri-Neb™
brand of

STERILE UNIT DOSE VIALS • NOT FOR INJECTION
FOR ORAL INHALATION USE ONLY

PLACE YOUR PRESCRIPTION LABEL HERE

0.5 mg/2.5 mL UNIT DOSE VIALS

Ipratropium Bromide Inhalation Solution
0.02% (0.5 mg/2.5 mL)
Rx Only

2000

FOR ORAL INHALATION USE ONLY

Each Low Density Polyethylene Vial Contains: 2.5 mL Ipratropium Bromide 0.02% preservative-free, sterile, isotonic aqueous solution containing sodium chloride. Adjusted to pH 3.4 (3 to 4) with hydrochloric acid.

Attention Pharmacist Detach "Patient's Instructions for Use" from package insert and dispense with solution. See package insert for Dosage and Administration.

Store between 15° and 30°C (59° and 86°F). Protect from light.

Retain in foil pouch until time of use.

5 x 2.5 mL Sterile Unit Dose Vials

Manufactured for:

Zenith Goldline Pharmaceuticals Inc., MIAMI, FL 33137

By: Steripak Ltd., Runcorn, Cheshire WA7 1QF England

1199J

PP1232

LOT :

EXP :



0046.