




**TAMIFLU™**  
**(oseltamivir phosphate)**  
**CAPSULES**

**DESCRIPTION:** TAMIFLU (oseltamivir phosphate) is available as a capsule containing 75 mg oseltamivir for oral use, in the form of oseltamivir phosphate. In addition to the active ingredient, each capsule contains pregelatinized starch, talc, povidone K 30, croscarmellose sodium, sodium stearyl fumarate, ethanol, and purified water. The capsule shell contains gelatin, titanium dioxide, yellow iron oxide, black iron oxide, and red iron oxide. Each capsule is printed with blue ink, which includes FD&C Blue #2 as the colorant. Oseltamivir phosphate is a white crystalline solid with the chemical name (3R,4R,5S)-4-acetylamino-5-amino-3(1-ethylpropoxy)-1-cyclohexene-1-carboxylic acid, ethyl ester, phosphate (1:1). The chemical formula is  $C_{16}H_{28}N_2O_4$  (free base). The molecular weight is 312.4 for oseltamivir free base and 410.4 for oseltamivir phosphate salt. The structural formula is as follows:



**MICROBIOLOGY: Mechanism of Action:** Oseltamivir is an ethyl ester prodrug requiring ester hydrolysis for conversion to the active form, oseltamivir carboxylate. The proposed mechanism of action of oseltamivir is via inhibition of influenza virus neuraminidase with the possibility of alteration of virus particle aggregation and release.

**Antiviral Activity In Vitro:** The antiviral activity of oseltamivir carboxylate against laboratory strains and clinical isolates of influenza virus was determined in cell culture assays. The concentrations of oseltamivir carboxylate required for inhibition of influenza virus were highly variable depending on the assay method used and the virus tested. The 50% and 90% inhibitory concentrations (IC<sub>50</sub> and IC<sub>90</sub>) were in the range of 0.0008  $\mu$ M to >35  $\mu$ M and 0.004  $\mu$ M to >100  $\mu$ M, respectively (1  $\mu$ M=0.284  $\mu$ g/mL). The relationship between the in vitro antiviral activity in cell culture and the inhibition of influenza virus replication in humans has not been established.

**Drug Resistance:** Influenza A virus isolates with reduced susceptibility to oseltamivir carboxylate have been recovered in vitro by passage of virus in the presence of increasing concentrations of oseltamivir carboxylate. Genetic analysis of these isolates showed that reduced susceptibility to oseltamivir carboxylate is associated with mutations that result in amino acid changes in the viral neuraminidase or viral hemagglutinin or both.

In challenge studies in the treatment of human subjects infected with influenza virus, 3% (3/102) of the post-treatment isolates showed emergence of influenza variants with decreased neuraminidase susceptibility to oseltamivir carboxylate. Genotypic analysis of these variants showed a specific mutation in the active site of neuraminidase compared to challenge virus.

In clinical studies of post-exposure and seasonal prophylaxis, determination of resistance was limited by the low overall incidence rate of influenza infection and prophylactic effect of TAMIFLU.

In clinical studies in the treatment of naturally acquired infection with influenza virus, 1.3% (4/301) of post-treatment isolates showed emergence of influenza variants with decreased neuraminidase susceptibility to oseltamivir carboxylate.

Genotypic analysis of these variants showed a specific mutation in the active site of neuraminidase compared to pretreatment isolates. The contribution of resistance due to alterations in the viral hemagglutinin has not been fully evaluated.

**Cross-resistance:** Cross-resistance between zanamivir-resistant influenza mutants and oseltamivir-resistant influenza mutants has been observed in vitro.

Due to limitations in the assays available to detect drug-induced shifts in virus susceptibility, an estimate of the incidence of oseltamivir resistance and possible cross-resistance to zanamivir in clinical isolates cannot be made. However, one of the three oseltamivir-induced mutations in the viral neuraminidase from clinical isolates is the same as one of the three mutations observed in zanamivir-resistant virus. Insufficient information is available to fully characterize the risk of emergence of TAMIFLU resistance in clinical use.

**Immune Response:** No influenza vaccine interaction study has been conducted. In studies of naturally acquired and experimental influenza, treatment with TAMIFLU did not impair normal humoral antibody response to infection.

**Influenza Challenge Studies:** Antiviral activity of TAMIFLU was supported for influenza A and B by experimental challenge studies in volunteers who received intranasal inoculations of challenge strains of influenza virus. These subjects received TAMIFLU either 24 hours following or 24 hours before virus challenge.

#### **CLINICAL PHARMACOLOGY: PHARMACOKINETICS:**

**Absorption and Bioavailability:** Oseltamivir is readily absorbed from the gastrointestinal tract after oral administration of oseltamivir phosphate and is extensively converted predominantly by hepatic esterases to oseltamivir carboxylate. At least 75% of an oral dose reaches the systemic circulation as oseltamivir carboxylate. Exposure to oseltamivir is less than 5% of the total exposure after oral dosing (Table 1).

**Table 1. Mean (% CV) Pharmacokinetic Parameters of Oseltamivir and Oseltamivir Carboxylate After a Multiple 75 mg Twice Daily Oral Dose (n=20)**

<b>Parameter</b>	<b>Oseltamivir</b>	<b>Oseltamivir Carboxylate</b>
$C_{\max}$ (ng/mL)	65.2 (26)	348 (18)
$AUC_{0-12h}$ (ng·h/mL)	112 (25)	2719 (20)

Plasma concentrations of oseltamivir carboxylate are proportional to doses up to 500 mg given twice daily (see DOSAGE AND ADMINISTRATION).

Coadministration with food has no significant effect on the peak plasma concentration (551 ng/mL under fasted conditions and 441 ng/mL under fed conditions) and the area under the plasma concentration time curve (6218 ng·h/mL under fasted conditions and 6069 ng·h/mL under fed conditions) of oseltamivir carboxylate.

**Distribution:** The volume of distribution ( $V_{ss}$ ) of oseltamivir carboxylate, following intravenous administration in 24 subjects, ranged between 23 and 26 liters.

The binding of oseltamivir carboxylate to human plasma protein is low (3%). The binding of oseltamivir to human plasma protein is 42%, which is insufficient to cause significant displacement-based drug interactions.

**Metabolism:** Oseltamivir is extensively converted to oseltamivir carboxylate by esterases located predominantly in the liver. Neither oseltamivir nor oseltamivir carboxylate is a substrate for, or inhibitor of, cytochrome P450 isoforms.

**Elimination:** Absorbed oseltamivir is primarily (>90%) eliminated by conversion to oseltamivir carboxylate. Plasma concentrations of oseltamivir declined with a half-life of 1 to 3 hours in most subjects after oral administration. Oseltamivir carboxylate is not further metabolized and is eliminated in the urine. Plasma concentrations of oseltamivir carboxylate declined with a half-life of 6 to 10 hours in most subjects after oral administration. Oseltamivir carboxylate is eliminated entirely (>99%) by renal excretion. Renal clearance (18.8 L/h) exceeds glomerular filtration rate (7.5 L/h) indicating that tubular secretion occurs, in addition to glomerular filtration. Less than 20% of an oral radiolabeled dose is eliminated in feces.

**Special Populations: Renal Impairment:** Administration of 100 mg of oseltamivir phosphate twice daily for 5 days to patients with various degrees of renal impairment showed that exposure to oseltamivir carboxylate is inversely proportional to declining renal function. Oseltamivir carboxylate exposures in patients with normal and abnormal renal function administered various dose regimens of oseltamivir are described in Table 2.

**Table 2. Oseltamivir Carboxylate Exposures in Patients With Normal and Reduced Serum Creatinine Clearance**

Parameter	Normal Renal Function			Impaired Renal Function			
	75 mg qd	75 mg bid	150 mg bid	Creatinine Clearance <10 mL/min		Creatinine Clearance >10 and <30 mL/min	
				CAPD	Hemodialysis		
				75 mg weekly	75 mg alternate HD cycle	75 mg daily	75 mg alternate days
C <sub>max</sub>	259*	348*	705*	1885*	2131*	1638	1175
C <sub>min</sub>	39*	138*	288*	155	120	864	209
AUC <sub>0-48</sub>	7476*	10876*	21864*	43451	31073	62636	21999

\*Observed values. All other values are predicted.

**Geriatric Patients:** Exposure to oseltamivir carboxylate at steady-state was 25% to 35% higher in geriatric patients (age range 65 to 78 years) compared to young adults given comparable doses of oseltamivir. Half-lives observed in the geriatric patients were similar to those seen in young adults. Based on drug exposure and tolerability, dose adjustments are not required for geriatric patients for either treatment or prophylaxis (see DOSAGE AND ADMINISTRATION: *Special Dosage Instructions*).

## INDICATIONS AND USAGE:

**Treatment of Influenza:** TAMIFLU is indicated for the treatment of uncomplicated acute illness due to influenza infection in adults who have been symptomatic for no more than 2 days. This indication is based on studies of naturally occurring influenza in which the predominant infection was influenza A, and influenza challenge studies in which antiviral activity of TAMIFLU was supported for influenza A and B (see *Description of Clinical Studies* and PRECAUTIONS).

**Prophylaxis of Influenza:** TAMIFLU is indicated for the prophylaxis of influenza in adults and

adolescents 13 years and older.

TAMIFLU is not a substitute for early vaccination on an annual basis as recommended by the Centers for Disease Control's Immunization Practices Advisory Committee.

***Description of Clinical Studies: Studies in Naturally Occurring Influenza:***

**Treatment of Influenza: Adults:** Two phase 3 placebo-controlled and double-blind clinical trials were conducted: one in the USA and one outside the USA. Patients were eligible for these trials if they had fever >100°F, accompanied by at least one respiratory symptom (cough, nasal symptoms or sore throat) and at least one systemic symptom (myalgia, chills/sweats, malaise, fatigue or headache) and influenza virus was known to be circulating in the community. In addition, all patients enrolled in the trials were allowed to take fever-reducing medications.

Of 1355 patients enrolled in these two trials, 849 (63%) patients were influenza-infected (age range 18 to 65 years; median age 34 years; 52% male; 90% Caucasian; 31% smokers). Of the 849 influenza infected patients, 95% were infected with influenza A, 3% with influenza B, and 2% with influenza of unknown type.

TAMIFLU was started within 40 hours of onset of symptoms. Subjects participating in the trials were required to self-assess the influenza-associated symptoms as “none,” “mild,” “moderate” or “severe”. Time to improvement was calculated from the time of treatment initiation to the time when all symptoms (nasal congestion, sore throat, cough, aches, fatigue, headaches, and chills/sweats) were assessed as “none” or “mild”. In both studies, at the recommended dose of TAMIFLU 75 mg twice daily for 5 days, there was a 1.3 day reduction in the median time to improvement in influenza-infected subjects receiving TAMIFLU compared to subjects receiving placebo. Subgroup analyses of these studies by gender showed no differences in the treatment effect of TAMIFLU in men and women.

In the treatment of influenza, no increased efficacy was demonstrated in subjects receiving treatment of 150 mg TAMIFLU twice daily for 5 days.

**Prophylaxis of Influenza:** The efficacy of TAMIFLU in preventing naturally occurring influenza illness has been demonstrated in three seasonal prophylaxis studies and a post-exposure prophylaxis study in households. The primary efficacy parameter for all these studies was the incidence of laboratory confirmed clinical influenza. Laboratory confirmed clinical influenza was defined as oral temperature • 99.0°F/37.2°C plus at least one respiratory symptom (cough, sore throat, nasal congestion) and at least one constitutional symptom (aches and pain, fatigue, headache, chills/sweats), all recorded within 24 hours, plus either a positive virus isolation or a fourfold increase in virus antibody titers from baseline.

In a pooled analysis of two seasonal prophylaxis studies in healthy unvaccinated adults (aged 13 to 65 years), TAMIFLU 75 mg once daily taken for 42 days during a community outbreak reduced the incidence of laboratory confirmed clinical influenza from 4.8% (25/519) for the placebo group to 1.2% (6/520) for the TAMIFLU group.

In a seasonal prophylaxis study in elderly residents of skilled nursing homes, TAMIFLU 75 mg once daily taken for 42 days reduced the incidence of laboratory confirmed clinical influenza from 4.4% (12/272) for the placebo group to 0.4% (1/276) for the TAMIFLU group. About 80% of this elderly population were vaccinated, 14% of subjects had chronic airway obstructive disorders, and 43% had cardiac disorders.

In a study of post-exposure prophylaxis in household contacts (aged • 13 years) of an index case, TAMIFLU 75 mg once daily administered within 2 days of onset of symptoms in the index case and continued for 7 days reduced the incidence of laboratory confirmed clinical influenza from 12% (24/200) in the placebo group to 1% (2/205) for the TAMIFLU group. Index cases did not receive TAMIFLU in the study.

**CONTRAINDICATIONS:** TAMIFLU is contraindicated in patients with known hypersensitivity to any of the components of the product.

**PRECAUTIONS: General:** There is no evidence for efficacy of TAMIFLU in any illness caused by agents other than influenza viruses Types A and B. Data on treatment of influenza B are limited (see INDICATIONS AND USAGE: *Description of Clinical Studies*).

Use of TAMIFLU should not affect the evaluation of individuals for annual influenza vaccination in accordance with guidelines of the Center for Disease Controls and Prevention Advisory Committee on Immunization Practices.

Efficacy of TAMIFLU in patients who begin treatment after 40 hours of symptoms has not been established.

Efficacy of TAMIFLU in the treatment of subjects with chronic cardiac disease and/or respiratory disease has not been established. No difference in the incidence of complications was observed between the treatment and placebo groups in this population. No information is available regarding treatment of influenza in patients with any medical condition sufficiently severe or unstable to be considered at imminent risk of requiring hospitalization.

Safety and efficacy of repeated treatment or prophylaxis courses have not been studied.

Efficacy of TAMIFLU for treatment or prophylaxis has not been established in immunocompromised patients.

Serious bacterial infections may begin with influenza-like symptoms or may coexist with or occur as complications during the course of influenza. TAMIFLU has not been shown to prevent such complications.

**Hepatic Impairment:** The safety and pharmacokinetics in patients with hepatic impairment have not been evaluated.

**Renal Impairment:** Dose adjustment is recommended for patients with a serum creatinine clearance <30 mL/min (see DOSAGE AND ADMINISTRATION).

**Information for Patients:** Patients should be instructed to begin treatment with TAMIFLU as soon as possible from the first appearance of flu symptoms. Similarly, prevention should begin as soon as possible after exposure, at the recommendation of a physician.

Patients should be instructed to take any missed doses as soon as they remember, except if it is near the next scheduled dose (within 2 hours), and then continue to take TAMIFLU at the usual times.

TAMIFLU is not a substitute for a flu vaccination. Patients should continue receiving an annual flu vaccination according to guidelines on immunization practices.

**Drug Interactions:** Information derived from pharmacology and pharmacokinetic studies of oseltamivir suggests that clinically significant drug interactions are unlikely.

Oseltamivir is extensively converted to oseltamivir carboxylate by esterases, located predominantly in the liver. Drug interactions involving competition for esterases have not been extensively reported in literature. Low protein binding of oseltamivir and oseltamivir carboxylate suggests that the probability of drug displacement interactions is low.

In vitro studies demonstrate that neither oseltamivir nor oseltamivir carboxylate is a good substrate for P450 mixed-function oxidases or for glucuronyl transferases.

Cimetidine, a non-specific inhibitor of cytochrome P450 isoforms and competitor for renal tubular secretion of basic or cationic drugs, has no effect on plasma levels of oseltamivir or oseltamivir carboxylate.

Clinically important drug interactions involving competition for renal tubular secretion are unlikely due to the known safety margin for most of these drugs, the elimination characteristics of oseltamivir carboxylate (glomerular filtration and anionic tubular secretion) and the excretion capacity of these pathways. Coadministration of probenecid results in an approximate twofold increase in exposure to oseltamivir carboxylate due to a decrease in active anionic tubular secretion in the kidney. However, due to the safety margin of oseltamivir carboxylate, no dose adjustments are required when coadministering with probenecid.

Coadministration with amoxicillin does not alter plasma levels of either compound, indicating that competition for the anionic secretion pathway is weak.

In six subjects, multiple doses of oseltamivir did not affect the single-dose pharmacokinetics of acetaminophen.

**Carcinogenesis, Mutagenesis, and Impairment of Fertility:** Long-term carcinogenicity tests with oseltamivir are underway but have not been completed. However, a 26-week dermal carcinogenicity study of oseltamivir carboxylate in FVB/Tg.AC transgenic mice was negative. The animals were dosed at 40, 140, 400 or 780 mg/kg/day in two divided doses. The highest dose represents the maximum feasible dose based on the solubility of the compound in the control vehicle. A positive control, tetradecanoyl phorbol-13-acetate administered at 2.5 µg per dose three times per week gave a positive response.

Oseltamivir was found to be non-mutagenic in the Ames test and the human lymphocyte chromosome assay with and without enzymatic activation and negative in the mouse micronucleus test. It was found to be positive in a Syrian Hamster Embryo (SHE) cell transformation test. Oseltamivir carboxylate was non-mutagenic in the Ames test and the L5178Y mouse lymphoma assay with and without enzymatic activation and negative in the SHE cell transformation test.

In a fertility and early embryonic development study in rats, doses of oseltamivir at 50, 250, and 1500 mg/kg/day were administered to females for 2 weeks before mating, during mating and until day 6 of pregnancy. Males were dosed for 4 weeks before mating, during and for 2 weeks after mating. There were no effects on fertility, mating performance or early embryonic development at any dose level. The highest dose was approximately 100 times the human systemic exposure ( $AUC_{0-24h}$ ) of oseltamivir carboxylate.

Long-term carcinogenicity tests with oseltamivir have not been completed.

**Pregnancy:** Pregnancy Category C: There are insufficient human data upon which to base an evaluation of risk of TAMIFLU to the pregnant woman or developing fetus. Studies for effects on embryo-fetal development were conducted in rats (50, 250, and 1500 mg/kg/day) and rabbits (50, 150, and 500 mg/kg/day) by the oral route. Relative exposures at these doses were, respectively, 2, 13, and 100 times human exposure in the rat and 4, 8, and 50 times human exposure in the rabbit. Pharmacokinetic studies indicated that fetal exposure was seen in both species. In the rat study, minimal maternal toxicity was reported in the 1500 mg/kg/day group. In the rabbit study, slight and marked maternal toxicities were observed, respectively, in the 150 and 500 mg/kg/day groups. There was a dose-dependent increase in the incidence rates of a variety of minor skeletal abnormalities and variants in the exposed offspring in these studies. However, the individual incidence rate of each skeletal abnormality or variant remained within the background rates of occurrence in the species studied.

Because animal reproductive studies may not be predictive of human response and there are no adequate and well-controlled studies in pregnant women, TAMIFLU should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

**Nursing Mothers:** In lactating rats, oseltamivir and oseltamivir carboxylate are excreted in the milk. It

is not known whether oseltamivir or oseltamivir carboxylate is excreted in human milk. TAMIFLU should, therefore, be used only if the potential benefit for the lactating mother justifies the potential risk to the breast-fed infant.

**Pediatric Use:** The safety and efficacy of TAMIFLU in children (<18 years) have not been established.

**Geriatric Use:** In an ongoing treatment study in otherwise healthy elderly patients, >65 years (n=168), given the recommended dosing regimen of TAMIFLU, there was a reduction in the median time to improvement in the subjects receiving TAMIFLU similar to that seen in younger adults. No overall difference in safety was observed between these subjects and younger adults. Safety and efficacy have been demonstrated in elderly residents of nursing homes who took TAMIFLU for up to 42 days for the prevention of influenza. Many of these individuals had cardiac and/or respiratory disease, and most had received vaccine that season (see *Description of Clinical Studies*).

**ADVERSE REACTIONS: Treatment Studies:** A total of 1171 patients who participated in adult phase 3 controlled clinical trials for the treatment of influenza were treated with TAMIFLU. The most frequently reported adverse events in these studies were nausea and vomiting. These events were generally of mild to moderate degree and usually occurred on the first 2 days of administration. Less than 1% of subjects discontinued prematurely from clinical trials due to nausea and vomiting.

Adverse events that occurred with an incidence of  $\geq 1\%$  in 1440 patients taking placebo or TAMIFLU 75 mg twice daily in adult phase 3 treatment studies are shown in Table 3. This summary includes 945 healthy young adults and 495 “at risk” patients (elderly patients and patients with chronic cardiac or respiratory disease). Those events reported numerically more frequently in patients taking TAMIFLU compared with placebo were nausea, vomiting, bronchitis, insomnia, and vertigo.

**Table 3. Most Frequent Adverse Events in Studies in Naturally Acquired Influenza**

Adverse Event	Treatment		Prophylaxis	
	Placebo N=716	Oseltamivir 75 mg bid N=724	Placebo N=1434	Oseltamivir 75 mg qd N=1480
Nausea (without vomiting)	40 (5.6%)	72 (9.9%)	56 (3.9%)	104 (7.0%)
Vomiting	21 (2.9%)	68 (9.4%)	15 (1.0%)	31 (2.1%)
Diarrhea	70 (9.8%)	48 (6.6%)	38 (2.6%)	48 (3.2%)
Bronchitis	15 (2.1%)	17 (2.3%)	17 (1.2%)	11 (0.7%)
Abdominal pain	16 (2.2%)	16 (2.2%)	23 (1.6%)	30 (2.0%)
Dizziness	25 (3.5%)	15 (2.1%)	21 (1.5%)	24 (1.6%)
Headache	14 (2.0%)	13 (1.8%)	251 (17.5%)	298 (20.1%)
Cough	12 (1.7%)	9 (1.2%)	86 (6.0%)	83 (5.6%)
Insomnia	6 (0.8%)	8 (1.1%)	14 (1.0%)	18 (1.2%)
Vertigo	4 (0.6%)	7 (1.0%)	3 (0.2%)	4 (0.3%)
Fatigue	7 (1.0%)	7 (1.0%)	107 (7.5%)	117 (7.9%)

Adverse events included are: all events reported in the treatment studies with frequency  $\geq 1\%$  in the oseltamivir 75 mg bid group.

Additional adverse events occurring in <1% of patients receiving TAMIFLU for treatment included unstable angina, anemia, pseudomembranous colitis, humerus fracture, pneumonia, pyrexia, and peritonsillar abscess.

**Prophylaxis Studies:** A total of 3434 subjects (adolescents, healthy adults and elderly) participated in phase III prophylaxis studies, of whom 1480 received the recommended dose of 75 mg once daily for up to 6 weeks. Adverse events were qualitatively very similar to those seen in the treatment studies, despite a longer duration of dosing (Table 3). Events reported more frequently in subjects receiving

TAMIFLU compared to subjects receiving placebo in prophylaxis studies, and more commonly than in treatment studies, were aches and pains, rhinorrhea, dyspepsia and upper respiratory tract infections. However, the difference in incidence between TAMIFLU and placebo for these events was less than 1%. There were no clinically relevant differences in the safety profile of the 942 elderly subjects who received TAMIFLU or placebo, compared with the younger population.

**Observed During Clinical Practice for Treatment:** The following adverse reactions have been identified during post-marketing use of TAMIFLU. Because these reactions are reported voluntarily from a population of uncertain size, it is not possible to reliably estimate their frequency or establish a causal relationship to TAMIFLU exposure.

General: Rash, swelling of the face or tongue

Cardiac: Arrhythmia

Neurologic: Seizure, confusion

Metabolic: Aggravation of diabetes

**OVERDOSAGE:** At present, there has been no experience with overdose. Single doses of up to 1000 mg of TAMIFLU have been associated with nausea and/or vomiting. A complete pack of ten capsules of TAMIFLU contains a total of 750 mg of oseltamivir.

**DOSAGE AND ADMINISTRATION:** TAMIFLU may be taken with or without food (see *PHARMACOKINETICS*). However, when taken with food, tolerability may be enhanced in some patients.

**Standard Dosage – Treatment of Influenza:** The recommended oral dose of TAMIFLU for treatment of influenza is 75 mg twice daily for 5 days. Treatment should begin within 2 days of onset of symptoms of influenza.

**Standard Dosage – Prophylaxis of Influenza:** The recommended oral dose of TAMIFLU for prophylaxis of influenza following close contact with an infected individual is 75 mg once daily for at least 7 days. Therapy should begin within 2 days of exposure. The recommended dose for prophylaxis during a community outbreak of influenza is 75 mg once daily. Safety and efficacy have been demonstrated for up to 6 weeks. The duration of protection lasts for as long as dosing is continued.

**Special Dosage Instructions: Hepatic Impairment:** The safety and pharmacokinetics in patients with hepatic impairment have not been evaluated.

**Renal Impairment:** For plasma concentrations of oseltamivir carboxylate predicted to occur following various dosing schedules in patients with renal impairment, see *CLINICAL PHARMACOLOGY: PHARMACOKINETICS: Special Populations*.

**Treatment of Influenza:** Dose adjustment is recommended for patients with creatinine clearance between 10 and 30 mL/min receiving TAMIFLU for the treatment of influenza. In these patients it is recommended that the dose be reduced to 75 mg of TAMIFLU once daily for 5 days. No recommended dosing regimens are available for patients undergoing routine hemodialysis and continuous peritoneal dialysis treatment with end-stage renal disease.

**Prophylaxis of Influenza:** For the prophylaxis of influenza, dose adjustment is recommended for patients with creatinine clearance between 10 and 30 mL/min receiving TAMIFLU. In these patients it is recommended that the dose be reduced to 75 mg of TAMIFLU every other day. No recommended dosing regimens are available for patients undergoing routine hemodialysis and continuous peritoneal dialysis treatment with end-stage renal disease.

**Pediatric Patients:** The safety and efficacy of TAMIFLU in children have not been established.

**Geriatric Patients:** No dose adjustment is required for geriatric patients (see *PHARMACOKINETICS: Special Populations* and *PRECAUTIONS*).

**HOW SUPPLIED:** TAMIFLU is supplied as 75-mg (75 mg free base equivalent of the phosphate salt) grey/light yellow hard gelatin capsules. "ROCHE" is printed in blue ink on the grey body and "75 mg" is printed in blue ink on the light yellow cap. Available in blister packages of 10 (NDC 0004-0800-85).

**Storage:** Store at 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F). [See USP Controlled

Room Temperature]  
Manufactured by:

F. Hoffmann-La Roche Ltd.  
Basel, Switzerland

Distributed by:



**Pharmaceuticals**

Roche Laboratories Inc.  
340 Kingsland Street  
Nutley, New Jersey 07110-1199

Licensors:

Gilead Sciences, Inc.  
Foster City, California 94404

XXXXXXXX-XXXX

XXXXXXXX-XXXX

Revised: November 2000

Printed in USA

Copyright © 1999-2000 by Roche Laboratories Inc. All rights reserved.

## Patient Information

### TAMIFLU™ (oseltamivir phosphate)

This leaflet contains important information about TAMIFLU (TAM-ih-flew). Read it well before you begin treatment. This information does not take the place of talking with your health care professional about your medical condition or your treatment. This leaflet does not list all the benefits and risks of TAMIFLU. If you have any questions about TAMIFLU, ask your health care professional. Only your health care professional can determine if TAMIFLU is right for you.

#### What is TAMIFLU?

TAMIFLU attacks the influenza virus and stops it from spreading inside your body. TAMIFLU treats flu at its source, by attacking the virus that causes the flu, rather than simply masking symptoms.

TAMIFLU is for treating adults with the flu whose flu symptoms started within the last day or two. TAMIFLU can also reduce the chance of getting the flu in people age 13 and older who have a higher chance of getting the flu because they spend time with someone who has the flu. TAMIFLU can also reduce the chance of getting the flu if there is a flu outbreak in the community.

#### What is “Flu”?

“The flu” is an infection caused by the influenza virus. Flu symptoms include fever (usually 100°F to 103°F in adults, and sometimes higher in children) and problems such as cough, sore throat, runny or stuffy nose, headaches, muscle aches, fever, and extreme tiredness. Many people use the term “flu” to mean any combination of these symptoms, such as the common cold, but true influenza infection is often worse and may last longer than a cold.

Flu outbreaks happen about once a year, usually in the winter, when the influenza virus spreads widely in the community. Outside of those outbreaks, only a very tiny number of respiratory infections are caused by the influenza virus.

#### Should I get a flu shot?

TAMIFLU is not a substitute for a flu vaccination. You should continue to get a flu vaccination every year, according to your health care professional’s advice.

#### Who should not take TAMIFLU?

Do not take TAMIFLU if you are allergic to the main ingredient, oseltamivir phosphate, or to any other ingredients of TAMIFLU. Before starting treatment, make sure your health care professional knows if you take any other medicines, or are pregnant, planning to become pregnant, or breastfeeding. TAMIFLU is normally not recommended for use during pregnancy or nursing, as the effects on the unborn child or nursing infant are unknown.

Tell your health care professional if you have any type of kidney disease, heart disease, respiratory disease, or any serious health condition.

#### How should I take TAMIFLU?

It is important that you begin your treatment with TAMIFLU as soon as possible from the first appearance of your flu symptoms or soon after you are exposed to the flu. If you feel worse or develop new symptoms during treatment with TAMIFLU, or if your flu symptoms do not start to get better, you should contact your health care professional.

If you have the flu: Take TAMIFLU twice a day for 5 days, once in the morning and once in the evening. You should complete the entire treatment of 10 capsules, even if you feel better.

To prevent the flu: If someone in your home has the flu, take TAMIFLU once a day for at least 7 days or for as long as prescribed. You can take TAMIFLU for up to 6 weeks if you are exposed to the flu because of an outbreak in your community. Follow your health care professional's advice on how long to take TAMIFLU.

You can take TAMIFLU with food or without food. There is less chance of stomach upset if you take it with a light snack, milk, or a meal.

If you forget to take your medicine, take the missed dose as soon as you remember, except if it is 2 hours or less before your next dose. Then continue to take TAMIFLU at the usual times. Do not take 2 doses at a time to make up for a missed dose. If you miss several doses, tell your health care professional and follow the advice given to you.

### **What are the possible side effects of TAMIFLU?**

The most common side effects of TAMIFLU are nausea and vomiting. These are usually mild to moderate. They usually happen in the first 2 days of treatment. Taking TAMIFLU with food may reduce the chance of getting these side effects.

If you notice any side effects not mentioned in this leaflet, or if you have any concerns about the side effects you get, tell your health care professional.

### **How and where should I store TAMIFLU?**

TAMIFLU capsules should be stored at room temperature below 77°F (25°C) and kept in a dry place. Keep this medication out of reach of children.

### **General advice about prescription medicines:**

Medicines are sometimes prescribed for conditions that are not mentioned in patient information leaflets. Do not use TAMIFLU for a condition for which it was not prescribed. Do not give TAMIFLU to other people, even if they have the same symptoms you have. It may not be right for them.

This leaflet summarizes the most important information about TAMIFLU. If you would like more information, talk with your health care professional. You can ask your pharmacist or health care professional for information about TAMIFLU that is written.

/s/

-----  
Debra Birnkrant  
11/17/00 03:33:50 PM  
NDA 21-087/S-002