

Cures Athlete's Foot

Relieves Itching & Burning  
Greaseless, Nonstaining

LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF

clotrimazole cream

LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF

Net Wt 12g (0.42oz)

Schering-Plough



LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF

NO  
CODE AREA  
NO VARNISH

NO  
CODE AREA  
NO VARNISH

LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF  
NO INK  
CODE AREA  
NO VARNISH

LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF  
NO INK  
CODE AREA  
NO VARNISH

FOR BEST RESULTS FOLLOW DIRECTIONS AND CONTINUE TREATMENT FOR LENGTH OF TIME INDICATED.

INDICATIONS: LOTTRIMIN AF CREAM cures athlete's foot (tinea pedis), jock itch (tinea cruris) and ringworm (tinea corporis). For effective relief of the itching, cracking, burning and discomfort which can accompany these conditions, athlete's foot and ringworm are usually obtained with 4 weeks use of this product, and in jock itch, with 2 weeks use. It is helpful to wear well-fitting, ventilated shoes and socks at least once daily. Best results in athlete's foot and ringworm are usually obtained with 4 weeks use of this product, and in jock itch, with 2 weeks use. If satisfactory results have not occurred within these times, consult a doctor or pharmacist. Children under 12 years of age should be supervised in the use of this product. This product is not effective on the scalp or nails.

WARNINGS: For external use only. Avoid contact with the eyes. Do not use on children under 2 years of age except under the advice and supervision of a doctor. If irritation occurs or if there is no improvement within 4 weeks (for athlete's foot or ringworm) or within 2 weeks (for jock itch), discontinue use and consult a doctor or pharmacist. Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately.

Use only as directed.

ACTIVE INGREDIENT: Clotrimazole 1%. INACTIVE INGREDIENTS: Cetylalcohol, benzyl alcohol (1%), cetyl esters wax, oxydodecanol, polyorbate, sorbitan monostearate, water and, as a preservative, benzyl alcohol (1%).

© 1989, 1997, Distributed by Schering-Plough HealthCare Products, Inc., Kenilworth, NJ 07033

Made in USA. All rights reserved.

16558630

- Cures athlete's foot
- Relieves itching & burning

LOTTRIMIN<sup>®</sup>  
CREAM ANTIFUNGAL AF

6321

**ACTIVE INGREDIENT:** Clotrimazole 1%. **Store between 2° and 30°C (36° and 86°F).**  
© 1989, 1997, Distributed by Schering-Plough Healthcare Products, Inc., Memphis, TN 38151 USA.  
All rights reserved. Control No. and Exp. Date—see crimp.  
**TO OPEN: USE POINTED END ON CAP TO PUNCTURE SEAL. FOR BEST RESULTS, FOLLOW DIRECTIONS AND CONTINUE TREATMENT FOR LENGTH OF TIME INDICATED.**

Clotrimazole cream  
**LOTTRIMIN<sup>AF</sup>**  
CREAM ANTIFUNGAL  
Net Wt 12g (0.42oz)

**INDICATIONS:** LOTTRIMIN AF CREAM cures athlete's foot (tinea pedis), jock itch (tinea cruris) and ringworm (tinea corporis). For effective relief of the itching, cracking, burning and discomfort which can accompany these conditions. **DIRECTIONS:** Cleanse skin with soap and water and dry thoroughly. Apply a thin layer over affected area morning and evening or as directed by a doctor. For athlete's foot, pay special attention to the spaces between the toes. It is also helpful to wear well-fitting, ventilated shoes and to change shoes and socks at least once daily. Best results in athlete's foot and ringworm are usually obtained with 4 weeks use of this product, and in jock itch, with 2 weeks use. If satisfactory results have not occurred within these times, consult a doctor or pharmacist. Children under 12 years of age should be supervised in the use of this product. This product is not effective on the scalp or nails. **WARNINGS:** For external use only. Avoid contact with the eyes. Do not use on children under 2 years of age except under the advice and supervision of a doctor. If irritation occurs or if there is no improvement within 4 weeks (for athlete's foot or ringworm) or within 2 weeks (for jock itch), discontinue use and consult a doctor or pharmacist. Keep this and all drugs out of the reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately. Use only as directed.

