

Carton Label, Drug Facts, Back Panel

Chlorhexidine Gluconate 2% w/v and Isopropyl Alcohol 70% v/v  
Patient Preoperative Skin Preparation • 0.67 mL Sepp® Applicator  
**WARNING. FLAMMABLE. KEEP AWAY FROM FIRE OR FLAME.**  
**DO NOT USE WITH ELECTROCAUTERY PROCEDURES.**

---

## Drug Facts

---

### Active ingredients

	<i>Purpose</i>
Chlorhexidine gluconate 2% w/v.	..... Antiseptic
Isopropyl alcohol 70% v/v .....	..... Antiseptic

---

Use for the preparation of the patient's skin prior to surgery or injection. The maximum treatment area for one applicator is approximately 2.5 by 2.5 inches.

---

### Warnings

**Flammable**, keep away from fire or flame  
Do not use with electrocautery procedures  
**For external use only**

---

### Do not use

- in children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

---

**When using this product** keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a physician.

---

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

---

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

---

### Directions

- pinch the barrel of the applicator to break the ampule and release the antiseptic. Do not touch the applicator tip. Press the applicator tip against the treatment area until liquid is visible on the skin.
- **dry surgical sites** (such as abdomen or arm): Use repeated back-and-forth strokes of the applicator for approximately 30 seconds. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately 30 seconds. Do not blot or wipe away.
- **moist surgical sites** (such as the inguinal fold): Use repeated back-and-forth strokes of the applicator for approximately 2 minutes. Completely wet the treatment area with antiseptic. Allow the area to air dry for approximately one (1) minute. Do not blot or wipe away.
- discard the applicator after a single use.

---

### Other information

- store between 20-25 °C (68-77 °F)
- avoid freezing and excessive heat above 40 °C (104 °F)

---

**Inactive ingredients** USP purified water

---

**Questions?** Call 1-800-523-0502 (8 a.m.-5 p.m. CST)

---

B. Individual Lidding Applicator Label

ChloroPrep®  
**Sepp**  
Chlorhexidine  
Gluconate 2% w/v;  
70% Isopropyl  
Alcohol v/v  
0.67-mL Applicator

Warning.  
Flammable.  
Keep away from  
fire or flame.  
Do not use with  
electrocautery  
procedures.  
Applicator sterile  
unless seal is  
broken.

Medi-Flex, Inc.  
Overland Park, KS

Lot No:  
Exp.:

-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Janice Soreth  
10/4/02 03:59:38 PM

Charles Ganley  
10/7/02 11:03:01 AM