

**CENTER FOR DRUG EVALUATION AND
RESEARCH**

APPLICATION NUMBER:

75803

DRAFT FINAL PRINTED LABELING

APPROVED MAR 20 2002

**Lessina™
21 Tablets**

(levonorgestrel and ethinyl estradiol tablets, USP)

31590130101

Revised MAY 2001

**COMBINATION DETAILED PATIENT
LABELING AND BRIEF SUMMARY**



DETAILED PATIENT LABELING

Rx only

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

INTRODUCTION:

Any woman who considers using oral contraceptives (the "birth-control pill" or "the pill") should understand the benefits and risks of using this form of birth control. This leaflet will give you much of the information you will need to make this decision and will also help you determine if you are at risk of developing any of the serious side effects of the pill. It will tell you how to use the pill properly so that it will be as effective as possible. However, this leaflet is not a replacement for a careful discussion between you and your healthcare provider. You should discuss the information provided in this leaflet with him or her, both when you first start taking the pill and during your revisits. You should also follow your healthcare provider's advice with regard to regular check-ups while you are on the pill.

EFFECTIVENESS OF ORAL CONTRACEPTIVES:

Oral contraceptives or "birth-control pills" or "the pill" are used to prevent pregnancy and are more effective than other nonsurgical methods of birth control. When they are taken correctly, the chance of becoming pregnant is less than 1.0% when used perfectly, without missing any pills. Typical failure rates are less than 3.0% per year. The chance of becoming pregnant increases with each missed pill during a menstrual cycle. In comparison, typical failure rates for other nonsurgical methods of birth control during the first year of use are as follows:

Percentage of women experiencing an unintended pregnancy during the first year of typical use and first year of perfect use of contraception and the percentage continuing use at the end of the first year, United States.

Method (1)	% of Women Experiencing an Unintended Pregnancy within the First Year of Use Typical Use ¹ (2)	% of Women Experiencing an Unintended Pregnancy within the First Year of Use Perfect Use ² (3)	% of Women Continuing Use at One Year ³ (4)
Chance ⁴	85	85	
Spermicides ⁵	26	6	40
Periodic abstinence	25		63
Calendar		9	
Ovulation method		3	
Sympto-therm ⁶		2	
Post ovulation		2	
Withdrawal	19	4	
Cap ⁷			
Parous women	40	26	42
Nulliparous women	20	9	56
Sponge			
Parous women	40	20	42
Nulliparous women	20	9	56
Diaphragm ⁸	20	6	56
Condom ⁹			
Female (Reality)	21	5	56
Male	14	3	61
Pill	5		71
progestin only		0.5	
combined		0.1	
IUD			
Progesterone T	2.0	1.5	81
Copper T380A	0.8	0.6	78
Leq 20	0.1	0.1	81
Depo Provera	0.3	0.3	70
Norplant and Norplant-2	0.05	0.05	88
Female sterilization	0.5	0.5	100
Male sterilization	0.15	0.10	100

Source: Trussell J. Contraceptive efficacy. In Hatcher RA, Trussell J, Stewart F, Cates W, Stewart GK, Kowal D, Guest F, *Contraceptive Technology: Seventeenth Revised Edition*. New York NY: Irvington Publishers, 1998, in press.

- Among typical couples who initiate use of a method (not necessarily for the first time), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason.
- Among couples who initiate use of a method (not necessarily for the first time) and who use it perfectly (both consistently and correctly), the percentage who experience an accidental pregnancy during the first year if they do not stop use for any other reason.
- Among couples attempting to avoid pregnancy, the percentage who continue to use a method for one year.
- The percentages becoming pregnant in columns (2) and (3) are based on data from populations where contraception is not used and from women who cease using contraception in order to become pregnant. Among such populations, about 89% become pregnant within one year. This estimate was lowered slightly (to 85%) to represent the percentage who would become pregnant within one year among women now relying on reversible methods of contraception if they abandoned contraception altogether.
- Foams, creams, gels, vaginal suppositories, vaginal film.
- Cervical mucus (ovulation) method supplemented by calendar in the pre-ovulatory and basal body temperature in the post-ovulatory phases.
- With spermicidal cream or jelly.
- Without spermicides.

WHO SHOULD NOT TAKE ORAL CONTRACEPTIVES:

Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels from oral contraceptive use. This risk increases with age and with heavy smoking (15 or more cigarettes per day) and is quite marked in women over 35 years of age. Women who use oral contraceptives should not smoke.

Some women should not use the pill. For example, you should not take the pill if you are pregnant or think you may be pregnant. You should also not use the pill if you have had any of the following conditions:

- A history of heart attack or stroke
 - Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), or eyes
 - A history of blood clots in the deep veins of your legs
 - Chest pain (angina pectoris)
 - Known or suspected breast cancer or cancer of the lining of the uterus, cervix or vagina
 - Unexplained vaginal bleeding (until a diagnosis is reached by your doctor)
 - Yellowing of the whites of the eyes or of the skin (jaundice) during pregnancy or during previous use of the pill
 - Liver tumor (benign or cancerous)
 - Known or suspected pregnancy
- Tell your healthcare provider if you have ever had any of these conditions. Your healthcare provider can recommend another method of birth control.

OTHER CONSIDERATIONS BEFORE TAKING ORAL CONTRACEPTIVES:

- Tell your healthcare provider if you or any family member has ever had:
- Breast nodules, fibrocystic disease of the breast, an abnormal breast X-ray or mammogram
 - Diabetes
 - Elevated cholesterol or triglycerides
 - High blood pressure
 - Migraine or other headaches or epilepsy
 - Mental depression
 - Gallbladder, heart or kidney disease
 - History of scanty or irregular menstrual periods
- Women with any of these conditions should be checked often by their healthcare provider if they choose to use oral contraceptives. Also, be sure to inform your doctor or healthcare provider if you smoke or are on any medications.

RISKS OF TAKING ORAL CONTRACEPTIVES:

- Risk of developing blood clots:**
Blood clots and blockage of blood vessels are the most serious side effects of taking oral contraceptives and can be fatal. In particular, a clot in the legs can cause thrombophlebitis and a clot that travels to the lungs can cause sudden blockage of the vessel carrying blood to the lungs. Rarely, clots occur in the blood vessels of the eye and may cause blindness, double vision, or impaired vision. If you take oral contraceptives and need elective surgery, need to stay in bed for a prolonged illness or have recently delivered a baby, you may be at risk of developing blood clots. You should consult your doctor about stopping oral contraceptives three to four weeks before surgery and not taking oral contraceptives for two weeks after surgery or during bed rest. You should also not take oral contraceptives soon after delivery of a baby or a mid-trimester pregnancy termination. It is advisable to wait for at least four weeks after delivery if you are not breast-feeding. If you are breast-feeding, you should wait until you have weaned your child before using the pill. (See also the section on breast-feeding in "GENERAL PRECAUTIONS".)
- Heart attacks and strokes:**
Oral contraceptives may increase the tendency to develop strokes (stoppage or rupture of blood vessels in the brain) and angina pectoris and heart attacks (blockage of blood vessels in the heart). Any of these conditions can cause death or serious disability. Smoking greatly increases the possibility of suffering heart attacks and strokes. Furthermore, smoking and the use of oral contraceptives greatly increase the chances of developing and dying of heart disease.
- Gallbladder disease:**
Oral contraceptive users probably have a greater risk than nonusers of having gallbladder disease, although this risk may be related to pills containing high doses of estrogens.
- Liver tumors:**
In rare cases, oral contraceptives can cause benign but dangerous liver tumors. These benign liver tumors can rupture and cause fatal inter-

nal bleeding. In addition, a possible but not definite association has been found with the pill and liver cancers in two studies, in which a few women who developed these very rare cancers were found to have used oral contraceptives for long periods. However, liver cancers are extremely rare. The chance of developing liver cancer from using the pill is thus even rarer.

5. Cancer of the reproductive organs:

There is, at present, no confirmed evidence that oral contraceptives increase the risk of cancer of the reproductive organs in human studies. Several studies have found no overall increase in the risk of developing breast cancer. However, women who use oral contraceptives and have a strong family history of breast cancer or who have breast nodules or abnormal mammograms should be closely followed by their doctors. Some studies have found an increase in the incidence of cancer of the cervix in women who use oral contraceptives. However, this finding may be related to factors other than the use of oral contraceptives.

TABLE V: ANNUAL NUMBER OF BIRTH-RELATED OR METHOD-RELATED DEATHS ASSOCIATED WITH CONTROL OF FERTILITY PER 100,000 NON-STERILE WOMEN, BY FERTILITY-CONTROL METHOD ACCORDING TO AGE

Method of Control and Outcome	15-19	20-24	25-29	30-34	35-39	40-44
No fertility control methods ⁽¹⁾	7.0	7.4	9.1	14.8	25.7	28.2
Oral contraceptives non-smoker ⁽²⁾	0.3	0.5	0.9	1.9	13.8	31.6
Oral contraceptives smoker ⁽²⁾	2.2	3.4	6.6	13.5	51.1	117.2
IUD ⁽²⁾	0.8	0.8	1.0	1.0	1.4	1.4
Condom ⁽¹⁾	1.1	1.6	0.7	0.2	0.3	0.4
Diaphragm/ ⁽¹⁾ spermicide ⁽¹⁾	1.9	1.2	1.2	1.3	2.2	2.8
Periodic abstinence ⁽¹⁾	2.5	1.6	1.6	1.7	2.9	3.6

Adapted from H.W. Ory, *Family Planning Perspectives* 15:57-63, 1983.

In the above table, the risk of death from any birth-control method is less than the risk of childbirth, except for oral contraceptive users over the age of 35 who smoke and pill users over the age of 40 even if they do not smoke. It can be seen in the table that for women aged 15 to 39, the risk of death was highest with pregnancy (7-26 deaths per 100,000 women, depending on age). Among pill users who do not smoke, the risk of death was always lower than that associated with pregnancy for any age group, except for those women over the age of 40, when the risk increases to 32 deaths per 100,000 women, compared to 28 associated with pregnancy at that age. However, for pill users who smoke and are over the age of 35, the estimated number of deaths exceeds those for other methods of birth control. If a woman is over the age of 40 and smokes, her estimated risk of death is four times higher (117/100,000 women) than the estimated risk associated with pregnancy (28/100,000 women) in that age group.

The suggestion that women over 40 who do not smoke should not take oral contraceptives is based on information from older high-dose pills and on less-selective use of pills than is practiced today. An Advisory Committee of the FDA discussed this issue in 1989 and recommended that the benefits of oral contraceptive use by healthy, non-smoking women over 40 years of age may outweigh the possible risks. However, all women, especially older women, are cautioned to use the lowest-dose pill that is effective.

WARNING SIGNALS:

- If any of these adverse effects occur while you are taking oral contraceptives, call your doctor immediately:
- Sharp chest pain, coughing of blood, or sudden shortness of breath (indicating a possible clot in the lung)
 - Pain in the calf (indicating a possible clot in the leg)
 - Crushing chest pain or heaviness in the chest (indicating a possible heart attack)
 - Sudden severe headache or vomiting, dizziness or fainting, disturbances of vision or speech, weakness, or numbness in an arm or leg (indicating a possible stroke)
 - Sudden partial or complete loss of vision (indicating a possible clot in the eye)
 - Breast lumps (indicating possible breast cancer or fibrocystic disease of the breast; ask your doctor or healthcare provider to show you how to examine your breasts)
 - Severe pain or tenderness in the stomach area (indicating a possible ruptured liver tumor)
 - Difficulty in sleeping, weakness, lack of energy, fatigue, or change in mood (possibly indicating severe depression)
 - Jaundice or a yellowing of the skin or eyeballs, accompanied frequently by fever, fatigue, loss of appetite, dark-colored urine, or light-colored bowel movements (indicating possible liver problems)

SIDE EFFECTS OF ORAL CONTRACEPTIVES:

- Vaginal bleeding:**
Irregular vaginal bleeding or spotting may occur while you are taking the pills. Irregular bleeding may vary from slight staining between menstrual periods, or breakthrough bleeding, which is a flow much like a regular period. Irregular bleeding occurs most often during the first few months of oral contraceptive use, but may also occur after you have been taking the pill for some time. Such bleeding may be temporary and usually does not indicate any serious problems. It is important to continue taking your pills on schedule. If the bleeding occurs in more than one cycle or lasts for more than a few days, talk to your doctor or healthcare provider.

2. Contact lenses:

If you wear contact lenses and notice a change in vision or an inability to wear your lenses, contact your doctor or healthcare provider.

3. Fluid retention:

Oral contraceptives may cause edema (fluid retention) with swelling of the fingers or ankles and may raise your blood pressure. If you experience fluid retention, contact your doctor or healthcare provider.

4. Melasma:

A spotty darkening of the skin is possible, particularly of the face.

5. Other side effects:

Other side effects may include change in appetite, headache, nervousness, depression, dizziness, loss of scalp hair, rash, and vaginal infections.

If any of these side effects bother you, call your doctor or healthcare provider.

GENERAL PRECAUTIONS:

- Missed periods and use of oral contraceptives before or during early pregnancy:**
There may be times when you may not menstruate regularly after you have completed taking a cycle of pills. If you have taken your pills regularly and miss one menstrual period, continue taking your pills for the next cycle but be sure to inform your healthcare provider before doing so. If you have not taken the pills daily as instructed and missed a menstrual period, or if you missed two consecutive menstrual periods, you may be pregnant. Check with your healthcare provider immediately to determine whether you are pregnant. Do not continue to take oral contraceptives until you are sure you are not pregnant, but continue to use another method of contraception. There is no conclusive evidence that oral contraceptive use is associated with an increase in birth defects when taken inadvertently during early pregnancy. Previously, a few studies had reported that oral contraceptives might be associated with birth defects, but these studies have not been confirmed. Nevertheless, oral contraceptives or any other drugs should not be used during pregnancy unless clearly necessary and prescribed by your doctor. You should check with your doctor about risks to your unborn child of any medication taken during pregnancy.
- While breast-feeding:**
If you are breast-feeding, consult your doctor before starting oral contraceptives. Some of the drug will be passed on to the child in the milk. A few adverse effects on the child have been reported, including yellowing of the skin (jaundice) and breast enlargement. In addition, oral contraceptives may decrease the amount and quality of your milk. If possible, do not use oral contraceptives while breast-feeding. You should use another method of contraception since breast-feeding provides only partial protection from becoming pregnant, and this partial protection decreases significantly as you breast-feed for longer periods of time. You should consider starting oral contraceptives only after you have weaned your child completely.

3. Laboratory tests:

If you are scheduled for any laboratory tests, tell your doctor you are taking birth-control pills. Certain blood tests may be affected by birth-control pills.

4. Drug Interactions:

Certain drugs may interact with birth-control pills to make them less effective in preventing pregnancy or cause an increase in breakthrough bleeding. Such drugs include rifampin, drugs used for epilepsy such as barbiturates (for example, phenobarbital) and phenytoin (Dilantin is one brand of this drug), phenylbutazone (Butazolidin is one brand) and possibly certain antibiotics. You may need to use an additional method of contraception during any cycle in which you take drugs that can make oral contraceptives less effective.

5. Sexually transmitted diseases:

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against transmission of HIV (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

HOW TO TAKE THE PILL

IMPORTANT POINTS TO REMEMBER

BEFORE YOU START TAKING YOUR PILLS:

- BE SURE TO READ THESE DIRECTIONS:**
Before you start taking your pills.
Anytime you are not sure what to do.
- THE RIGHT WAY TO TAKE THE PILL IS TO TAKE ONE PILL EVERY DAY AT THE SAME TIME.**
If you miss pills you could get pregnant. This includes starting the pack late. The more pills you miss, the more likely you are to get pregnant.
- MANY WOMEN HAVE SPOTTING OR LIGHT BLEEDING, OR MAY FEEL SICK TO THEIR STOMACH DURING THE FIRST 1-3 PACKS OF PILLS.**
If you do feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it does not go away, check with your doctor or clinic.
- MISSING PILLS CAN ALSO CAUSE SPOTTING OR LIGHT BLEEDING.** even when you make up these missed pills. On the days you take two pills, to make up for missed pills, you could also feel a little sick to your stomach.
- IF YOU HAVE VOMITING OR DIARRHEA, for any reason, or IF YOU TAKE SOME MEDICINES,** including some antibiotics, your pills may not work as well.
Use a back-up method (such as condoms, foam, or sponge) until you check with your doctor or clinic.
- IF YOU HAVE TROUBLE REMEMBERING TO TAKE THE PILL,** talk to your doctor or clinic about how to make pill-taking easier or about using another method of birth control.

(over)

← Tear here at perforation.

Lessina™

(levonorgestrel and ethinyl estradiol tablets, USP)
21 Tablets

BRIEF SUMMARY PATIENT PACKAGE INSERT

Rx only

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases.

21 tablets containing the following:

21 pink "active" tablets

Oral contraceptives, also known as "birth-control pills" or "the pill", are taken to prevent pregnancy, and when taken correctly, have a failure rate of less than 1% per year when used without missing any pills. The typical failure rate of large numbers of pill users is less than 3% per year when women who miss pills are included. For most women oral contraceptives are also free of serious or unpleasant side effects. However, forgetting to take pills considerably increases the chances of pregnancy.

For the majority of women, oral contraceptives can be taken safely. But there are some women who are at high risk of developing certain serious diseases that can be life-threatening or may cause temporary or permanent disability or death. The risks associated with taking oral contraceptives increase significantly if you:

- smoke
- have high blood pressure, diabetes, high cholesterol
- have or have had clotting disorders, heart attack, stroke, angina pectoris, cancer of the breast or sex organs, jaundice, or malignant or benign liver tumors.

You should not take the pill if you suspect you are pregnant or have unexplained vaginal bleeding.

Cigarette smoking increases the risk of serious adverse effects on the heart and blood vessels from oral contraceptive use. This risk increases with age and with heavy smoking (15 or more cigarettes per day) and is quite marked in women over 35 years of age. Women who use oral contraceptives should not smoke.

Most side effects of the pill are not serious. The most common such effects are nausea, vomiting, bleeding between menstrual periods, weight gain, breast tenderness, and difficulty wearing contact lenses. These side effects, especially nausea and vomiting may subside within the first three months of use.

The serious side effects of the pill occur very infrequently, especially if you are in good health and do not smoke. However, you should know that the following medical conditions have been associated with or made worse by the pill.

1. Blood clots in the legs (thrombophlebitis), lungs (pulmonary embolism), stoppage or rupture of a blood vessel in the brain (stroke), blockage of blood vessels in the heart (heart attack and angina pectoris) or other organs of the body. As mentioned above, smoking increases the risk of heart attacks and strokes and subsequent serious medical consequences.

2. Liver tumors, which may rupture and cause severe bleeding. A possible but not definite association has been found with the pill and liver cancer. However, liver cancers are extremely rare. The chance of developing liver cancer from using the pill is thus even rarer.

3. High blood pressure, although blood pressure usually returns to normal when the pill is stopped.

The symptoms associated with these serious side effects are discussed in the detailed leaflet given to you with your supply of pills. Notify your doctor or healthcare provider if you notice any unusual physical disturbances while taking the pill. In addition, drugs such as rifampin, as well as some anticonvulsants and some antibiotics, may decrease oral contraceptive effectiveness.

Studies to date of women taking the pill have not shown an increase in the incidence of cancer of the breast or cervix. There is, however, insufficient evidence to rule out the possibility that pills may cause such cancers.

Taking the pill provides some important non-contraceptive benefits. These include less painful menstruation, less menstrual blood loss and anemia, fewer pelvic infections, and fewer cancers of the ovary and the lining of the uterus.


Be sure to discuss any medical condition you may have with your healthcare provider. Your healthcare provider will take a medical and family history before prescribing oral contraceptives and will examine you. The physical examination may be delayed to another time if you request it and the healthcare provider believes that it is appropriate to postpone it. You should be re-examined at least once a year while taking oral contraceptives. The Detailed Patient Information Leaflet gives you further information which you should read and discuss with your healthcare provider.

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhea, hepatitis B, and syphilis.

INSTRUCTIONS TO PATIENTS

How to Use the Lessina™ 21 Tablets Blister Card:

1. Pick the Days of the Week Sticker that starts the first day of your period. (This is the day you begin bleeding or spotting, even if it is midnight when bleeding begins.) When you have picked the right sticker, throw away the others and place the sticker on the blister card over the pre-printed days of the week and make sure it lines up with the pills.
2. Your blister package consists of two parts, the sleeve and a blister card containing 21 individually sealed pills. Note that the pills are arranged in three numbered rows of 7 pills, with the pre-printed days of the week printed above them. All 21 pills are "active" birth control pills. Refer to the sample of the blister card below:



3. On the first day of your period, take the first pink pill in row #1. To remove a pill, push down on the pill with your thumb and forefinger so that the pill releases through the back of the blister card. Each day, take one pill. Always go from left to right along the row. Each new row will begin on the same day of the week.

If your physician has instructed you to use a "Sunday Start" method, then use the blister card which is set up for a Sunday start, and begin with the first pink pill on the Sunday after your period begins. If the first day of your bleeding falls on Sunday, begin taking the first pink pill that day. If you use a Sunday start, you must use back-up contraception for at least 7 days of pills.

4. Your period will usually begin about three days after you take the last pink pill. Don't be alarmed if the amount of bleeding is not the same as before. No matter when your period starts, wait 7 days after taking the last pink pill before you begin a new package of pills. For example, if you took the last pink pill on Tuesday, you would not take the first pink pill from a new package until Wednesday of the following week. The basic schedule is 21 days on the pills - seven days off the pills. Any time you start the pills later than directed, protect yourself by using another method of birth control until you have taken a pill a day for seven consecutive days (7 pink).

THREE WAYS TO REMEMBER IN WHAT ORDER TO TAKE THE PILLS.

1. Follow the sticker with the days of the week (placed above the pills).
2. Always go from left to right.
3. Always finish all your pink pills.
4. After 7 pill-free days have passed, start a new package of pills on the eighth day. Anytime you start the pills later than directed, protect yourself by using another method of birth control until you have taken 7 pink pills for 7 consecutive days. If you have taken the pills as directed, the first pink pill in every package will always be taken on the same day of the week that you first began taking Lessina™ 21 tablets.
5. Take the pills in each new package as before. Start with the pink pill at the left of the first row and take one pill each day left to right, until the last pink pill has been taken. Always wait 7 days before starting a new blister card.

HOW TO TAKE THE PILL

IMPORTANT POINTS TO REMEMBER

BEFORE YOU START TAKING YOUR PILLS:

1. **BE SURE TO READ THESE DIRECTIONS:**
Before you start taking your pills.
Anytime you are not sure what to do.
2. **THE RIGHT WAY TO TAKE THE PILL IS TO TAKE ONE PILL EVERY DAY AT THE SAME TIME.**
If you miss pills you could get pregnant. This includes starting the pack late. The more pills you miss, the more like you are to get pregnant.
3. **MANY WOMEN HAVE SPOTTING OR LIGHT BLEEDING, OR MAY FEEL SICK TO THEIR STOMACH DURING THE FIRST 1-3 PACKS OF PILLS.**
If you do feel sick to your stomach, do not stop taking the pill. The problem will usually go away. If it does not go away, check with your doctor or clinic.
4. **MISSING PILLS CAN ALSO CAUSE SPOTTING OR LIGHT BLEEDING,** even when you make up these missed pills.
On the days you take two pills, to make up for missed pills, you could also feel a little sick to your stomach.
5. **IF YOU HAVE VOMITING OR DIARRHEA,** for any reason, or **IF YOU TAKE SOME MEDICINES,** including some antiotics, your pills may not work as well. Use a back-up method (such as condoms, foam, or sponge) until you check with your doctor or clinic.

(over)

- IF YOU HAVE TROUBLE REMEMBERING TO TAKE THE PILL, talk to your doctor or clinic about how to make pill-taking easier or about using another method of birth control.
- IF YOU HAVE ANY QUESTIONS OR ARE UNSURE ABOUT THE INFORMATION IN THIS LEAFLET, call your doctor or clinic.

BEFORE YOU START TAKING YOUR PILLS

- DECIDE WHAT TIME OF DAY YOU WANT TO TAKE YOUR PILL.
It is important to take it at about the same time every day.
- LOOK AT YOUR PILL PACK TO SEE IF IT HAS 21 PILLS:
The 21-pill pack has 21 (pink) "active" pills (with hormones) to take for three weeks, followed by 1 week without pills.
- ALSO FIND:
 - where on the pack to start taking pills,
 - in what order to take the pills.
- BE SURE YOU HAVE READY AT ALL TIMES:
ANOTHER KIND OF BIRTH CONTROL (such as condoms, foam or sponge) to use as a back-up in case you miss pills.

AN EXTRA, FULL PILL PACK

WHEN TO START THE FIRST PACK OF PILLS

You have a choice for which day to start taking your first pack of pills. Decide with your doctor or clinic which is the best day for you. Pick a time of day which will be easy to remember.

DAY 1 START:

- Take the first (pink) "active" pill of the first pack during the first 24 hours of your period.
- You will not need to use a back-up method of birth control, since you are starting the pill at the beginning of your period.

SUNDAY START:

- Take the first (pink) "active" pill of the first pack on the Sunday after your period starts, even if you are still bleeding. If your period begins on Sunday, start the pack that same day.
- Use another method of birth control as a back-up method if you have sex anytime from the Sunday you start your first pack until the next Sunday (7 days). Condoms, foam, or the sponge are good back-up methods of birth control.

WHAT TO DO DURING THE MONTH

- TAKE ONE PILL AT THE SAME TIME EVERY DAY UNTIL THE PACK IS EMPTY**
Do not skip pills even if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea). Do not skip pills even if you do not have sex very often.
- WHEN YOU FINISH A PACK OR SWITCH YOUR BRAND OF PILLS:**
21 pills; Wait 7 days to start the next pack. You will probably have your period during that week. Be sure that no more than 7 days pass between 21-day packs.

WHAT TO DO IF YOU MISS PILLS

- If you MISS 1 (pink) "active" pill:
- Take it as soon as you remember. Take the next pill at your regular time. This means you may take two pills in one day.
 - You do not need to use a back-up birth control method if you have sex.

If you MISS 2 (pink) "active" pills in a row in WEEK 1 OR WEEK 2 of your pack:

- Take two pills on the day you remember and two pills the next day.
- Then take one pill a day until you finish the pack.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

If you MISS 2 (pink) "active" pills in a row in THE 3rd WEEK:

- If you are a Day 1 Starter:**
THROW OUT the rest of the pill pack and start a new pack that same day.
If you are a Sunday Starter:
Keep taking one pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.
- You may not have your period this month but this is expected. However, if you miss your period two months in a row, call your doctor or clinic because you might be pregnant.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

If you MISS 3 OR MORE (pink) "active" pills in a row (during the first 3 weeks):

- If you are a Day 1 Starter:**
THROW OUT the rest of the pill pack and start a new pack that same day.
If you are a Sunday Starter:
Keep taking one pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.
- You may not have your period this month but this is expected. However, if you miss your period two months in a row, call your doctor or clinic because you might be pregnant.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

FINALLY, IF YOU ARE STILL NOT SURE WHAT TO DO ABOUT THE PILLS YOU HAVE MISSED:

Use a BACK-UP METHOD anytime you have sex.
KEEP TAKING ONE ACTIVE PILL EACH DAY until you can reach your doctor or clinic.
For additional information see "Detailed Patient Labeling".

MANUFACTURED BY
BARR LABORATORIES, INC.
POMONA, NY 10970


Revised MAY 2001
BS-9013

- IF YOU HAVE ANY QUESTIONS OR ARE UNSURE ABOUT THE INFORMATION IN THIS LEAFLET, call your doctor or clinic.

BEFORE YOU START TAKING YOUR PILLS

- DECIDE WHAT TIME OF DAY YOU WANT TO TAKE YOUR PILL.
It is important to take it at about the same time every day.
- LOOK AT YOUR PILL PACK TO SEE IF IT HAS 21 OR 28 PILLS:
The 21-pill pack has 21 (pink) "active" pills (with hormones) to take for three weeks, followed by 1 week without pill:
The 28-pill pack has 21 (pink) "active" pills (with hormones) to take for three weeks, followed by 1 week of reminder (without hormones).

Refer to the sample of the blister card below:



- ALSO FIND:
 - where on the pack to start taking pills,
 - in what order to take the pills.
- BE SURE YOU HAVE READY AT ALL TIMES:
ANOTHER KIND OF BIRTH CONTROL (such as condoms, foam or sponge) to use as a back-up in case you miss pills. A FULL PILL PACK.

WHEN TO START THE FIRST PACK OF PILLS

You have a choice for which day to start taking your first pack of pills. Decide with your doctor or clinic which is the best day of day which will be easy to remember.

DAY 1 START:

- Take the first (pink) "active" pill of the first pack during the first 24 hours of your period.
- You will not need to use a back-up method of birth control, since you are starting the pill at the beginning of your period.

SUNDAY START:

- Take the first (pink) "active" pill of the first pack on the Sunday after your period starts, even if you are still bleeding, begins on Sunday, start the pack that same day.
- Use another method of birth control as a back-up method if you have sex anytime from the Sunday you start your first next Sunday (7 days). Condoms, foam, or the sponge are good back-up methods of birth control.

WHAT TO DO DURING THE MONTH

- TAKE ONE PILL AT THE SAME TIME EVERY DAY UNTIL THE PACK IS EMPTY**
Do not skip pills if you are spotting or bleeding between monthly periods or feel sick to your stomach (nausea). Do not skip pills even if you do not have sex very often.
- WHEN YOU FINISH A PACK OR SWITCH YOUR BRAND OF PILLS:**
21 pills; Wait 7 days to start the next pack. You will probably have your period during that week. Be sure that no more than 7 days pass between 21-day packs.
28 pills; Start the next pack on the day after your last (white) "reminder" pill. Do not wait any days between packs.

WHAT TO DO IF YOU MISS PILLS

- If you MISS 1 (pink) "active" pill:
- Take it as soon as you remember. Take the next pill at your regular time. This means you may take two pills in one day.
 - You do not need to use a back-up birth control method if you have sex.

If you MISS 2 (pink) "active" pills in a row in WEEK 1 OR WEEK 2 of your pack:

- Take two pills on the day you remember and two pills the next day.
- Then take one pill a day until you finish the pack.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

If you MISS 2 (pink) "active" pills in a row in THE 3rd WEEK:

- If you are a Day 1 Starter:**
THROW OUT the rest of the pill pack and start a new pack that same day.
If you are a Sunday Starter:
Keep taking one pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.
- You may not have your period this month but this is expected. However, if you miss your period two months in a row, call your doctor or clinic because you might be pregnant.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

If you MISS 3 OR MORE (pink) "active" pills in a row (during the first 3 weeks):

- If you are a Day 1 Starter:**
THROW OUT the rest of the pill pack and start a new pack that same day.
If you are a Sunday Starter:
Keep taking one pill every day until Sunday. On Sunday, THROW OUT the rest of the pack and start a new pack of pills that same day.
- You may not have your period this month but this is expected. However, if you miss your period two months in a row, call your doctor or clinic because you might be pregnant.
- You MAY BECOME PREGNANT if you have sex in the 7 days after you miss pills. You MUST use another birth control method (such as condoms, foam, or sponge) as a back-up for those 7 days.

A REMINDER FOR THOSE ON 28-DAY PACKS:

If you forget any of the 7 (white) "reminder" pills in Week 4:

THROW AWAY the pills you missed.
Keep taking one pill each day until the pack is empty.
You do not need a back-up method.

FINALLY, IF YOU ARE STILL NOT SURE WHAT TO DO ABOUT THE PILLS YOU HAVE MISSED:

Use a BACK-UP METHOD anytime you have sex.
KEEP TAKING ONE ACTIVE PILL EACH DAY until you can reach your doctor or clinic.

PREGNANCY DUE TO PILL FAILURE:

The incidence of pill failure resulting in pregnancy is approximately less than 1.0% if taken every day as directed, but more typical rates are less than 3.0%. If failure does occur, the risk to the fetus is minimal.

PREGNANCY AFTER STOPPING THE PILL:

There may be some delay in becoming pregnant after you stop using oral contraceptives, especially if you had irregular menstruation before you used oral contraceptives. It may be advisable to postpone conception until you begin menstruating regularly once you stopped taking the pill and desire pregnancy. There does not appear to be any increase in birth defects in newborn babies when pregnancy occurs soon after stopping the pill.

OVERDOSAGE:

Serious ill effects have not been reported following ingestion of large doses of oral contraceptives by young children. Overdosage cause nausea and withdrawal bleeding in females. In case of overdosage, contact your healthcare provider or pharmacist.

OTHER INFORMATION:

Your healthcare provider will take a medical and family history before prescribing oral contraceptives and will examine you. The examination may be delayed to another time if you request it and the healthcare provider believes that it is appropriate to postpone should be re-examined at least once a year. Be sure to inform your healthcare provider if there is a family history of any of the conditions listed previously in this leaflet. Be sure to keep all appointments with your healthcare provider, because this is a time to determine early signs of side effects of oral contraceptive use.
Do not use the drug for any condition other than the one for which it was prescribed. This drug has been prescribed specifically for you and do not give it to others who may want birth-control pills.

HEALTH BENEFITS FROM ORAL CONTRACEPTIVES:

In addition to preventing pregnancy, use of oral contraceptives may provide certain benefits. They are:

- Menstrual cycles may become more regular
- Blood flow during menstruation may be lighter and less iron may be lost. Therefore, anemia due to iron deficiency is less likely to occur.
- Pain or other symptoms during menstruation may be encountered less frequently
- Ovarian cysts may occur less frequently
- Ectopic (tubal) pregnancy may occur less frequently
- Noncancerous cysts or lumps in the breast may occur less frequently
- Acute pelvic inflammatory disease may occur less frequently
- Oral contraceptive use may provide some protection against developing two forms of cancer: cancer of the ovaries and cancer of the lining of the uterus.

If you want more information about birth-control pills, ask your doctor or pharmacist. They have a more technical leaflet called Prescribing Information which you may wish to read.

MANUFACTURED BY
BARR LABORATORIES, INC.
POMONA, NY 10970

Revised MAY 2001

PL-9013, 1

MAY 90

2050 Wardrobe Avenue • Merced, CA 95340 • 209.384.1000 • 800.388.1990 • FAX 209.384.0322 • graphics@labeltech.co

To better serve you PLEASE "check off" each of the following items as you complete your examination of this proof, then sign and fax/send back to u

Colors Copy Barcodes Other _____ Approved Approved w/ Changes Submit 2nd Proo

Customer: **Barr Laboratories**


Job No.: 62061 Date: 06.22.01 **Signed:**

Date:

Labeltechnology Proof

Please use this electronic proof for color breakdown and type placement. Refer to Pantone Matching System® for exact spot color match. Request a "dye-sub" proof for accurate 4-color process match.

Actual Size:
7.75" x 6.00"



FRONT PANEL



← Press Direction

• pink lines are for size reference only- will not print

48



Mango

SAMPLE

BARR LABORATORIES, INC.

NDC 0555-9013-57



Lessina™ 21

(levonorgestrel and ethinyl estradiol tablets, USP)

MAR 20 2002

APPROVED

BARR LABORATORIES, INC.

NDC 0555-9013-57



Lessina™ 21

(levonorgestrel and ethinyl estradiol tablets, USP)

Store at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F).

This product (like all oral contraceptives) is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted diseases.



Barr Laboratories, Inc.
Shaping Women's Health

52

Rx only

6 Blister Cards, 21 Tablets Each

BA
Po

PROOF

SCOTT & DANIELS, Inc. 264 PRESTON AVE. • PORTLAND, CT 06480
TEL. 860.342.1922 • FAX. 860.342.2436
W.O.# 9466

DD _____ Date: 6/28/01 Proof No. 1
Black _____ 2 PMS 293 3 PMS 325
PMS 190 _____ 5 _____ 6 _____


if carefully, for typographic as well as layout errors, then check
rite box, sign, and date below. Scott & Daniels, Inc. is not
e for errors or omissions found after final approval.

Print (final approval)
Print w/Corrections (final approval given in-house after
re are made and proofed)
ew Proof w/Corrections

_____ Date: _____
n-House By: _____ Date: _____

PRINTED SIDE


SCO-DAN
#BL60133
06/08/2001
0.016LDSBS
Barr Laboratories
001106
6 x 3+5/8 x 3+1/8
TT AUTO BTM
L=132 A=140



BARR LABORATORIES, INC.

NDC 0555-9013-5

Lessina™ 21



Lessina™ 21

(levonorgestrel and ethinyl estradiol tablets, USP)

(levonorgestrel and ethinyl estradiol tablets, USP)

Contains 6 blister cards, each containing 21 pink tablets. Each
tablet contains 0.1 mg levonorgestrel and 0.02 mg ethinyl estradiol.

Usual Dosage: One pink tablet daily for 21 consecutive days per
menstrual cycle as prescribed.

See enclosed package information.

To the Dispenser: This carton contains one piece of information intended
for the patient. All informational pieces are to be provided to the
patient with each prescription.

BARR LABORATORIES, INC.
10970

R5-01
2169013570103

6 Blister Cards, 21 Tablets Each

E

Lessina 21ct carton

■ Black

■ PMS 293

■ PMS 325

■ PMS 190

NDC 0555-9013-57

Lessina™ 21

Ethinyl Estradiol Tablets (SP)

containing 21 pink tablets. Each
tablet contains 0.02 mg ethinyl estradiol.
Take one tablet daily for 21 consecutive days per
cycle.

This leaflet contains one piece of information intended
to provide you with information. All other necessary
instructions are to be provided to the patient.

21 Tablets Each



KO Varnish



SCOTT &
DANIELLS, Inc.

PROOF

264 FRESTONE AVE. • PORTLAND, CT 06480
TEL: 860-342-1932 • FAX: 860-342-1933

NDC 0555-9013-57

aTM 21



21 pink tablets. Each
0.02 mg ethinyl estradiol.
consecutive days per



KO Varnish

piece of information intended
to be provided to the

ets Each



PROOF

SCOTT & DANIELS, Inc.

264 PRESTONE AVE. • PORTLAND, CT 06480
TEL: 860.342.1922 • FAX: 860.342.7436

W.O.# 9466

Prod. By: DD	Date: 6/28/01	Proof No. 1
Colors: 1. Black	2. PMS 293	3. PMS 325
4. PMS 190	5.	6.

Revisions

Please proof carefully, for typographic as well as layout errors, then check the appropriate box, sign, and date below. Scott & Daniels, Inc. is not responsible for errors or omissions found after final approval.

- O.K. to Print (final approval)
- O.K. to Print w/Corrections (final approval given in-house after corrections are made and proofed)
- Send New Proof w/Corrections

Cust. Sign: _____ Date: _____

Received In-House By: _____ Date: _____