CENTER FOR DRUG EVALUATION AND RESEARCH

Approved Labeling for:

APPLICATION NUMBER:

50-755/010

Trade Name: Augmentin ES-600™

Generic Name: Amoxicillin/clavulanate potassium

Sponsor: GlaxoSmithKline

Approval Date: June 3, 2004

AE:L9

PRESCRIBING INFORMATION

AUGMENTIN ES-600®

(amoxicillin/clavulanate potassium)
Powder for Oral Suspension

To reduce the development of drug-resistant bacteria and maintain the effectiveness of AUGMENTIN ES-600 (amoxicillin/clavulanate potassium) and other antibacterial drugs, AUGMENTIN ES-600 should be used only to treat or prevent infections that are proven or strongly suspected to be caused by bacteria.

DESCRIPTION

AUGMENTIN ES-600 is an oral antibacterial combination consisting of the semisynthetic antibiotic amoxicillin and the β-lactamase inhibitor, clavulanate potassium (the potassium salt of clavulanic acid). Amoxicillin is an analog of ampicillin, derived from the basic penicillin nucleus, 6-aminopenicillanic acid. The amoxicillin molecular formula is $C_{16}H_{19}N_3O_5S{\bullet}3H_2O$, and the molecular weight is 419.46. Chemically, amoxicillin is (2*S*,5*R*,6*R*)-6-[(*R*)-(-)-2-Amino-2-(*p*-hydroxyphenyl)acetamido]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylic acid trihydrate and may be represented structurally as:



Clavulanic acid is produced by the fermentation of *Streptomyces clavuligerus*. It is a β-lactam structurally related to the penicillins and possesses the ability to inactivate a wide variety of β-lactamases by blocking the active sites of these enzymes. Clavulanic acid is particularly active against the clinically important plasmid-mediated β-lactamases frequently responsible for transferred drug resistance to penicillins and cephalosporins. The clavulanate potassium molecular formula is $C_8H_8KNO_5$ and the molecular weight is 237.25. Chemically, clavulanate potassium is potassium (*Z*)-(2*R*,5*R*)-3-(2-hydroxyethylidene)-7-oxo-4-oxa-1-azabicyclo[3.2.0]-heptane-2-carboxylate and may be represented structurally as:



Inactive Ingredients: Powder for Oral Suspension—Colloidal silicon dioxide, orange flavorings, xanthan gum, aspartame•, hypromellose, and silicon dioxide.

• See PRECAUTIONS–Information for the Patient/Phenylketonurics.

Each 5 mL of reconstituted 600 mg/5 mL oral suspension of AUGMENTIN ES-600 contains 0.23 mEq potassium.

CLINICAL PHARMACOLOGY

The pharmacokinetics of amoxicillin and clavulanate were determined in a study of 19 pediatric patients, 8 months to 11 years, given AUGMENTIN ES-600 at an amoxicillin dose of 45 mg/kg q12h with a snack or meal. The mean plasma amoxicillin and clavulanate pharmacokinetic parameter values are listed in the following table.

Table 1. Mean (±SD) Plasma Amoxicillin and Clavulanate Pharmacokinetic Parameter Values Following Administration of 45 mg/kg of AUGMENTIN ES-600 Every 12 Hours to Pediatric Patients

Parameter*	Amoxicillin	Clavulanate
C_{max} (mcg/mL)	15.7 ± 7.7	1.7 ± 0.9
T_{max} (hr)	2.0 (1.0 – 4.0)	1.1 (1.0 – 4.0)
AUC_{0-t} (mcg•hr/mL)	59.8 ± 20.0	4.0 ± 1.9
T½ (hr)	1.4 ± 0.3	1.1 ± 0.3
CL/F (L/hr/kg)	0.9 ± 0.4	1.1 ± 1.1

*Arithmetic mean ± standard deviation, except T_{max} values which are medians (ranges).

The effect of food on the oral absorption of AUGMENTIN ES-600 has not been studied.

Approximately 50% to 70% of the amoxicillin and approximately 25% to 40% of the clavulanic acid are excreted unchanged in urine during the first 6 hours after administration of 10 mL of 250 mg/5 mL suspension of AUGMENTIN.

Concurrent administration of probenecid delays amoxicillin excretion but does not delay renal excretion of clavulanic acid.

Neither component in AUGMENTIN ES-600 is highly protein-bound; clavulanic acid has been found to be approximately 25% bound to human serum and amoxicillin approximately 18% bound.

Oral administration of a single dose of AUGMENTIN ES-600 at 45 mg/kg (based on the amoxicillin component) to pediatric patients, 9 months to 8 years, yielded the following pharmacokinetic data for amoxicillin in plasma and middle ear fluid (MEF):

Table 2. Amoxicillin Concentrations in Plasma and Middle Ear Fluid Following Administration of 45 mg/kg of AUGMENTIN ES-600 to Pediatric Patients

Timepoint		Amoxicillin concentration in plasma (mcg/mL)	Amoxicillin concentration in MEF (mcg/mL)
1 hour	mean	7.7	3.2
	median	9.3	3.5
	range	1.5 – 14.0	0.2 – 5.5
		(n = 5)	(n = 4)
2 hour	mean	15.7	3.3
	median	13.0	2.4
	range	11.0 – 25.0	1.9 – 6
		(n = 7)	(n = 5)
3 hour	mean	13.0	5.8
	median	12.0	6.5
	range	5.5 – 21.0	3.9 – 7.4
		(n = 5)	(n = 5)

Dose administered immediately prior to eating.

Amoxicillin diffuses readily into most body tissues and fluids with the exception of the brain and spinal fluid. The results of experiments involving the administration of clavulanic acid to animals suggest that this compound, like amoxicillin, is well distributed in body tissues.

Microbiology: Amoxicillin is a semisynthetic antibiotic with a broad spectrum of bactericidal activity against many gram-positive and gram-negative microorganisms. Amoxicillin is, however, susceptible to degradation by β-lactamases, and therefore, the spectrum of activity does not include organisms which produce these enzymes. Clavulanic acid is a β-lactam, structurally related to the penicillins, which possesses the ability to inactivate a wide range of β-lactamase enzymes commonly found in microorganisms resistant to penicillins and cephalosporins. In particular, it has good activity against the clinically important plasmid-mediated β-lactamases frequently responsible for transferred drug resistance.

The clavulanic acid component in AUGMENTIN ES-600 protects amoxicillin from degradation by β-lactamase enzymes and effectively extends the antibiotic spectrum of amoxicillin to include many bacteria normally resistant to amoxicillin and other β-lactam antibiotics. Thus, AUGMENTIN ES-600 possesses the distinctive properties of a broad-spectrum antibiotic and a β-lactamase inhibitor.

Amoxicillin/clavulanic acid has been shown to be active against most strains of the following microorganisms, both in vitro and in clinical infections as described in INDICATIONS AND USAGE.

Aerobic Gram-Positive Microorganisms:

Streptococcus pneumoniae (including isolates with penicillin MICs ≤2 mcg/mL)

Aerobic Gram-Negative Microorganisms:

Haemophilus influenzae (including β-lactamase–producing strains)

Moraxella catarrhalis (including β-lactamase–producing strains)

The following in vitro data are available, but their clinical significance is unknown.

At least 90% of the following microorganisms exhibit an in vitro minimum inhibitory concentration (MIC) less than or equal to the susceptible breakpoint for amoxicillin/clavulanic acid. However, with the exception of organisms shown to respond to amoxicillin alone, the safety and effectiveness of amoxicillin/clavulanic acid in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials.

Aerobic Gram-Positive Microorganisms:

Staphylococcus aureus (including β-lactamase–producing strains)

Streptococcus pyogenes

NOTE: Staphylococci which are resistant to methicillin/oxacillin must be considered resistant to amoxicillin/clavulanic acid.

NOTE: *S. pyogenes* do not produce β-lactamase, and therefore, are susceptible to amoxicillin alone. Adequate and well-controlled clinical trials have established the effectiveness of amoxicillin alone in treating certain clinical infections due to *S. pyogenes*.

Susceptibility Testing: *Dilution Techniques:* Quantitative methods are used to determine antimicrobial minimum inhibitory concentrations (MICs). These MICs provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MICs should be determined using a standardized procedure.[1,2] Standardized procedures are based on a dilution method (broth for *S. pneumoniae* and *H. influenzae)* or equivalent with standardized inoculum concentrations and standardized concentrations of amoxicillin/clavulanate potassium powder.

The recommended dilution pattern utilizes a constant amoxicillin/clavulanate potassium ratio of 2 to 1 in all tubes with varying amounts of amoxicillin. MICs are expressed in terms of the amoxicillin concentration in the presence of clavulanic acid at a constant 2 parts amoxicillin to 1 part clavulanic acid. The MIC values should be interpreted according to the following criteria:

For testing *Streptococcus pneumoniae*[a]:

MIC (mcg/mL)	Interpretation	
≤2/1	Susceptible	(S)
4/2	Intermediate	(I)
≥8/4	Resistant	(R)

[a] These interpretive standards are applicable only to broth microdilution susceptibility tests using cation-adjusted Mueller-Hinton broth with 2-5% lysed horse blood.[2]

For testing *Haemophilus influenzae*[b]:

MIC (mcg/mL)	Interpretation	
≤4/2	Susceptible	(S)
≥8/4	Resistant	(R)

[b] These interpretive standards are applicable only to broth microdilution susceptibility tests with *Haemophilus* spp. using *Haemophilus* Test Medium (HTM).[2]

A report of "Susceptible" indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentration usually achievable. A report of "Intermediate" indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone that prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of "Resistant" indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard amoxicillin/clavulanate potassium powder should provide the following MIC values:

Microorganism	MIC Range (mcg/mL)[c]
Escherichia coli ATCC 35218 (*H. influenzae* quality control)	4 to 16
Haemophilus influenzae[d] ATCC 49247	2 to 16
Streptococcus pneumoniae[e] ATCC 49619	0.03 to 0.12

[c] Expressed as concentration of amoxicillin in the presence of clavulanic acid at a constant 2 parts amoxicillin to 1 part clavulanic acid.

[d] This quality control range is applicable to *H. influenzae* ATCC 49247 tested by a broth microdilution procedure using HTM.[2]

[e] This quality control range is applicable to *S. pneumoniae* ATCC 49619 tested by a broth microdilution procedure using cation-adjusted Mueller-Hinton broth with 2-5% lysed horse blood.[2]

Diffusion Techniques: Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure[3] requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 30 mcg of amoxicillin/clavulanate potassium (20 mcg amoxicillin plus 10 mcg clavulanate potassium) to test the susceptibility of microorganisms to amoxicillin/clavulanic acid.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 30-mcg amoxicillin/clavulanate potassium (20 mcg amoxicillin plus 10 mcg clavulanate potassium) disk should be interpreted according to the following criteria:

For *H. influenzae*[f]:

Zone Diameter (mm)	Interpretation
≥20	Susceptible (S)
≤19	Resistant (R)

[f] These zone diameter standards are applicable only to tests conducted with *Haemophilus* spp. using HTM.[2]

NOTE: Beta-lactamase–negative, ampicillin-resistant *H. influenzae* strains must be considered resistant to amoxicillin/clavulanic acid.

For *Streptococcus pneumoniae:*

Susceptibility of *S. pneumoniae* should be determined using a 1-mcg oxacillin disk. Isolates with oxacillin zone sizes of ≥20 mm are susceptible to amoxicillin/clavulanic acid.[g] An amoxicillin/clavulanic acid MIC should be determined on isolates of *S. pneumoniae* with oxacillin zone sizes of ≤19 mm.

[g] These zone diameter standards for *S. pneumoniae* apply only to tests performed using Mueller-Hinton agar supplemented with 5% sheep blood incubated in 5% CO_2.[3]

Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for amoxicillin/clavulanic acid.

As with standardized dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 30-mcg amoxicillin/clavulanate potassium (20 mcg amoxicillin plus 10 mcg clavulanate potassium) disk should provide the following zone diameters in these laboratory quality control strains:

Microorganism	Zone Diameter (mm)
*Escherichia coli*ATCC 35218 (*H. influenzae* quality control)	18 to 22
Haemophilus influenzae[h] ATCC 49247	15 to 23

[h] This quality control limit applies only to tests conducted with *H. influenzae* ATCC 49247 using HTM.

INDICATIONS AND USAGE

AUGMENTIN ES-600 is indicated for the treatment of pediatric patients with recurrent or persistent acute otitis media due to *S. pneumoniae* (penicillin MICs ≤2 mcg/mL), *H. influenzae* (including β-lactamase–producing strains), or *M. catarrhalis* (including β-lactamase–producing strains) characterized by the following risk factors:

- antibiotic exposure for acute otitis media within the preceding 3 months, and either of the following:

– age ≤2 years

– daycare attendance

[See CLINICAL PHARMACOLOGY, Microbiology.]

Note: Acute otitis media due to *S. pneumoniae* alone can be treated with amoxicillin. AUGMENTIN ES-600 is not indicated for the treatment of acute otitis media due to *S. pneumoniae* with penicillin MIC ≥4 mcg /mL.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of AUGMENTIN ES-600 and other antibacterial drugs, AUGMENTIN ES-600 should be used only to treat or prevent infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy when there is reason to believe the infection may involve both *S. pneumoniae* (penicillin MIC ≤2 mcg/mL) and the β-lactamase–producing organisms listed above. Once the results are known, therapy should be adjusted appropriately.

CONTRAINDICATIONS

AUGMENTIN ES-600 is contraindicated in patients with a history of allergic reactions to any penicillin. It is also contraindicated in patients with a previous history of cholestatic jaundice/hepatic dysfunction associated with AUGMENTIN.

WARNINGS

SERIOUS AND OCCASIONALLY FATAL HYPERSENSITIVITY (ANAPHYLACTIC) REACTIONS HAVE BEEN REPORTED IN PATIENTS ON PENICILLIN THERAPY. THESE REACTIONS ARE MORE LIKELY TO OCCUR IN INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY AND/OR A HISTORY OF SENSITIVITY TO MULTIPLE ALLERGENS. THERE HAVE BEEN REPORTS OF INDIVIDUALS WITH A HISTORY OF PENICILLIN HYPERSENSITIVITY WHO HAVE EXPERIENCED SEVERE REACTIONS WHEN TREATED WITH CEPHALOSPORINS. BEFORE INITIATING THERAPY WITH AUGMENTIN ES-600, CAREFUL INQUIRY SHOULD BE MADE CONCERNING PREVIOUS HYPERSENSITIVITY REACTIONS TO PENICILLINS, CEPHALOSPORINS, OR OTHER ALLERGENS. IF AN ALLERGIC REACTION OCCURS, AUGMENTIN ES-600 SHOULD BE DISCONTINUED AND THE APPROPRIATE THERAPY INSTITUTED. **SERIOUS ANAPHYLACTIC REACTIONS REQUIRE IMMEDIATE EMERGENCY TREATMENT WITH EPINEPHRINE. OXYGEN, INTRAVENOUS STEROIDS, AND AIRWAY MANAGEMENT, INCLUDING INTUBATION, SHOULD ALSO BE ADMINISTERED AS INDICATED.**

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including amoxicillin/clavulanate potassium, and has ranged in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of "antibiotic-associated colitis."

After the diagnosis of pseudomembranous colitis has been established, appropriate therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

AUGMENTIN ES-600 should be used with caution in patients with evidence of hepatic dysfunction. Hepatic toxicity associated with the use of amoxicillin/clavulanate potassium is usually reversible. On rare occasions, deaths have been reported (less than 1 death reported per estimated 4 million prescriptions worldwide). These have generally been cases associated with serious underlying diseases or concomitant medications. (See CONTRAINDICATIONS and ADVERSE REACTIONS—Liver.)

PRECAUTIONS

General: While amoxicillin/clavulanate possesses the characteristic low toxicity of the penicillin group of antibiotics, periodic assessment of organ system functions, including renal, hepatic, and hematopoietic function, is advisable if therapy is for longer than the drug is approved for administration.

A high percentage of patients with mononucleosis who receive ampicillin develop an erythematous skin rash. Thus, ampicillin-class antibiotics should not be administered to patients with mononucleosis.

The possibility of superinfections with mycotic or bacterial pathogens should be kept in mind during therapy. If superinfections occur (usually involving *Pseudomonas* or *Candida*), the drug should be discontinued and/or appropriate therapy instituted.

Prescribing AUGMENTIN ES-600 in the absence of a proven or strongly suspected bacterial infection or a prophylactic indication is unlikely to provide benefit to the patient and increases the risk of the development of drug-resistant bacteria.

Information for the Patient: AUGMENTIN ES-600 should be taken every 12 hours with a meal or snack to reduce the possibility of gastrointestinal upset. If diarrhea develops and is severe or lasts more than 2 or 3 days, call your doctor.

Keep suspension refrigerated. Shake well before using. When dosing a child with the suspension (liquid) of AUGMENTIN ES-600, use a dosing spoon or medicine dropper. Be sure to rinse the spoon or dropper after each use. Bottles of suspension of AUGMENTIN ES-600 may contain more liquid than required. Follow your doctor's instructions about the amount to use and the days of treatment your child requires. Discard any unused medicine.

Patients should be counseled that antibacterial drugs including AUGMENTIN ES-600, should only be used to treat bacterial infections. They do not treat viral infections (e.g., the common cold). When AUGMENTIN ES-600 is prescribed to treat a bacterial infection, patients should be told that although it is common to feel better early in the course of therapy, the medication should be taken exactly as directed. Skipping doses or not completing the full course of therapy may: (1) decrease the effectiveness of the immediate treatment, and (2) increase the likelihood

262 that bacteria will develop resistance and will not be treatable by AUGMENTIN ES-600 or other
263 antibacterial drugs in the future.
264 **Phenylketonurics:** Each 5 mL of the 600 mg/5 mL suspension of AUGMENTIN ES-600
265 contains 7 mg phenylalanine.
266 **Drug Interactions:** Probenecid decreases the renal tubular secretion of amoxicillin. Concurrent
267 use with AUGMENTIN ES-600 may result in increased and prolonged blood levels of
268 amoxicillin. Co-administration of probenecid cannot be recommended.
269 The concurrent administration of allopurinol and ampicillin increases substantially the
270 incidence of rashes in patients receiving both drugs as compared to patients receiving ampicillin
271 alone. It is not known whether this potentiation of ampicillin rashes is due to allopurinol or the
272 hyperuricemia present in these patients. There are no data with AUGMENTIN ES-600 and
273 allopurinol administered concurrently.
274 In common with other broad-spectrum antibiotics, amoxicillin/clavulanate may reduce the
275 efficacy of oral contraceptives.
276 **Drug/Laboratory Test Interactions:** Oral administration of AUGMENTIN will result in
277 high urine concentrations of amoxicillin. High urine concentrations of ampicillin may result in
278 false-positive reactions when testing for the presence of glucose in urine using CLINITEST®,
279 Benedict's Solution, or Fehling's Solution. Since this effect may also occur with amoxicillin and
280 therefore AUGMENTIN ES-600, it is recommended that glucose tests based on enzymatic
281 glucose oxidase reactions (such as CLINISTIX®) be used.
282 Following administration of ampicillin to pregnant women, a transient decrease in plasma
283 concentration of total conjugated estriol, estriol-glucuronide, conjugated estrone, and estradiol
284 has been noted. This effect may also occur with amoxicillin and therefore
285 AUGMENTIN ES-600.
286 **Carcinogenesis, Mutagenesis, Impairment of Fertility:** Long-term studies in animals
287 have not been performed to evaluate carcinogenic potential. The mutagenic potential of
288 AUGMENTIN was investigated in vitro with an Ames test, a human lymphocyte cytogenetic
289 assay, a yeast test, and a mouse lymphoma forward mutation assay, and in vivo with mouse
290 micronucleus tests and a dominant lethal test. All were negative apart from the in vitro mouse
291 lymphoma assay where weak activity was found at very high, cytotoxic concentrations.
292 AUGMENTIN at oral doses of up to 1,200 mg/kg/day (5.7 times the maximum adult human
293 dose based on body surface area) was found to have no effect on fertility and reproductive
294 performance in rats, dosed with a 2:1 ratio formulation of amoxicillin:clavulanate.
295 **Teratogenic Effects:** Pregnancy (Category B). Reproduction studies performed in pregnant
296 rats and mice given AUGMENTIN at oral dosages up to 1,200 mg/kg/day (4.9 and 2.8 times the
297 maximum adult human oral dose based on body surface area, respectively), revealed no evidence
298 of harm to the fetus due to AUGMENTIN. There are, however, no adequate and well-controlled
299 studies in pregnant women. Because animal reproduction studies are not always predictive of
300 human response, this drug should be used during pregnancy only if clearly needed.

Labor and Delivery: Oral ampicillin-class antibiotics are generally poorly absorbed during labor. Studies in guinea pigs have shown that intravenous administration of ampicillin decreased the uterine tone, frequency of contractions, height of contractions, and duration of contractions. However, it is not known whether the use of AUGMENTIN in humans during labor or delivery has immediate or delayed adverse effects on the fetus, prolongs the duration of labor, or increases the likelihood that forceps delivery or other obstetrical intervention or resuscitation of the newborn will be necessary. In a single study in women with premature rupture of fetal membranes, it was reported that prophylactic treatment with AUGMENTIN may be associated with an increased risk of necrotizing enterocolitis in neonates.

Nursing Mothers: Ampicillin-class antibiotics are excreted in human milk; therefore, caution should be exercised when AUGMENTIN is administered to a nursing woman.

Pediatric Use: Safety and efficacy of AUGMENTIN ES-600 in infants younger than 3 months have not been established. Safety and efficacy of AUGMENTIN ES-600 have been demonstrated for treatment of acute otitis media in infants and children 3 months to 12 years (see Description of Clinical Studies).

ADVERSE REACTIONS

AUGMENTIN ES-600 is generally well tolerated. The majority of side effects observed in pediatric clinical trials of acute otitis media were either mild or moderate, and transient in nature; 4.4% of patients discontinued therapy because of drug-related side effects. The most commonly reported side effects with probable or suspected relationship to AUGMENTIN ES-600 were contact dermatitis, i.e., diaper rash (3.5%), diarrhea (2.9%), vomiting (2.2%), moniliasis (1.4%), and rash (1.1%). The most common adverse experiences leading to withdrawal that were of probable or suspected relationship to AUGMENTIN ES-600 were diarrhea (2.5%) and vomiting (1.4%).

The following adverse reactions have been reported for ampicillin-class antibiotics:

Gastrointestinal: Diarrhea, nausea, vomiting, indigestion, gastritis, stomatitis, glossitis, black "hairy" tongue, mucocutaneous candidiasis, enterocolitis, and hemorrhagic/pseudomembranous colitis. Onset of pseudomembranous colitis symptoms may occur during or after antibiotic treatment. (See WARNINGS.)

Hypersensitivity Reactions: Skin rashes, pruritus, urticaria, angioedema, serum sickness–like reactions (urticaria or skin rash accompanied by arthritis, arthralgia, myalgia, and frequently fever), erythema multiforme (rarely Stevens-Johnson syndrome), acute generalized exanthematous pustulosis, and an occasional case of exfoliative dermatitis (including toxic epidermal necrolysis) have been reported. These reactions may be controlled with antihistamines and, if necessary, systemic corticosteroids. Whenever such reactions occur, the drug should be discontinued, unless the opinion of the physician dictates otherwise. Serious and occasional fatal hypersensitivity (anaphylactic) reactions can occur with oral penicillin. (See WARNINGS.)

Liver: A moderate rise in AST (SGOT) and/or ALT (SGPT) has been noted in patients treated with ampicillin-class antibiotics, but the significance of these findings is unknown. Hepatic

dysfunction, including increases in serum transaminases (AST and/or ALT), serum bilirubin, and/or alkaline phosphatase, has been infrequently reported with AUGMENTIN. It has been reported more commonly in the elderly, in males, or in patients on prolonged treatment. The histologic findings on liver biopsy have consisted of predominantly cholestatic, hepatocellular, or mixed cholestatic-hepatocellular changes. The onset of signs/symptoms of hepatic dysfunction may occur during or several weeks after therapy has been discontinued. The hepatic dysfunction, which may be severe, is usually reversible. On rare occasions, deaths have been reported (less than 1 death reported per estimated 4 million prescriptions worldwide). These have generally been cases associated with serious underlying diseases or concomitant medications.

Renal: Interstitial nephritis and hematuria have been reported rarely. Crystalluria has also been reported (see **OVERDOSAGE**).

Hemic and Lymphatic Systems: Anemia, including hemolytic anemia, thrombocytopenia, thrombocytopenic purpura, eosinophilia, leukopenia, and agranulocytosis have been reported during therapy with penicillins. These reactions are usually reversible on discontinuation of therapy and are believed to be hypersensitivity phenomena. A slight thrombocytosis was noted in less than 1% of the patients treated with AUGMENTIN. There have been reports of increased prothrombin time in patients receiving AUGMENTIN and anticoagulant therapy concomitantly.

Central Nervous System: Agitation, anxiety, behavioral changes, confusion, convulsions, dizziness, insomnia, and reversible hyperactivity have been reported rarely.

Miscellaneous: Tooth discoloration (brown, yellow, or gray staining) has been rarely reported. Most reports occurred in pediatric patients. Discoloration was reduced or eliminated with brushing or dental cleaning in most cases.

OVERDOSAGE

Following overdosage, patients have experienced primarily gastrointestinal symptoms including stomach and abdominal pain, vomiting, and diarrhea. Rash, hyperactivity, or drowsiness have also been observed in a small number of patients.

In the case of overdosage, discontinue AUGMENTIN ES-600, treat symptomatically, and institute supportive measures as required. If the overdosage is very recent and there is no contraindication, an attempt at emesis or other means of removal of drug from the stomach may be performed. A prospective study of 51 pediatric patients at a poison control center suggested that overdosages of less than 250 mg/kg of amoxicillin are not associated with significant clinical symptoms and do not require gastric emptying.[4]

Interstitial nephritis resulting in oliguric renal failure has been reported in a small number of patients after overdosage with amoxicillin.

Crystalluria, in some cases leading to renal failure, has also been reported after amoxicillin overdosage in adult and pediatric patients. In case of overdosage, adequate fluid intake and diuresis should be maintained to reduce the risk of amoxicillin crystalluria.

Renal impairment appears to be reversible with cessation of drug administration. High blood levels may occur more readily in patients with impaired renal function because of decreased

379 renal clearance of both amoxicillin and clavulanate. Both amoxicillin and clavulanate are
380 removed from the circulation by hemodialysis.

381 **DOSAGE AND ADMINISTRATION**

382 **AUGMENTIN ES-600, 600 mg/5 mL, does not contain the same amount of clavulanic acid**
383 **(as the potassium salt) as any of the other suspensions of AUGMENTIN.**
384 **AUGMENTIN ES-600 contains 42.9 mg of clavulanic acid per 5 mL, whereas the**
385 **200 mg/5 mL suspension of AUGMENTIN contains 28.5 mg of clavulanic acid per 5 mL**
386 **and the 400 mg/5 mL suspension contains 57 mg of clavulanic acid per 5 mL. Therefore,**
387 **the 200 mg/5 mL and 400 mg/5 mL suspensions of AUGMENTIN should *not* be substituted**
388 **for AUGMENTIN ES-600, as they are not interchangeable.**
389 **Dosage:** ***Pediatric patients 3 months and older:*** Based on the amoxicillin component
390 (600 mg/5 mL), the recommended dose of AUGMENTIN ES-600 is 90 mg/kg/day divided every
391 12 hours, administered for 10 days (see chart below).

Body Weight (kg)	Volume of AUGMENTIN ES-600 providing 90 mg/kg/day
8	3.0 mL twice daily
12	4.5 mL twice daily
16	6.0 mL twice daily
20	7.5 mL twice daily
24	9.0 mL twice daily
28	10.5 mL twice daily
32	12.0 mL twice daily
36	13.5 mL twice daily

392 ***Pediatric patients weighing 40 kg and more:*** Experience with AUGMENTIN ES-600
393 (600 mg/5 mL formulation) in this group is not available.
394 ***Adults:*** Experience with AUGMENTIN ES-600 (600 mg/5 mL formulation) in adults is not
395 available and adults who have difficulty swallowing should not be given AUGMENTIN ES-600
396 (600 mg/5 mL) in place of the 500-mg or 875-mg tablet of AUGMENTIN.
397 Hepatically impaired patients should be dosed with caution and hepatic function monitored at
398 regular intervals. (See WARNINGS.)
399 ***Directions for Mixing Oral Suspension:*** Prepare a suspension at time of dispensing as
400 follows: Tap bottle until all the powder flows freely. Add approximately 2/3 of the total amount
401 of water for reconstitution (see table below) and shake vigorously to suspend powder. Add
402 remainder of the water and again shake vigorously.

AUGMENTIN ES-600 (600 mg/5 mL Suspension)

Bottle Size	Amount of Water Required for Reconstitution
75 mL	65 mL
125 mL	110 mL
200 mL	175 mL

Each teaspoonful (5 mL) will contain 600 mg amoxicillin as the trihydrate and 42.9 mg of clavulanic acid as the potassium salt.

Note: SHAKE ORAL SUSPENSION WELL BEFORE USING.

Administration: To minimize the potential for gastrointestinal intolerance, AUGMENTIN ES-600 should be taken at the start of a meal. Absorption of clavulanate potassium may be enhanced when AUGMENTIN ES-600 is administered at the start of a meal.

HOW SUPPLIED

AUGMENTIN ES-600, 600 mg/5 mL, for Oral Suspension: Each 5 mL of reconstituted orange-flavored suspension contains 600 mg amoxicillin and 42.9 mg clavulanic acid as the potassium salt.

NDC 0029-6094-39 .....75 mL bottle

NDC 0029-6094-45 .....125 mL bottle

NDC 0029-6094-24 .....200 mL bottle

STORAGE

Store reconstituted suspension under refrigeration. Discard unused suspension after 10 days. Store dry powder for oral suspension at or below 25°C (77°F). Dispense in original container.

Description of Clinical Studies

Two clinical studies were conducted in pediatric patients with acute otitis media.

A non-comparative, open-label study assessed the bacteriologic and clinical efficacy of AUGMENTIN ES-600 (90/6.4 mg/kg/day, divided every 12 hours) for 10 days in 521 pediatric patients (3 to 50 months) with acute otitis media. The primary objective was to assess bacteriological response in children with acute otitis media due to *S. pneumoniae* with amoxicillin/clavulanic acid MICs of 4 mcg/mL. The study sought the enrollment of patients with the following risk factors: Failure of antibiotic therapy for acute otitis media in the previous 3 months, history of recurrent episodes of acute otitis media, ≤2 years, or daycare attendance. Prior to receiving AUGMENTIN ES-600, all patients had tympanocentesis to obtain middle ear fluid for bacteriological evaluation. Patients from whom *S. pneumoniae* (alone or in combination with other bacteria) was isolated had a second tympanocentesis 4 to 6 days after the start of therapy. Clinical assessments were planned for all patients during treatment (4-6 days after starting therapy), as well as 2-4 days post-treatment and 15-18 days post-treatment. Bacteriological success was defined as the absence of the pretreatment pathogen from the on-therapy tympanocentesis specimen. Clinical success was defined as improvement or resolution of signs and symptoms. Clinical failure was defined as lack of improvement or worsening of signs and/or symptoms at any time following at least 72 hours of

434 AUGMENTIN ES-600; patients who received an additional systemic antibacterial drug for otitis
435 media after 3 days of therapy were considered clinical failures. Bacteriological eradication on
436 therapy (day 4-6 visit) in the per protocol population is summarized in the following table:
437
438 **Table 3. Bacteriologic Eradication Rates in the Per Protocol Population**

	Bacteriologic Eradication on Therapy		
Pathogen	**n/N**	**%**	**95% CI***
All *S. pneumoniae*	121/123	98.4	(94.3, 99.8)
S. pneumoniae with penicillin MIC = 2 mcg/mL	19/19	100	(82.4, 100.0)
S. pneumoniae with penicillin MIC = 4 mcg/mL	12/14	85.7	(57.2, 98.2)
H. influenzae	75/81	92.6	(84.6, 97.2)
M. catarrhalis	11/11	100	(71.5, 100.0)

439 *CI = confidence intervals; 95% CIs are not adjusted for multiple comparisons.
440
441 Clinical assessments were made in the per protocol population 2-4 days post-therapy and
442 15-18 days post-therapy. Patients who responded to therapy 2-4 days post-therapy were followed
443 for 15-18 days post-therapy to assess them for acute otitis media. Nonresponders at 2-4 days
444 post-therapy were considered failures at the latter timepoint.
445

446 **Table 4. Clinical Assessments in the Per Protocol Population (Includes *S. pneumoniae***
447 **Patients With Penicillin MICs = 2 or 4 mcg/mL*)**

Pathogen	2-4 Days Post-Therapy (Primary Endpoint)		
	n/N	**%**	**95% CI†**
All *S. pneumoniae*	122/137	89.1	(82.6, 93.7)
S. pneumoniae with penicillin MIC = 2 mcg/mL	17/20	85.0	(62.1, 96.8)
S. pneumoniae with penicillin MIC = 4 mcg/mL	11/14	78.6	(49.2, 95.3)
H. influenzae	141/162	87.0	(80.9, 91.8)
M. catarrhalis	22/26	84.6	(65.1, 95.6)
	15-18 Days Post-Therapy‡ (Secondary Endpoint)		
	n/N	**%**	**95% CI†**
All *S. pneumoniae*	95/136	69.9	(61.4, 77.4)
S. pneumoniae with penicillin MIC = 2 mcg/mL	11/20	55.0	(31.5, 76.9)
S. pneumoniae with penicillin MIC = 4 mcg/mL	5/14	35.7	(12.8, 64.9)
H. influenzae	106/156	67.9	(60.0, 75.2)
M. catarrhalis	14/25	56.0	(34.9, 75.6)

448 **S. pneumoniae* strains with penicillin MICs of 2 or 4 mcg/mL are considered resistant to
449 penicillin.
450 †CI = confidence intervals; 95% CIs are not adjusted for multiple comparisons.
451 ‡Clinical assessments at 15-18 days post-therapy may have been confounded by viral
452 infections and new episodes of acute otitis media with time elapsed post-treatment.
453
454 In the intent-to-treat analysis, overall clinical outcomes at 2-4 days and 15-18 days
455 post-treatment in patients with *S. pneumoniae* with penicillin MIC = 2 mcg/mL and 4 mcg/mL
456 were 29/41 (71%) and 17/41 (41.5%), respectively.
457 In the intent-to-treat population of 521 patients, the most frequently reported adverse events
458 were vomiting (6.9%), fever (6.1%), contact dermatitis (i.e., diaper rash) (6.1%), upper
459 respiratory tract infection (4.0%), and diarrhea (3.8%). Protocol-defined diarrhea (i.e., 3 or more
460 watery stools in one day or 2 watery stools per day for 2 consecutive days as recorded on diary
461 cards) occurred in 12.9% of patients.
462 A double-blind, randomized, clinical study compared AUGMENTIN ES-600
463 (90/6.4 mg/kg/day, divided every 12 hours) to AUGMENTIN (45/6.4 mg/kg/day, divided every
464 12 hours) for 10 days in 450 pediatric patients (3 months to 12 years) with acute otitis media.
465 The primary objective of the study was to compare the safety of AUGMENTIN ES-600 to
466 AUGMENTIN. There was no statistically significant difference between treatments in the
467 proportion of patients with 1 or more adverse events. The most frequently reported adverse

events for AUGMENTIN ES-600 and the comparator of AUGMENTIN were coughing (11.9% versus 6.8%), vomiting (6.5% versus 7.7%), contact dermatitis (i.e., diaper rash, 6.0% versus 4.8%), fever (5.5% versus 3.9%), and upper respiratory infection (3.0% versus 9.2%), respectively. The frequencies of protocol-defined diarrhea with AUGMENTIN ES-600 (11.1%) and AUGMENTIN (9.4%) were similar (95% confidence interval on difference: –4.2% to 7.7%). Only 2 patients in the group treated with AUGMENTIN ES-600 and 1 patient in the group treated with AUGMENTIN were withdrawn due to diarrhea.

REFERENCES

1. National Committee for Clinical Microbiology Standards. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically* – Sixth Edition. Approved Standard. NCCLS Document M7-A6, Vol. 23, No. 2. (ISBN-1-56238-486-4) NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2003.
2. National Committee for Clinical Microbiology Standards. *Performance Standards for Antimicrobial Susceptibility Testing* – Thirteenth Informational Supplement. NCCLS Document M100-S13 (M7), Vol. 23, No. 2. NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2003.
3. National Committee for Clinical Microbiology Standards. *Performance Standards for Antimicrobial Disk Susceptibility Tests* – Eighth Edition. Approved Standard. NCCLS Document M2-A8, Vol. 23, No. 1. (ISBN-1-56238-485-6). NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087-1898, January 2003.
4. Swanson-Biearman B, Dean BS, Lopez G, Krenzelok EP. The effects of penicillin and cephalosporin ingestions in children less than six years of age. *Vet Hum Toxicol* 1988; 30:66-67.

AUGMENTIN ES-600 is a registered trademark of GlaxoSmithKline.
CLINITEST is a registered trademark of Miles, Inc.
CLINISTIX is a registered trademark of Bayer Corporation.

GlaxoSmithKline
Research Triangle Park, NC 27709

©2004, GlaxoSmithKline. All rights reserved.

Month 2004 AE:L9