

**The Xyrem Success Program<sup>®</sup> for Patients - Video Script**  
***A Patient's Guide to Success with Xyrem<sup>®</sup> (sodium oxybate) oral solution***

This video is one of the components of a comprehensive program for patients. It is designed solely as an instructional tool to teach patients how to prepare and use the medication, about in-home safety, and issues regarding misuse of the drug. This video contains no product promotional claims.

Video	Audio
<p>Title: <i>The Xyrem Success Program<sup>®</sup>: A Patient's Guide to Success with Xyrem<sup>®</sup> (sodium oxybate) oral solution</i></p>	<p>Music up full</p>
<p>Host on camera</p> <p>Cut away shot of Xyrem box and Xyrem bottle</p>	<p><u>Host on Camera</u> Your doctor has determined that you may benefit from taking Xyrem. Xyrem has been shown to reduce excessive daytime sleepiness and cataplexy in patients with narcolepsy.</p>
<p>Host on camera</p>	<p>You are being asked to watch this video presentation before taking your first dose of Xyrem to help you feel confident about how to use the medication.</p>
<p>Shot of hands dispensing medication with text on screen:</p> <ul style="list-style-type: none"> <li>• <i>How to prepare and use Xyrem</i></li> <li>•</li> </ul>	<p><u>Narration</u> In this video you will learn: How to prepare and use Xyrem properly,</p>
<p>Shot of medication being placed in a secure, but not locked cabinet. With text on screen:</p> <ul style="list-style-type: none"> <li>• <i>In-Home Safety Procedures</i></li> <li>•</li> </ul>	<p>correct in-home safety procedures,</p>
<p>Patient reading patient brochure with text on screen:</p> <ul style="list-style-type: none"> <li>• <i>Precautions to consider</i></li> <li>•</li> </ul>	<p>and what precautions you should consider while you're using Xyrem.</p>
<p>Chapter Heading: <i>How to Use Xyrem Properly</i> Host on camera</p>	<p>Music up full</p>
<p>Host on camera</p>	<p><u>Host on Camera</u> Xyrem is different than other medications that</p>

Video	Audio
	<p>you may be taking so it's important that you follow your doctor's instructions.</p> <p>Take Xyrem two times each night or as directed by your doctor.</p>
<p>Cut to shot of physician showing the patient a dosing plan.</p>	<p><u>Narrator</u> You can expect to see some improvement in your symptoms soon after beginning Xyrem therapy. If you're not experiencing any change in your symptoms, contact your doctor for an assessment and possible dose adjustment.</p>
<p>Text on screen: <i>Never change the dose of Xyrem yourself.</i></p>	<p>Never change the dose of Xyrem without direction from your doctor.</p>
<p>Shot of physician and patient continues</p>	<p>Response to Xyrem varies from patient to patient and it may take several weeks for you to achieve the maximum beneficial effect of Xyrem.</p>
<p>Medium shot of carton, pan up Host on camera with carton, bottle, dispensing syringe, 2 containers, and a medication guide. Each element adds on to counter as it is mentioned.</p>	<p><u>Narrator</u> Your Xyrem carton contains one bottle of Xyrem oral solution, a measuring device, two dosing cups with child-resistant caps for preparing and storing your two night-time doses of Xyrem and a Xyrem Medication Guide.</p>
<p>Close up of Medication Guide</p>	<p><u>Narrator</u> Be sure to read the Medication Guide for additional important information before taking your first dose of Xyrem.</p>
<p>Host of camera with text on screen: <i>Prepare both doses at bedtime.</i></p>	<p><u>Host on camera</u> You will need to prepare both doses at bedtime.</p>
<p>Medium and close up shots of patient removing pieces from the box, unwrapping measuring device, and removing cap.</p>	<p><u>Narrator</u> Remove the Xyrem bottle, measuring device, and dosing cups from the box. - Unwrap the measuring device.</p>
<p>Cut to close-up shot of patient's hands turning cap on medicine bottle and setting bottle on counter.</p>	<p>Take the child resistant cap off the first dosing cup. To do this, push the cap down while turning it counterclockwise. Set the bottle upright on a tabletop.</p>

Video	Audio
Cut to medium shot of patient holding bottle and inserting measuring device with one hand.	While holding the bottle and the measuring device with one hand, insert the tip of the device into the center opening of the bottle and press down firmly.
Cut to close up shot of fingers pulling up plunger, then extreme close up of gram markers on the plunger.	Using your other hand, slowly pull the plunger of the measuring device up, until the point on the grams scale corresponds with half the total nightly dose that your doctor has prescribed.
Freeze frame shot of plunger and gram scale defocused: Text on screen: <i>Example</i> <i>Dose=6 grams/per night</i> <i>Extract 3 grams X 2 times</i>	For example, if your total nightly dose is 6 grams per night, you will need to extract 3 grams from the bottle for each nightly dose.
Cut to medium shot of bottle in upright position	Remember to keep the bottle upright or the medicine will not flow into the measuring device properly. (moved)
Cut to close-up shot of patient removing measuring device from bottle, putting tip into the dosing cup and pressing down plunger.	Now remove the measuring device from the Xyrem bottle and empty the medication into one of the dosing cups by slowly pressing down on the plunger.
Medium shot of patient extracting Xyrem and putting in dosing cup again	Extract the medication for the second dosing cup.
Medium shot of patient pouring 2 oz. of water into each dosing cup.	Next, pour approximately two ounces or one quarter cup of water into each dosing cup.
Cut to shot of patient securing caps on bottle and cups and checking. Medium shot of placing medicine in a cabinet.	Put the child-resistant caps back on both dosing cups, making sure that they are secure. Also make sure that you have securely replaced the cap on the bottle of Xyrem. Then place the medication in a safe location.
Dissolve to wider shot of patient placing dosing cups inside drawer of bedside table.	Place the containers near your bed in an area that is not visible to children or pets.
Shot of patient getting into bed and setting alarm clock	Before you take your first dose, set an alarm clock to wake you up in 4 hours so you can take

Video	Audio
	your second dose. If you wake up before the alarm goes off and it has been at least 2 1/2 hours, turn off your alarm and take your second dose.
Dissolve to shot of patient sitting up in bed, drinking first dose, and then laying down.	Drink all of the first dose right before bedtime. Because Xyrem is a medicine that produces sleepiness; as a safety precaution, it is best to be in bed when you take it.
Slow dissolve showing passage of time: shot of patient waking up, turning off alarm, and taking second dose. Shot of patient lies back down to sleep	When your alarm rings, remove the cap from the second dosing cup and drink the entire contents before going back to sleep.
Dissolve to shot of patient waking up happily and stretching in the morning.	It's important that you take Xyrem exactly as your doctor has prescribed. If you miss the second dose, skip that dose and don't take Xyrem again until the next night.
Text on screen: <i>Never take both doses at once.</i>	<u>Host on camera</u> Never take both Xyrem doses at once.
Host holds up Patient Success Program Brochure	The Xyrem Patient Success Program brochure includes tips on traveling with Xyrem. Be sure to read this material before leaving on a trip.
Chapter Heading: <i>Safe Disposal of the Medication</i>	Music up full
Close up shot of hands trying to dispense medication from an almost empty bottle.	<u>Narrator</u> When you can no longer pull medication out of the bottle with the dispensing device, it's time to throw the bottle away. Pour any unused Xyrem down the drain.
Cut to shot of patient marking label with a black permanent marker.	Use a pen to mark up the bottle's label so that it cannot be reused.
Cut to shot of patient throwing bottle in the trashcan.	It is now safe to throw the empty bottle away.
Text on screen: <i>Emergency Procedures</i>	Music up full

Video	Audio
Patient putting Xyrem in cabinet	Xyrem, like all medications, must be kept out of the reach of children and pets.
Medium shot of patient dialing telephone	In case of accidental ingestion or overdose, immediately call 911 or your local poison control center.
Host on camera	<u>Host on camera</u> Xyrem is a controlled substance, which means that it is only available by prescription and should only be used for appropriate medical purposes.
Text on screen: <i>Xyrem must be:</i> <ul style="list-style-type: none"> <li>• <i>Used only by you</i></li> <li>• <i>Used only as directed by your physician</i></li> <li>• <i>Always stored in a safe and secure place</i></li> </ul>	<u>Narrator</u> It is very important that your Xyrem be used only by you, only as directed by your physician, and that it always be stored in a safe and secure place.
Host on camera	<u>Host on camera</u> Anyone who knowingly distributes, shares or uses Xyrem for purposes other than it was prescribed will be subject to criminal penalties outlined in the Controlled Substances Act.
Jazz Pharmaceuticals logo animates on screen	At Jazz Pharmaceuticals we are working to deliver your prescribed Xyrem in a safe and timely manner.
Host on camera	<u>Host on camera</u> Working together, we can ensure that you continue to receive the medical benefits of Xyrem. If you have any questions or concerns about Xyrem, be sure to talk to your doctor...
On screen text: <i>The Xyrem Success Program</i> <sup>®</sup> <i>1-866-997-3688</i>	<u>Narrator</u> or call the pharmacist at The Xyrem Success Program, toll free at 1-866-997-3688.
Copyright Jazz Pharmaceuticals 2006	Total approximate length: 7:15