

28 Count Outer Carton



100% SIZE  
12 digit UPC  
(non-suppressed)  
For Position Only  
0 37000 35906 7

**Prilosec**  
OTC

Safety Feature—  
Do not use  
if tablet blister  
unit is open  
or torn.

NDC 37000-455-03

Easy Push-Through Package

Treats Frequent  
Heartburn!

**Prilosec**  
omeprazole delayed-release tablets  
20 mg / acid reducer **OTC**

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

28 TABLETS

Two 14-day courses of treatment

Dist. by Preder & Gambis, Cincinnati, OH 45202  
Product of Sweden 95227633

**Prilosec**  
OTC

**Prilosec**  
OTC

Questions? 1-800-289-9181

**Inactive ingredients** glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polyborate 80, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate

**Drug Facts (continued)**  
■ protect product from moisture  
■ keep product out of high heat and humidity  
■ store at 20-25C (68-77F)  
■ keep the carton and package insert. They contain important information.

**Ask a doctor or pharmacist before use if you are taking**  
■ warfarin (blood-thinning medicine)  
■ prescription antifungal or anti-yeast medicines  
■ diazepam (anxiety medicine) ■ digoxin (heart medicine)  
**Ask a doctor before use if you have**  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness  
■ frequent chest pain  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting ■ stomach pain  
**Do not use if you have**  
■ trouble or pain swallowing food  
■ vomiting with blood  
■ bloody or black stools  
These may be signs of a serious condition. See your doctor.

**Warnings**  
**Allergy alert:** Do not use if you are allergic to omeprazole  
**Use**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect  
**Purpose**  
Omeprazole magnesium delayed-release tablet 20.5 mg (equivalent to 20 mg omeprazole)  
Acid reducer

**Drug Facts (continued)**  
**Stop use and ask a doctor if**  
■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months  
**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions**  
■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours  
**14-Day Course of Treatment**  
■ swallow 1 tablet with a glass of water before eating in the morning  
■ take every day for 14 days  
■ do not take more than 1 tablet a day  
■ do not chew or crush the tablets  
■ do not use for more than 14 days unless directed by your doctor  
**Repeated 14-Day Courses (if needed)**  
■ do not crush tablets in food  
■ do not take more than 1 tablet a day  
■ do not take more than 14 days unless directed by your doctor  
**Other information**  
■ children under 18 years of age: ask a doctor  
■ every 4 months unless directed by a doctor  
■ do not take for more than 14 days or more often than  
■ you may repeat a 14-day course every 4 months  
■ read the directions, warnings and package insert before use

make a mark in this area  
FPO 128 code  
[Barcode]

make a mark in this area  
FPO 128 code  
[Barcode]



14 Count Inner Carton



**Drug Facts** (continued)

**Active ingredient (in each tablet) Purpose**  
Omeprazole magnesium delayed-release tablet 20.6 mg (equivalent to 20 mg omeprazole).....Acid reducer

**Use**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:** Do not use if you are allergic to omeprazole  
 Do not use if you have  
 ■ trouble or pain swallowing food  
 ■ vomiting with blood  
 ■ bloody or black stools  
 These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
 ■ frequent chest pain  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss  
 ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are taking**  
 ■ warfarin (blood-thinning medicine)  
 ■ prescription antifungal or anti-yeast medicines  
 ■ diazepam (anxiety medicine) ■ digoxin (heart medicine)

**Other information**  
 ■ read the directions, warnings and package insert before use

**Drug Facts** (continued)

**Stop use and ask a doctor if**  
 ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months

**if pregnant or breast-feeding**, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ it may take 1 to 4 days for full effect, although some people get complete relief of symptoms within 24 hours

**14-Day Course of Treatment**  
 ■ swallow 1 tablet with a glass of water before eating in the morning  
 ■ take every day for 14 days  
 ■ do not take more than 1 tablet a day  
 ■ do not chew or crush the tablets  
 ■ do not crush tablets in food  
 ■ do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor

■ children under 18 years of age: ask a doctor

FPO 128 code

FPO 128 code

**NOT FOR RESALE**

**Prilosec OTC**

Safety Feature—Do not use if tablet blister unit is open or torn.

NDC 37000-455-02

**Easy Push-Through Package**

**Treats Frequent Heartburn!**

# Prilosec

omeprazole delayed-release tablets 20 mg / acid reducer **OTC**®

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**14 TABLETS**

One 14-day course of treatment

**Drug Facts** (continued)

■ keep the carton and package insert. They contain important information.  
 ■ store at 20-25C (68-77F)  
 ■ keep product out of high heat and humidity ■ protect product from moisture

Dist. by  
 Procter & Gamble  
 Cincinnati, OH 45202

**Drug Facts** (continued)  
**Questions? 1-800-289-9181**

Product of Sweden 95227631

**Drug Facts** (continued)

**Inactive ingredients** glyceryl monostearate, hydroxypropyl cellulose, hypromellose, iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, paraffin, polyethylene glycol 6000, polysorbate 80, polyvinylpyrrolidone, sodium stearyl fumarate, starch, sucrose, talc, titanium dioxide, triethyl citrate



# Blister Card (Non-CR)

Do not  
chew or  
crush tablets

**Prilosec**  
OTC®

Do not  
crush tablets  
in food

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble, Cincinnati, OH 45202

**PACKAGE NOT CHILD-RESISTANT**

95437426

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202

**Prilosec**  
OTC®

1 omeprazole delayed-release tablet  
20 mg / acid reducer

**Push tablet through foil.**

Dist. by Procter & Gamble,  
Cincinnati, OH 45202