

NET CONTENTS: 3000 INDIVIDUAL SWABS

REORDER NO. B11400

NDC 10819-1080-2

Chlorascrub™ Swab

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]

Antiseptic

*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% w/v	Antiseptic

Use for the preparation of the patient's skin prior to an injection

Warnings

For external use only
Flammable, keep away from fire or flame
Do not use with electrocautery procedures

- Do not use**
- in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption
 - on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
 - for lumbal puncture or in contact with the meninges
 - on open skin wounds or as a general skin cleanser
 - under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)



80B11400

Each Swab contains 0.034 fl.oz. (1 mL)



PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdipdi.com
Chlorascrub™ is under license by American Solumed Inc.

2.5" width

REORDER NO. B11400
NDC 10819-1080-2

Chlorascrub™ Swab
[Chlorhexidine Gluconate (3.15%)* and
Isopropyl Alcohol (70%) Swab] Antiseptic

*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

Drug Facts	
Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v.....	Antiseptic
Isopropyl alcohol 70% w/v.....	Antiseptic

Use for the preparation of the patient's skin prior to an injection

Warnings

For external use only
Flammable, keep away from fire or flame
Do not use with electrocautery procedures

Do not use

- in premature or low birth weight infants, or on children less than 2 months of age because of the potential for excessive skin irritation and increased drug absorption
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient
USP purified water

Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

Each Swab contains 0.034 fl.oz. (1 mL)

PDI PDI®, The Healthcare Division of Nice-Pak Products, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdiipdi.com
Chlorascrub™ is under license by American Solumed Inc.

50B114IS

6.5" height

<p>NDC 10819-1080-2 REORDER NO. B11400</p>	<p>TEAR HERE</p>
<p>Chlorascrub™ Swab</p>	<p>LOT</p>
<p>(Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab)</p>	<p>CONTENTS: One swab premoistened with 1 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) isopropyl alcohol, and USP purified water.</p>
<p>Antiseptic</p>	<p>DIRECTIONS: Maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm). Tear open packet and remove swab. Do not unfold swab. Prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds. Allow the prepped area to air dry for 30 seconds. Do not blot or wipe dry. Discard after a single use. WARNINGS: For external use only. Flammable, keep away from fire or flame. Do not use with electrocautery procedures. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>*equivalent to 167 mg of Chlorhexidine Gluconate per pouch</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>■ For external use only</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>■ Use for the preparation of the patient's skin prior to an injection</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>■ Helps reduce bacteria that potentially can cause skin infection</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>■ For hospital and professional use only</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>■ Latex free</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>	<p>IMPORTANT: See insert for complete Drug Facts and safety information.</p>
<p>One Swab, 0.034 fl.oz. (1 mL) Each</p>	<p>2AB114</p>
<p>FDA</p>	<p>EXP.</p>
<p>Made in U.S.A.</p>	