



- If you miss your dose, do not double your next dose.** Take only your usual dose 1 to 3 hours before your next bedtime.
- Contact your doctor, if you stop taking ropinirole hydrochloride tablets for any reason. Do not restart without consulting your doctor.
- You can take ropinirole hydrochloride tablets with or without food. Taking ropinirole hydrochloride with food may decrease the chances of feeling nauseated.

#### What are the possible side effects of ropinirole hydrochloride?

- Most people who take ropinirole hydrochloride tolerate it well. The most commonly reported side effects in people taking ropinirole hydrochloride for RLS are nausea, vomiting, dizziness, and drowsiness or sleepiness. You should be careful until you know if ropinirole hydrochloride affects your ability to remain alert while doing normal daily activities, and you should watch for the development of significant daytime sleepiness or episodes of falling asleep. It is possible that you could fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole hydrochloride. Your chances of falling asleep while doing normal activities while taking ropinirole hydrochloride are greater if you are taking other medicines that cause drowsiness.
- When you start taking ropinirole hydrochloride or when you increase your dose, you may feel dizzy, nauseated, sweaty or faint, when first standing up from sitting or lying down. Therefore, do not stand up quickly after sitting or lying down, particularly if you have been sitting or lying down for a long period of time. Take a minute sitting on the edge of the bed or chair before you get up.
- Hallucinations (unreal sounds, visions, or sensations) have been reported in patients taking ropinirole hydrochloride. These were uncommon in patients taking ropinirole hydrochloride for RLS. The risk is greater in patients taking ropinirole hydrochloride with L-dopa, or taking higher doses of ropinirole hydrochloride than recommended for RLS.

This is not a complete list of side effects and should not take the place of discussions with your healthcare providers. Your doctor or pharmacist can give you a more complete list of possible side effects. Talk to your doctor about any side effects or problems you may have.

#### Other information about ropinirole hydrochloride

Patients being treated with ropinirole hydrochloride should have periodic skin examinations for melanoma.

- Take ropinirole hydrochloride exactly as your doctor prescribes it.
- Do not share ropinirole hydrochloride with other people, even if they have the same symptoms you have.
- Keep ropinirole hydrochloride out of the reach of children.
- Store ropinirole hydrochloride at room temperature out of direct sunlight.
- Keep ropinirole hydrochloride in a tightly closed container.

This leaflet summarizes important information about ropinirole hydrochloride. Medicines are sometimes prescribed for purposes other than those listed in this leaflet. Do not take ropinirole hydrochloride for a condition for which it was not prescribed. For more information, talk with your doctor or pharmacist. They can give you information about ropinirole hydrochloride that is written for healthcare professionals.

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008

- If you miss your dose, do not double your next dose.** Take only your usual dose 1 to 3 hours before your next bedtime.
- Contact your doctor, if you stop taking ropinirole hydrochloride tablets for any reason. Do not restart without consulting your doctor.
- You can take ropinirole hydrochloride tablets with or without food. Taking ropinirole hydrochloride with food may decrease the chances of feeling nauseated.

#### What are the possible side effects of ropinirole hydrochloride?

- Most people who take ropinirole hydrochloride tolerate it well. The most commonly reported side effects in people taking ropinirole hydrochloride for RLS are nausea, vomiting, dizziness, and drowsiness or sleepiness. You should be careful until you know if ropinirole hydrochloride affects your ability to remain alert while doing normal daily activities, and you should watch for the development of significant daytime sleepiness or episodes of falling asleep. It is possible that you could fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole hydrochloride. Your chances of falling asleep while doing normal activities while taking ropinirole hydrochloride are greater if you are taking other medicines that cause drowsiness.
- When you start taking ropinirole hydrochloride or when you increase your dose, you may feel dizzy, nauseated, sweaty or faint, when first standing up from sitting or lying down. Therefore, do not stand up quickly after sitting or lying down, particularly if you have been sitting or lying down for a long period of time. Take a minute sitting on the edge of the bed or chair before you get up.
- Hallucinations (unreal sounds, visions, or sensations) have been reported in patients taking ropinirole hydrochloride. These were uncommon in patients taking ropinirole hydrochloride for RLS. The risk is greater in patients taking ropinirole hydrochloride with L-dopa, or taking higher doses of ropinirole hydrochloride than recommended for RLS.

This is not a complete list of side effects and should not take the place of discussions with your healthcare providers. Your doctor or pharmacist can give you a more complete list of possible side effects. Talk to your doctor about any side effects or problems you may have.

#### Other information about ropinirole hydrochloride

Patients being treated with ropinirole hydrochloride should have periodic skin examinations for melanoma.

- Take ropinirole hydrochloride exactly as your doctor prescribes it.
- Do not share ropinirole hydrochloride with other people, even if they have the same symptoms you have.
- Keep ropinirole hydrochloride out of the reach of children.
- Store ropinirole hydrochloride at room temperature out of direct sunlight.
- Keep ropinirole hydrochloride in a tightly closed container.

This leaflet summarizes important information about ropinirole hydrochloride. Medicines are sometimes prescribed for purposes other than those listed in this leaflet. Do not take ropinirole hydrochloride for a condition for which it was not prescribed. For more information, talk with your doctor or pharmacist. They can give you information about ropinirole hydrochloride that is written for healthcare professionals.

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008

- If you miss your dose, do not double your next dose.** Take only your usual dose 1 to 3 hours before your next bedtime.
- Contact your doctor, if you stop taking ropinirole hydrochloride tablets for any reason. Do not restart without consulting your doctor.
- You can take ropinirole hydrochloride tablets with or without food. Taking ropinirole hydrochloride with food may decrease the chances of feeling nauseated.

#### What are the possible side effects of ropinirole hydrochloride?

- Most people who take ropinirole hydrochloride tolerate it well. The most commonly reported side effects in people taking ropinirole hydrochloride for RLS are nausea, vomiting, dizziness, and drowsiness or sleepiness. You should be careful until you know if ropinirole hydrochloride affects your ability to remain alert while doing normal daily activities, and you should watch for the development of significant daytime sleepiness or episodes of falling asleep. It is possible that you could fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole hydrochloride. Your chances of falling asleep while doing normal activities while taking ropinirole hydrochloride are greater if you are taking other medicines that cause drowsiness.
- When you start taking ropinirole hydrochloride or when you increase your dose, you may feel dizzy, nauseated, sweaty or faint, when first standing up from sitting or lying down. Therefore, do not stand up quickly after sitting or lying down, particularly if you have been sitting or lying down for a long period of time. Take a minute sitting on the edge of the bed or chair before you get up.
- Hallucinations (unreal sounds, visions, or sensations) have been reported in patients taking ropinirole hydrochloride. These were uncommon in patients taking ropinirole hydrochloride for RLS. The risk is greater in patients taking ropinirole hydrochloride with L-dopa, or taking higher doses of ropinirole hydrochloride than recommended for RLS.

This is not a complete list of side effects and should not take the place of discussions with your healthcare providers. Your doctor or pharmacist can give you a more complete list of possible side effects. Talk to your doctor about any side effects or problems you may have.

#### Other information about ropinirole hydrochloride

Patients being treated with ropinirole hydrochloride should have periodic skin examinations for melanoma.

- Take ropinirole hydrochloride exactly as your doctor prescribes it.
- Do not share ropinirole hydrochloride with other people, even if they have the same symptoms you have.
- Keep ropinirole hydrochloride out of the reach of children.
- Store ropinirole hydrochloride at room temperature out of direct sunlight.
- Keep ropinirole hydrochloride in a tightly closed container.

This leaflet summarizes important information about ropinirole hydrochloride. Medicines are sometimes prescribed for purposes other than those listed in this leaflet. Do not take ropinirole hydrochloride for a condition for which it was not prescribed. For more information, talk with your doctor or pharmacist. They can give you information about ropinirole hydrochloride that is written for healthcare professionals.

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008

- If you miss your dose, do not double your next dose.** Take only your usual dose 1 to 3 hours before your next bedtime.
- Contact your doctor, if you stop taking ropinirole hydrochloride tablets for any reason. Do not restart without consulting your doctor.
- You can take ropinirole hydrochloride tablets with or without food. Taking ropinirole hydrochloride with food may decrease the chances of feeling nauseated.

#### What are the possible side effects of ropinirole hydrochloride?

- Most people who take ropinirole hydrochloride tolerate it well. The most commonly reported side effects in people taking ropinirole hydrochloride for RLS are nausea, vomiting, dizziness, and drowsiness or sleepiness. You should be careful until you know if ropinirole hydrochloride affects your ability to remain alert while doing normal daily activities, and you should watch for the development of significant daytime sleepiness or episodes of falling asleep. It is possible that you could fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole hydrochloride. Your chances of falling asleep while doing normal activities while taking ropinirole hydrochloride are greater if you are taking other medicines that cause drowsiness.
- When you start taking ropinirole hydrochloride or when you increase your dose, you may feel dizzy, nauseated, sweaty or faint, when first standing up from sitting or lying down. Therefore, do not stand up quickly after sitting or lying down, particularly if you have been sitting or lying down for a long period of time. Take a minute sitting on the edge of the bed or chair before you get up.
- Hallucinations (unreal sounds, visions, or sensations) have been reported in patients taking ropinirole hydrochloride. These were uncommon in patients taking ropinirole hydrochloride for RLS. The risk is greater in patients taking ropinirole hydrochloride with L-dopa, or taking higher doses of ropinirole hydrochloride than recommended for RLS.

This is not a complete list of side effects and should not take the place of discussions with your healthcare providers. Your doctor or pharmacist can give you a more complete list of possible side effects. Talk to your doctor about any side effects or problems you may have.

#### Other information about ropinirole hydrochloride

Patients being treated with ropinirole hydrochloride should have periodic skin examinations for melanoma.

- Take ropinirole hydrochloride exactly as your doctor prescribes it.
- Do not share ropinirole hydrochloride with other people, even if they have the same symptoms you have.
- Keep ropinirole hydrochloride out of the reach of children.
- Store ropinirole hydrochloride at room temperature out of direct sunlight.
- Keep ropinirole hydrochloride in a tightly closed container.

This leaflet summarizes important information about ropinirole hydrochloride. Medicines are sometimes prescribed for purposes other than those listed in this leaflet. Do not take ropinirole hydrochloride for a condition for which it was not prescribed. For more information, talk with your doctor or pharmacist. They can give you information about ropinirole hydrochloride that is written for healthcare professionals.

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008

pneumonia, respiratory infection, respiratory tract infection, tonsillitis, tooth infection, vaginal candidiasis, vaginal infection, vaginal mycosis, viral infection, viral upper respiratory tract infection, wound infection.

***Injury, Poisoning, and Procedural Complications:*** *Inrequent:* Concussion, lower limb fracture, post procedural hemorrhage, road traffic accident.

***Investigations:*** *Inrequent:* Blood cholesterol increased, blood iron decreased, blood pressure increased, blood urine present, hemoglobin decreased, heart rate increased, protein urine present, weight decreased, weight increased.

***Metabolism and Nutrition Disorders:*** *Inrequent:* Anorexia, decreased appetite, diabetes mellitus non-insulin-dependent, fluid retention, gout, hypercholesterolemia.

***Musculoskeletal and Connective Tissue Disorders:*** *Frequent:* Muscle spasms, musculoskeletal stiffness, myalgia, neck pain, osteoarthritis, tendonitis. *Inrequent:* Arthritis, aseptic necrosis bone, bone pain, bone spur, bursitis, grain pain, intervertebral disc degeneration, intervertebral disc protrusion, joint stiffness, joint swelling, localized osteoarthritis, monoarthritis, muscle contracture, muscle tightness, muscle twitching, osteoporosis, rotator cuff syndrome, sacroiliitis, synovitis.

***Neoplasms Benign, Malignant, and Unspecified:*** *Inrequent:* Anaplastic thyroid cancer, angiomylipoma, basal cell carcinoma, breast cancer, gastric cancer, gastrointestinal stromal tumor, malignant melanoma, prostate cancer, skin papilloma, squamous cell carcinoma, uterine leiomyoma.

***Nervous System Disorders:*** *Frequent:* Hypoesthesia, migraine. *Inrequent:* Amnesia, aphasia, ataxia, balance disorder, benign intracranial hypertension, burning sensation, carpal tunnel syndrome, disturbance in attention, dizziness postural, dysgnusia, dyskinesia, head discomfort, hyperesthesia, hypersomnia, lethargy, loss of consciousness, memory impairment, migraine with aura, migraine without aura, neuritis, sciatica, sinus headache, sleep apnea syndrome, syncope vasovagal, tension headache, transient ischemic attack, tremor.

***Psychiatric Disorders:*** *Frequent:* Anxiety, depression, irritability, sleep disorder. *Inrequent:* Abnormal dreams, agitation, bruising, confusional state, depressed mood, disorientation, early morning awakening, libido decreased, loss of libido, mood swings, nervousness, nightmares, panic attack, stress symptoms, tension.

***Renal and Urinary Disorders:*** *Inrequent:* Dysuria, hematuria, hypertonic bladder, micturition disorder, nephrolithiasis, nocturia, pollakiuria, proteinuria, urinary retention.

***Reproductive System and Breast Disorders:*** *Frequent:* Erectile dysfunction. *Inrequent:* Breast cyst, dysmenorrhea, menorrhagia, pelvic, peritoneal adhesions, postmenopausal hemorrhage, premenstrual syndrome, prostatitis.

***Respiratory, Thoracic and Mediastinal Disorders:*** *Frequent:* Asthma, pharyngolaryngeal pain. *Inrequent:* Dry throat, dyspnea, epistaxis, hemoptysis, hoarseness, interstitial lung disease, nasal mucosal disorder, nasal polyps, respiratory tract congestion, rhinorrhea, sinus congestion, sneezing, wheezing, yawning.

***Skin and Subcutaneous Tissue Disorders:*** *Frequent:* Night sweats, rash. *Inrequent:* Acne, actinic keratosis, alopecia, cold sores, dermatitis, dermatitis allergic, dermatitis contact, eczema, exanthem, face edema, photosensitivity reaction, pruritis, psoriasis, rash, pruritic skin lesion, urticaria.

***Vascular Disorders:*** *Frequent:* Hot flush, hypertension, hypotension. *Inrequent:* Atherosclerosis, circulatory collapse, flushing, hematoma, thrombosis, varicose vein.

##### DRUG ABUSE AND DEPENDENCE

##### Controlled Substance Class

Ropinirole hydrochloride is not a controlled substance.

##### Physical and Psychological Dependence

Animal studies and human clinical trials with ropinirole hydrochloride did not reveal any potential for drug-seeking behavior or physical dependence.

##### OVERDOSAGE

Of patients who received a dose greater than 24 mg/day, reported symptoms included adverse events commonly reported during dopaminergic therapy (nausea, dizziness), as well as visual hallucinations, hyperhidrosis, claustrophobia, chorea, palpitations, asthenia, and nightmares. Additional symptoms reported for doses of 24 mg or less or for overdoses of unknown amount included vomiting, increased coughing, fatigue, syncope, vasovagal syncope, dyskinesia, agitation, chest pain, orthostatic hypotension, somnolence, and confusional state.

##### Overdose Management

It is anticipated that the symptoms of overdose with ropinirole hydrochloride will be related to its dopaminergic activity. General supportive measures are recommended. Vital signs should be maintained, if necessary. Removal of any unabsorbed material (e.g., by gastric lavage) should be considered.

##### DOUSAGE AND ADMINISTRATION

##### General Dosing Considerations for RLS

Ropinirole hydrochloride tablets can be taken with or without food. Patients may be advised that taking ropinirole hydrochloride tablets with food may reduce the occurrence of nausea. However, this has not been established in controlled clinical trials.

If a significant interruption in therapy with ropinirole hydrochloride has occurred, reinitiation of therapy may be warranted.

##### Geriatric Use

Pharmacokinetic studies demonstrated a reduced clearance of ropinirole in the elderly (see **CLINICAL PHARMACOLOGY**). Dose adjustment is not necessary since the dose is individually titrated to clinical response.

##### Renal Impairment

The pharmacokinetics of ropinirole were not altered in patients with moderate renal impairment (see **CLINICAL PHARMACOLOGY**). Therefore, no dosage adjustment is necessary in patients with moderate renal impairment. The use of ropinirole hydrochloride in patients with severe renal impairment has not been studied.

##### Hepatic Impairment

The pharmacokinetics of ropinirole have not been studied in patients with hepatic impairment. Since patients with hepatic impairment may have higher plasma levels and lower clearance, ropinirole hydrochloride should be titrated with caution in these patients.

##### Dosing for Restless Legs Syndrome

In all clinical trials, the dose for ropinirole hydrochloride was initiated at 0.25 mg once daily, 1 to 3 hours before bedtime. Patients were titrated based on clinical response and tolerability. The recommended adult starting dosage for RLS is 0.25 mg once daily, 1 to 3 hours before bedtime. After 2 days, the dosage can be increased to 0.5 mg once daily and to 1 mg once daily at the end of the first week of dosing, then as shown in **Table 3** as needed to achieve efficacy. For RLS, the safety and effectiveness of doses greater than 4 mg once daily have not been established.

<b>Table 3. Dose Titration Schedule for RLS</b>	
Day/Week	Dosage to be taken once daily, 1 to 3 hours before bedtime
Days 1 and 2	0.25 mg
Days 3 to 7	0.5 mg
Week 2	1 mg
Week 3	1.5 mg
Week 4	2 mg
Week 5	2.5 mg
Week 6	3 mg
Week 7	4 mg

In clinical trials of patients being treated for RLS with doses up to 4 mg once daily, ropinirole hydrochloride was discontinued without a taper.

##### HOW SUPPLIED

Ropinirole hydrochloride tablets are supplied as follows:

0.25 mg are white, film-coated, convex, round tablets, debossed with “93” on one side and “5282” on the other side in bottles of 100 and 1000.

0.5 mg are yellow, film-coated, convex, round tablets, debossed with “93” on one side and “5283” on the other side in bottles of 100 and 1000.

1 mg are green, film-coated, convex, round tablets, debossed with “93” on one side and “5284” on the other side in bottles of 100 and 1000.

2 mg are pink, film-coated, convex, round tablets, debossed with “93” on one side and “5285” on the other side in bottles of 100 and 1000.

3 mg are purple, film-coated, convex, round tablets, debossed with “93” on one side and “5286” on the other side in bottles of 100.

4 mg are beige, film-coated, convex, round tablets, debossed with “93” on one side and “5287” on the other side in bottles of 100.

5 mg are blue, film-coated, convex, round tablets, debossed with “93” on one side and “5288” on the other side in bottles of 100.

##### STORAGE

Protect from light and moisture. Close container tightly after each use.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Dispense in a light, light-resistant container as defined in the USP, with a child-resistant closure (as required).

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008

##### PATIENT INFORMATION

##### ROPINIROLE HYDROCHLORIDE TABLETS

##### ℞ only

**Read this information completely before you start taking ropinirole hydrochloride.**

Read the information each time you get more medicine. There may be new information. This leaflet provides a summary about ropinirole hydrochloride. It does not include everything there is to know about your medicine. This information should not take the place of discussions with your doctor about your medical condition or ropinirole hydrochloride.

##### What is ropinirole hydrochloride?

Ropinirole hydrochloride tablet is a prescription medicine to treat primary moderate-to-severe Restless Legs Syndrome.

##### What is the most important information I should know about ropinirole hydrochloride?

- A lower dose of ropinirole hydrochloride is generally needed for patients with RLS, and is taken once daily before bedtime (see **How should I take ropinirole hydrochloride tablets for RLS?** for the recommended dosing).**

- There are known side effects of ropinirole hydrochloride. If you fall asleep or feel very sleepy while doing normal activities such as driving, faint, feel dizzy, nauseated, or sweaty when you stand up from sitting or lying down, you should talk with your doctor (see **What are the possible side effects of ropinirole hydrochloride?**).

- Before starting ropinirole hydrochloride tablet, be sure to tell your doctor if you are taking any medicines that make you drowsy.

##### Who should not take ropinirole hydrochloride?

You should not take ropinirole hydrochloride tablets if you are allergic to the active ingredient ropinirole or to any of the inactive ingredients. Your doctor and pharmacist have a list of the inactive ingredients.

##### What should I tell my doctor?

Be sure to tell your doctor if:

- you are pregnant or plan to become pregnant.

- you are breast-feeding.
- you have daytime sleepiness from a sleep disorder other than RLS or have unexpected sleepiness or periods of sleep while taking ropinirole hydrochloride.
- you are taking any other prescription or over-the-counter medicines. Some of these medicines may increase your chances of getting side effects while taking ropinirole hydrochloride.

- you start or stop taking other medicines while you are taking ropinirole hydrochloride. This may increase your chances of getting side effects.

- you start or stop smoking while you are taking ropinirole hydrochloride. Smoking may decrease the treatment effect of ropinirole hydrochloride.

- you feel dizzy, nauseated, sweaty, or faint when you stand up from sitting or lying down.
- you drink alcoholic beverages. This may increase your chances of becoming drowsy or sleepy while taking ropinirole hydrochloride.

##### How should I take ropinirole hydrochloride tablets for RLS?

- Be sure to take ropinirole hydrochloride tablets exactly as directed by your doctor or healthcare provider.

- The usual way to take ropinirole hydrochloride tablets is once in the evening, 1 to 3 hours before bedtime.

- Your doctor will start you on a low dose of ropinirole hydrochloride tablets. Your doctor may change the dose until you are taking the amount of medicine that is right for you to control your symptoms.

- If you miss your dose, do not double your next dose.** Take only your usual dose 1 to 3 hours before your next bedtime.

- Contact your doctor, if you stop taking ropinirole hydrochloride tablets for any reason. Do not restart without consulting your doctor.

- You can take ropinirole hydrochloride tablets with or without food. Taking ropinirole hydrochloride with food may decrease the chances of feeling nauseated.

##### What are the possible side effects of ropinirole hydrochloride?

- Most people who take ropinirole hydrochloride tolerate it well. The most commonly reported side effects in people taking ropinirole hydrochloride for RLS are nausea, vomiting, dizziness, and drowsiness or sleepiness. You should be careful until you know if ropinirole hydrochloride affects your ability to remain alert while doing normal daily activities, and you should watch for the development of significant daytime sleepiness or episodes of falling asleep. It is possible that you could fall asleep while doing normal activities such as driving a car, doing physical tasks, or using hazardous machinery while taking ropinirole hydrochloride. Your chances of falling asleep while doing normal activities while taking ropinirole hydrochloride are greater if you are taking other medicines that cause drowsiness.

- When you start taking ropinirole hydrochloride or when you increase your dose, you may feel dizzy, nauseated, sweaty or faint, when first standing up from sitting or lying down. Therefore, do not stand up quickly after sitting or lying down, particularly if you have been sitting or lying down for a long period of time. Take a minute sitting on the edge of the bed or chair before you get up.

- Hallucinations (unreal sounds, visions, or sensations) have been reported in patients taking ropinirole hydrochloride. These were uncommon in patients taking ropinirole hydrochloride for RLS. The risk is greater in patients taking ropinirole hydrochloride with L-dopa, or taking higher doses of ropinirole hydrochloride than recommended for RLS.

This is not a complete list of side effects and should not take the place of discussions with your healthcare providers. Your doctor or pharmacist can give you a more complete list of possible side effects. Talk to your doctor about any side effects or problems you may have.

##### Other information about ropinirole hydrochloride

Patients being treated with ropinirole hydrochloride should have periodic skin examinations for melanoma.

- Take ropinirole hydrochloride exactly as your doctor prescribes it.
- Do not share ropinirole hydrochloride with other people, even if they have the same symptoms you have.
- Keep ropinirole hydrochloride out of the reach of children.
- Store ropinirole hydrochloride at room temperature out of direct sunlight.
- Keep ropinirole hydrochloride in a tightly closed container.

This leaflet summarizes important information about ropinirole hydrochloride. Medicines are sometimes prescribed for purposes other than those listed in this leaflet. Do not take ropinirole hydrochloride for a condition for which it was not prescribed. For more information, talk with your doctor or pharmacist. They can give you information about ropinirole hydrochloride that is written for healthcare professionals.

Manufactured By:  
**TEVA PHARMACEUTICALS USA**  
Sellersville, PA 18960

Iss. 5/2008