









EXPIRES tor oral suspension) (mycophenolate mofetil

Oral Suspension [®] tqə Əllə Ə

NDC 0004-0591-59

NDC 0004-0261-29

CellCept® Oral Suspension

(mycophenolate mofetil for oral suspension)

200 mg/mL

Net contents after constitution: 175 mL, containing 35.0 grams mycophenolate mofetil. Deliverable volume after constitution: 160-165 mL.

Phenylketonurics: Contains 0.56 mg phenylalanine per mL after constitution. Usual dosage: For dosage recommendations, handling and other important prescribing information, read accompanying insert.

Store dry powder at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F).

Store constituted suspension at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). Storage in a refrigerator at 2°-8°C (36°-46°F) is acceptable.

Do not freeze. Shake closed bottle well before using. Discard any unused portion 60 days after constitution.

To the pharmacist:

Tap the closed bottle several times to loosen the powder. Measure 94 mL of water in a graduated cylinder.

Add approximately half the total amount of water for constitution to the bottle and shake the closed bottle well for about 1 minute.

Add the remainder of water and shake the closed bottle well for about 1 minute.

Remove the child-resistant cap and push bottle adapter into neck of bottle.

Close bottle with child-resistant cap tightly. This will assure the proper seating of the bottle adapter in the bottle, and child-resistant status of the cap. Dispense with patient instructions

and oral dispensers. See enclosed patient instructions for use of oral dispenser.

Contains 225 mL bottle with bottle adapter and 2 oral dispensers.

Made in Ireland Distributed by: **Roche Laboratories Inc.** Nutley, New Jersey 07110

CellCept® Oral Suspension

(mycophenolate mofetil

Each mL contains 200 mg mycophenolate mofetil after constitution.

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient. For additional Medication **Guides call 1-800-617-8191 or visit** www.rocheusa.com/products/CellCept.



NDC 0004-0261-29

for oral suspension)

200 mg/mL




Ronly


27899633(A)








27899637-L (D) 26Jan09:Layout 1 1/29/09 1:48 PM Page 1



may not be the latest approve ormation, please visit https://v 27899624 L (D) 22Jan09: NDC 0004-0298-30 CellCept® Intravenous (mycophenolate mofetil hydrochloride for injection) 500 ma Ronly FOR INTRAVENOUS INFUSION ONLY Each vial contains the equivalent of 500 mg mycophenolate mofetil (as the hydrochloride salt). 25 mg pólysorbate 80, and 5 mg citric acid. USUAL DOSAGE: See package insert. Attention Pharmacist: Dispense the accompanying Medication Guide to each patient. Store powder at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). Made in Ireland Distributed by: Roche Laboratories Inc. Nutley, New Jersey 07110 i-Wate-WWW (01) 103 0004 0298 30 4 3000553B


This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda

27899632-L (D) 23Jan09:27899207-L-(D) 1/29/09 1:57 PM Page 1




This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda

27899626-L (D) 26Jan09:27899205-L (D) 01May07 1/29/09 1:25 PM Page



27899630-L (D) 23Jan09:27899204-L (D) 1/29/09 1:31 PM Page 1



Store dry powder at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). Store constituted suspension at 25°C (77°F); excursions permitted to 15°-30°C (59°-86°F). Storage in a refrigerator at 2°-8°C (36°-46°F) is acceptable. Do not freeze.

Shake closed bottle well before using. Discard any unused portion 60 days after constitution.

Constitute Date:

27899634

NDC 0004-0261-29

CellCept® Oral Suspension

(mycophenolate mofetil for oral suspension)

200 mg/mL

Each mL contains 200 mg mycophenolate mofetil after constitution.

Attention Pharmacist: Dispense the accompanying Medication Guide to each patient.




Usual dosage: See package insert. Net contents after constitution: 175 mL, containing 35.0 grams mycophenolate mofetil. Deliverable volume after constitution: 160-165 mL. Phenylketonurics: Contains 0.56 mg phenylalanine per mL after constitution.


To the pharmacist: Tap the closed bottle several times to loosen the powder. Measure 94 mL of water in a graduated cylinder. Add approximately half the total amount of water for constitution to the bottle and shake the closed bottle well for about 1 minute. Add the remainder of water and shake the closed bottle well for about 1 minute. Remove the childresistant cap and push bottle adapter into neck of bottle. Close bottle with child-resistant cap tightly. This will assure the proper seating of the bottle adapter in the bottle, and child-resistant status of the cap. Dispense with patient instructions and



Made in Ireland Distributed by: Roche Laboratories Inc. Nutley, New Jersey 07110

oral dispensers. See enclosed patient instructions for use of oral dispenser. 27899628-L (D) 26Jan09:Layout 1 1/29/09 1:29 PM Page 1





Ozlem Belen

2/27/2009 05:13:38 PM