

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.  
Dissolves in any beverage  
Distributed by Kremers Urban, LLC, Princeton, NJ 08540, USA

### Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

#### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

**Questions or comments?**  
1-800-477-7877

NDC 62175-195-07

Original  
Prescription  
Strength

## Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool

**CAP** is not a dosing **CUP**.

Use dosing **CUP** provided to measure dose.

NET WT 4.1 OZ (119 g) 7 ONCE-DAILY DOSES

### Drug Facts

| Active ingredient (in each dose)                               | Purpose  |
|--|----------|
| Polyethylene Glycol 3350, 17 g (dosing cup filled to line).... | Laxative |

#### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

#### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

#### Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product** you may have loose, watery, more frequent stools

#### Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

4010359 Rev. 04/09

Exp.:

Lot.:



Dissolves in Any Beverage

### Drug Facts (continued)

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877

NDC 62175-195-15

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

\* Relieves Occasional Constipation

\* Softens Stool

**CAP** is not a dosing **CUP**.  
Use dosing **CUP** provided to measure dose.

NET WT 8.9 OZ (255 g)

15 ONCE-DAILY DOSES

Distributed by Kremers Urban, LLC, Princeton, NJ 08540 USA

Polyethylene Glycol 3350 Powder for Solution increases frequency of bowel movements and softens the stool.

### Drug Facts

| Active ingredient (in each dose)                               | Purpose  |
|--|----------|
| Polyethylene Glycol 3350, 17 g (dosing cup filled to line).... | Laxative |

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



Lot: Exp:

4010420 Rev. 04/09

Dissolves in Any Beverage

NDC 62175-195-31

Original  
Prescription  
Strength

Polyethylene Glycol 3350 Powder for Solution  
increases frequency of bowel movements  
and softens the stool.

### Drug Facts (continued)

If **pregnant or breast feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not take more than directed unless advised by your doctor
- the dosing cup supplied is marked to contain 17 grams of powder when filled to the indicated line
- adults and children 17 years of age and older:
  - fill to indicated line for the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877

Distributed by Kremers Urban, LLC, Princeton, NJ 08540 USA

# Polyethylene Glycol 3350 Powder for Solution, Laxative

\* *Relieves Occasional Constipation*

\* *Softens Stool*

**CAP** is not a dosing **CUP**.

Use dosing **CUP** provided to measure dose.

**NET WT 18.5 OZ (527 g) 31 ONCE-DAILY DOSES**

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

### Drug Facts

| Active ingredient (in each dose)                              | Purpose  |
|---|----------|
| Polyethylene Glycol 3350, 17 g (dosing cup filled to line)... | Laxative |

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are** taking a prescription drug

**When using this product** you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



Lot: Exp:

4010421  
Rev. 04/09

FOLD AT LINE  
TEAR AT ARROW

NDC 62175-195-01

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* *Relieves Occasional Constipation*
- \* *Softens Stool*

NET WT 0.5 OZ (17g)

1 Dose

## Drug Facts

**Active ingredient (in each dose)** Purpose  
Polyethylene Glycol 3350, 17 g.....Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

#### Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product** you may have loose, watery, more frequent stools

#### Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

## Drug Facts (continued)

**If pregnant or breast feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- **do not take more than directed unless advised by your doctor**
- adults and children 17 years of age and older:
  - stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - use once daily
  - use no more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20° - 25°C (68° - 77°F)
- tamper-evident: do not use if foil is open or broken

**Inactive ingredients** none

### Questions or comments?

1-800-477-7877



Distributed by Kremers Urban, LLC  
Princeton, NJ 08540 USA L5480 Rev. 04/09

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

NDC 62175-195-10

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage

10 PACKETS, EACH  
PACKET NET WT 0.5 OZ (17g)

**10**  
ONCE-DAILY DOSES

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage

CONTAINS 10 ONCE-DAILY DOSES  
EACH PACKET CONTAINS ONE DOSE NET WT 0.5 OZ (17G)

Original  
Prescription  
Strength

# Polyethylene Glycol 3350 Powder for Solution, Laxative

- \* Relieves Occasional Constipation
- \* Softens Stool
- \* Dissolves in Any Beverage



**Drug Facts (continued)**  
If pregnant or breast feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ do not take more than directed unless advised by your doctor  
 ■ adults and children 17 years of age and older:  
 ■ stir and dissolve one packet of powder (17 g) in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink  
 ■ use once daily  
 ■ use no more than 7 days  
 ■ children 16 years of age or under: ask a doctor

**Other information**  
 ■ store at 20° - 25°C (68° - 77°F)  
 ■ tamper-evident: do not use if foil is open or broken

**Inactive ingredients** none

**Questions or comments?**  
1-800-477-7877

**Drug Facts**  
**Active ingredient (in each dose)** Polyethylene Glycol 3350, 17 g.....  
**Purpose** Laxative

**Use**  
 ■ relieves occasional constipation (irregularity)  
 ■ generally produces a bowel movement in 1 to 3 days

**Warnings**  
**Allergy alert:** Do not use if you are allergic to polyethylene glycol  
**Do not use** if you have kidney disease, except under the advice and supervision of a doctor  
**Ask a doctor before use if you have**  
 ■ nausea, vomiting or abdominal pain  
 ■ a sudden change in bowel habits that lasts over 2 weeks  
 ■ irritable bowel syndrome  
**Ask a doctor or pharmacist before use if you are** taking a prescription drug  
**When using this product** you may have loose, watery, more frequent stools  
**Stop use and ask a doctor if**  
 ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.  
 ■ you get diarrhea  
 ■ you need to use a laxative for longer than 1 week

Unvarnished Area

Control No.  
Exp. Date



Barcode Area



CR5481  
Rev. 04/09