



PEGASYS®

(peginterferon alfa-2a)

Rx only

Alpha interferons, including PEGASYS (peginterferon alfa-2a), may cause or aggravate fatal or life-threatening neuropsychiatric, autoimmune, ischemic, and infectious disorders. Patients should be monitored closely with periodic clinical and laboratory evaluations. Therapy should be withdrawn in patients with persistently severe or worsening signs or symptoms of these conditions. In many, but not all cases, these disorders resolve after stopping PEGASYS therapy (see WARNINGS and ADVERSE REACTIONS).

Use with Ribavirin. Ribavirin, including COPEGUS®, may cause birth defects and/or death of the fetus. Extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients. Ribavirin causes hemolytic anemia. The anemia associated with ribavirin therapy may result in a worsening of cardiac disease. Ribavirin is genotoxic and mutagenic and should be considered a potential carcinogen (see COPEGUS Package Insert for additional information and other WARNINGS).

DESCRIPTION

PEGASYS, peginterferon alfa-2a, is a covalent conjugate of recombinant alfa-2a interferon (approximate molecular weight [MW] 20,000 daltons) with a single branched bis-monomethoxy polyethylene glycol (PEG) chain (approximate MW 40,000 daltons). The PEG moiety is linked at a single site to the interferon alfa moiety via a stable amide bond to lysine. Peginterferon alfa-2a has an approximate molecular weight of 60,000 daltons. Interferon alfa-2a is produced using recombinant DNA technology in which a cloned human leukocyte interferon gene is inserted into and expressed in *Escherichia coli*.

PEGASYS is supplied as an injectable solution in vials and prefilled syringes.

180 µg/1.0 mL Vial: A vial contains approximately 1.2 mL of solution to deliver 1.0 mL of drug product. Subcutaneous (sc) administration of 1.0 mL delivers 180 µg of drug product (expressed as the amount of interferon alfa-2a), 8.0 mg sodium chloride, 0.05 mg polysorbate 80, 10.0 mg benzyl alcohol, 2.62 mg sodium acetate trihydrate, and 0.05 mg acetic acid. The solution is colorless to light yellow and the pH is 6.0 ± 0.5.

180 µg/0.5 mL Prefilled Syringe: Each syringe contains 0.6 mL of solution to deliver 0.5 mL of drug product. Subcutaneous (sc) administration of 0.5 mL delivers 180 µg of drug product (expressed as the amount of interferon alfa-2a), 4.0 mg sodium chloride, 0.025 mg polysorbate 80, 5.0 mg benzyl alcohol, 1.3085 mg sodium acetate trihydrate, and 0.0231 mg acetic acid. The solution is colorless to light yellow and the pH is 6.0 ± 0.5.

PEGASYS® (peginterferon alfa-2a)

CLINICAL PHARMACOLOGY

Pharmacodynamics

Interferons bind to specific receptors on the cell surface initiating intracellular signaling via a complex cascade of protein-protein interactions leading to rapid activation of gene transcription. Interferon-stimulated genes modulate many biological effects including the inhibition of viral replication in infected cells, inhibition of cell proliferation and immunomodulation. The clinical relevance of these in vitro activities is not known.

PEGASYS stimulates the production of effector proteins such as serum neopterin and 2', 5'-oligoadenylate synthetase.

Pharmacokinetics

Maximal serum concentrations (C_{max}) and AUC increased in a nonlinear dose related manner following administration of 90 to 270 µg of PEGASYS. Maximal serum concentrations (C_{max}) occur between 72 to 96 hours post-dose.

Week 48 mean trough concentrations (16 ng/mL; range 4 to 28) at 168 hours post-dose are approximately 2-fold higher than week 1 mean trough concentrations (9 ng/mL; range 0 to 15). Steady-state serum levels are reached within 5 to 8 weeks of once weekly dosing. The peak to trough ratio at week 48 is approximately 2. The mean systemic clearance in healthy subjects given PEGASYS was 94 mL/h, which is approximately 100-fold lower than that for interferon alfa-2a (ROFERON®-A). The mean terminal half-life after sc dosing in patients with chronic hepatitis C was 160 hours (range 84 to 353 hours) compared to 5 hours (range 3.7 to 8.5 hours) for ROFERON-A.

Special Populations

Gender and Age

PEGASYS administration yielded similar pharmacokinetics in male and female healthy subjects. The AUC was increased from 1295 to 1663 ng·h/mL in subjects older than 62 years taking 180 µg PEGASYS, but peak concentrations were similar (9 vs. 10 ng/mL) in those older and younger than 62 years.

Pediatric Patients

In a population pharmacokinetics study, 14 children 2 to 8 years of age with CHC received PEGASYS based on their body surface area (BSA of the child \times 180 µg/1.73m²). The clearance of PEGASYS in children was nearly 4-fold lower compared to the clearance reported in adults.

Steady-state trough levels in children with the BSA-adjusted dosing were similar to trough levels observed in adults with 180 µg fixed dosing. Time to reach the steady state in children is approximately 12 weeks, whereas in adults, steady state is reached within 5 to 8 weeks. In these children receiving the BSA adjusted dose, the mean exposure (AUC) during the dosing interval is predicted to be 25% to 70% higher than that observed in adults receiving 180 µg fixed dosing. The safety and effectiveness of PEGASYS in patients below the age of 18 years have not been established (see **PRECAUTIONS: Pediatric Use**).

PEGASYS® (peginterferon alfa-2a)

80 Renal Dysfunction

81 In patients with end stage renal disease undergoing hemodialysis, there is a 25% to 45%
82 reduction in PEGASYS clearance (see **PRECAUTIONS: Renal Impairment**).

83 The pharmacokinetics of ribavirin following administration of COPEGUS have not been
84 studied in patients with renal impairment and there are limited data from clinical trials on
85 administration of COPEGUS in patients with creatinine clearance <50 mL/min.
86 Therefore, patients with creatinine clearance <50 mL/min should not be treated with
87 COPEGUS (see **WARNINGS** and **DOSAGE AND ADMINISTRATION**).

88 Effect of Food on Absorption of Ribavirin

89 Bioavailability of a single oral dose of ribavirin was increased by co-administration with
90 a high-fat meal. The absorption was slowed (T_{max} was doubled) and the AUC_{0-192h} and
91 C_{max} increased by 42% and 66%, respectively, when COPEGUS was taken with a high-
92 fat meal compared with fasting conditions (see **DOSAGE AND ADMINISTRATION**).

93 Drug Interactions

94 Nucleoside Analogues

95 In vitro data indicate ribavirin reduces phosphorylation of lamivudine, stavudine, and
96 zidovudine. However, no pharmacokinetic (e.g., plasma concentrations or intracellular
97 triphosphorylated active metabolite concentrations) or pharmacodynamic (e.g., loss of
98 HIV/HCV virologic suppression) interaction was observed when ribavirin and
99 lamivudine (n=18), stavudine (n=10), or zidovudine (n=6) were co-administered as part
100 of a multi-drug regimen to HCV/HIV coinfecting patients (see **PRECAUTIONS: Drug**
101 **Interactions**).

102 In vitro, didanosine or its active metabolite (dideoxyadenosine 5'-triphosphate) is
103 increased when didanosine is co-administered with ribavirin (see **PRECAUTIONS:**
104 **Drug Interactions**).

105 Drugs Metabolized by Cytochrome P450

106 There was no effect on the pharmacokinetics of representative drugs metabolized by CYP
107 2C9, CYP 2C19, CYP 2D6 or CYP 3A4.

108 Treatment with PEGASYS once weekly for 4 weeks in healthy subjects was associated
109 with an inhibition of P450 1A2 and a 25% increase in theophylline AUC (see
110 **PRECAUTIONS: Drug Interactions**).

111 Methadone

112 The pharmacokinetics of concomitant administration of methadone and PEGASYS were
113 evaluated in 24 PEGASYS naive chronic hepatitis C (CHC) patients (15 male, 9 female)
114 who received 180 µg PEGASYS subcutaneously weekly. All patients were on stable
115 methadone maintenance therapy (median dose 95 mg, range 30 mg to 150 mg) prior to
116 receiving PEGASYS. Mean methadone PK parameters were 10% to 15% higher after 4
117 weeks of PEGASYS treatment as compared to baseline (see **PRECAUTIONS: Drug**
118 **Interactions**). Methadone did not significantly alter the PK of PEGASYS as compared to
119 a PK study of 6 chronic hepatitis C patients not receiving methadone.

PEGASYS® (peginterferon alfa-2a)

CLINICAL STUDIES

Chronic Hepatitis C Studies 1, 2, and 3: PEGASYS Monotherapy

The safety and effectiveness of PEGASYS for the treatment of hepatitis C virus infection were assessed in three randomized, open-label, active-controlled clinical studies. All patients were adults, had compensated liver disease, detectable hepatitis C virus (HCV), liver biopsy diagnosis of chronic hepatitis, and were previously untreated with interferon. All patients received therapy by sc injection for 48 weeks, and were followed for an additional 24 weeks to assess the durability of response. In studies 1 and 2, approximately 20% of subjects had cirrhosis or bridging fibrosis. Study 3 enrolled patients with a histological diagnosis of cirrhosis (78%) or bridging fibrosis (22%).

In Study 1 (n=630), patients received either ROFERON-A (interferon alfa-2a) 3 MIU three times/week (tiw), PEGASYS 135 µg once each week (qw) or PEGASYS 180 µg qw. In Study 2 (n=526), patients received either ROFERON-A 6 MIU tiw for 12 weeks followed by 3 MIU tiw for 36 weeks or PEGASYS 180 µg qw. In Study 3 (n=269), patients received ROFERON-A 3 MIU tiw, PEGASYS 90 µg qw or PEGASYS 180 µg once each week.

In all three studies, treatment with PEGASYS 180 µg resulted in significantly more patients who experienced a sustained response (defined as undetectable HCV RNA [<50 IU/mL] using the COBAS AMPLICOR® HCV Test, version 2.0 and normalization of ALT on or after study week 68) compared to treatment with ROFERON-A. In Study 1, response to PEGASYS 135 µg was not different from response to 180 µg. In Study 3, response to PEGASYS 90 µg was intermediate between PEGASYS 180 µg and ROFERON-A.

Table 1 Sustained Response to Monotherapy Treatment

| | Study 1 | | | Study 2 | | | Study 3 | | |
|--|-------------------------------|------------------------------|-------------------|---------------------------------|------------------------------|-------------------|------------------------------|-----------------------------|-------------------|
| | ROFERON-A 3 MIU (N=207) | PEGASYS 180 µg (N=208) | DIFF* (95% CI) | ROFERON-A 6/3 MIU (N=261) | PEGASYS 180 µg (N=265) | DIFF* (95% CI) | ROFERON-A 3 MIU (N=86) | PEGASYS 180 µg (N=87) | DIFF* (95% CI) |
| Combined Virologic and Biologic Sustained Response | 11% | 24% | 13 (6, 20) | 17% | 35% | 18 (11, 25) | 7% | 23% | 16 (6, 26) |
| Sustained Virologic Response | 11% | 26% | 15 (8, 23) | 19% | 38% | 19 (11, 26) | 8% | 30% | 22 (11, 33) |

*Percent difference between PEGASYS and ROFERON-A treatment.

Matched pre- and post-treatment liver biopsies were obtained in approximately 70% of patients. Similar modest reductions in inflammation compared to baseline were observed in all treatment groups.

Of the patients who did not demonstrate either undetectable HCV RNA or at least a 2log₁₀ drop in HCV RNA titer from baseline by 12 weeks of PEGASYS 180 µg therapy,

PEGASYS® (peginterferon alfa-2a)

2% (3/156) achieved a sustained virologic response (see **DOSAGE AND ADMINISTRATION**).

Averaged over Study 1, Study 2, and Study 3, response rates to PEGASYS were 23% among patients with viral genotype 1 and 48% in patients with other viral genotypes. The treatment response rates were similar in men and women.

Chronic Hepatitis C Studies 4 and 5: PEGASYS/COPEGUS Combination Therapy

The safety and effectiveness of PEGASYS in combination with COPEGUS for the treatment of hepatitis C virus infection were assessed in two randomized controlled clinical trials. All patients were adults, had compensated liver disease, detectable hepatitis C virus, liver biopsy diagnosis of chronic hepatitis, and were previously untreated with interferon. Approximately 20% of patients in both studies had compensated cirrhosis (Child-Pugh class A). Patients coinfectd with HIV were excluded from these studies.

In Study 4, patients were randomized to receive either PEGASYS 180 µg sc once weekly (qw) with an oral placebo, PEGASYS 180 µg qw with COPEGUS 1000 mg po (body weight <75 kg) or 1200 mg po (body weight ≥75 kg) or REBETRON® (interferon alfa-2b 3 MIU sc tiw plus ribavirin 1000 mg or 1200 mg po). All patients received 48 weeks of therapy followed by 24 weeks of treatment-free follow-up. COPEGUS or placebo treatment assignment was blinded. Sustained virological response was defined as undetectable (<50 IU/mL) HCV RNA on or after study week 68. PEGASYS in combination with COPEGUS resulted in a higher SVR compared to PEGASYS alone or interferon alfa-2b and ribavirin (**Table 2**). In all treatment arms, patients with viral genotype 1, regardless of viral load, had a lower response rate.

Table 2 Sustained Virologic Response to Combination Therapy (Study 4)

| | Interferon alfa-2b + Ribavirin 1000 mg or 1200 mg | PEGASYS + Placebo | PEGASYS + COPEGUS 1000 mg or 1200 mg |
|---------------|--|----------------------|---|
| All patients | 197/444 (44%)* | 65/224 (29%) | 241/453 (53%)* |
| Genotype 1 | 103/285 (36%) | 29/145 (20%) | 132/298 (44%) |
| Genotypes 2-6 | 94/159 (59%) | 36/79 (46%) | 109/155 (70%) |

*Difference in overall treatment response (PEGASYS/COPEGUS – Interferon alfa-2b/ribavirin) was 9% (95% CI 2.3, 15.3).

In Study 5 (see **Table 3**), all patients received PEGASYS 180 µg sc qw and were randomized to treatment for either 24 or 48 weeks and to a COPEGUS dose of either 800 mg or 1000 mg/1200 mg (for body weight <75 kg / ≥75 kg). Assignment to the four treatment arms was stratified by viral genotype and baseline HCV viral titer. Patients with genotype 1 and high viral titer (defined as >2 x 10⁶ HCV RNA copies/mL serum) were preferentially assigned to treatment for 48 weeks.

PEGASYS® (peginterferon alfa-2a)

185 HCV Genotypes

186 HCV 1 and 4 – Irrespective of baseline viral titer, treatment for 48 weeks with
187 PEGASYS and 1000 mg or 1200 mg of COPEGUS resulted in higher SVR (defined as
188 undetectable HCV RNA at the end of the 24-week treatment-free follow-up period)
189 compared to shorter treatment (24 weeks) and/or 800 mg COPEGUS.

190 HCV 2 and 3 – Irrespective of baseline viral titer, treatment for 24 weeks with
191 PEGASYS and 800 mg of COPEGUS resulted in a similar SVR compared to longer
192 treatment (48 weeks) and/or 1000 mg or 1200 mg of COPEGUS (see **Table 3**).

193 The numbers of patients with genotype 5 and 6 were too few to allow for meaningful
194 assessment.

195 **Table 3 Sustained Virologic Response as a Function of Genotype**
196 **(Study 5)**

| | 24 Weeks Treatment | | 48 Weeks Treatment | |
|-----------------------|---|--|---|--|
| | PEGASYS + COPEGUS 800 mg (N=207) | PEGASYS + COPEGUS 1000 mg or 1200 mg* (N=280) | PEGASYS + COPEGUS 800 mg (N=361) | PEGASYS + COPEGUS 1000 mg or 1200 mg* (N=436) |
| Genotype 1 | 29/101 (29%) | 48/118 (41%) | 99/250 (40%) | 138/271 (51%) |
| Genotypes 2, 3 | 79/96 (82%) | 116/144 (81%) | 75/99 (76%) | 117/153 (76%) |
| Genotype 4 | 0/5 (0%) | 7/12 (58%) | 5/8 (63%) | 9/11 (82%) |

197 *1000 mg for body weight <75 kg; 1200 mg for body weight ≥75 kg.

198 Other Treatment Response Predictors

199 Treatment response rates are lower in patients with poor prognostic factors receiving
200 pegylated interferon alpha therapy. In studies 4 and 5, treatment response rates were
201 lower in patients older than 40 years (50% vs. 66%), in patients with cirrhosis (47% vs.
202 59%), in patients weighing over 85 kg (49% vs. 60%), and in patients with genotype 1
203 with high vs. low viral load (43% vs. 56%). African-American patients had lower
204 response rates compared to Caucasians.

205 Paired liver biopsies were performed on approximately 20% of patients in studies 4 and
206 5. Modest reductions in inflammation compared to baseline were seen in all treatment
207 groups.

208 In studies 4 and 5, lack of early virologic response by 12 weeks (defined as HCV RNA
209 undetectable or >2log₁₀ lower than baseline) was grounds for discontinuation of
210 treatment. Of patients who lacked an early viral response by 12 weeks and completed a
211 recommended course of therapy despite a protocol-defined option to discontinue therapy,
212 5/39 (13%) achieved an SVR. Of patients who lacked an early viral response by 24
213 weeks, 19 completed a full course of therapy and none achieved an SVR.

PEGASYS® (peginterferon alfa-2a)

Chronic Hepatitis C and Coinfection with HIV (CHC/HIV) Study 6: PEGASYS Monotherapy and PEGASYS/COPEGUS Combination Therapy

In Study 6, patients with CHC/HIV were randomized to receive either PEGASYS 180 µg sc once weekly (qw) plus an oral placebo, PEGASYS 180 µg qw plus COPEGUS 800 mg po daily or ROFERON-A (interferon alfa-2a), 3 MIU sc tiw plus COPEGUS 800 mg po daily. All patients received 48 weeks of therapy and sustained virologic response (SVR) was assessed at 24 weeks of treatment-free follow-up. COPEGUS or placebo treatment assignment was blinded in the PEGASYS treatment arms. All patients were adults, had compensated liver disease, detectable hepatitis C virus, liver biopsy diagnosis of chronic hepatitis C, and were previously untreated with interferon. Patients also had CD4+ cell count ≥200 cells/µL or CD4+ cell count ≥100 cells/µL but <200 cells/µL and HIV-1 RNA <5000 copies/mL, and stable status of HIV. Approximately 15% of patients in the study had cirrhosis. Results are shown in Table 4.

Table 4 Sustained Virologic Response in Patients with Chronic Hepatitis C Coinfected with HIV (Study 6)

| | ROFERON-A + COPEGUS 800 mg (N=289) | PEGASYS + Placebo (N=289) | PEGASYS + COPEGUS 800 mg (N=290) |
|----------------|--|---------------------------------|--|
| All patients | 33 (11%)* | 58 (20%)* | 116 (40%) |
| Genotype 1 | 12/171 (7%) | 24/175 (14%) | 51/176 (29%) |
| Genotypes 2, 3 | 18/89 (20%) | 32/90 (36%) | 59/95 (62%) |

*PEGASYS + COPEGUS vs. PEGASYS; PEGASYS + COPEGUS vs. ROFERON-A + COPEGUS p-value <0.0001 (Cochran-Mantel-Haenszel).

Treatment response rates are lower in CHC/HIV patients with poor prognostic factors (including HCV genotype 1, HCV RNA >800,000 IU/mL, and cirrhosis) receiving pegylated interferon alpha therapy. Geographic region is not a prognostic factor for response. However, poor prognostic factors occur more frequently in the US population than in the non-US population.

Of the patients who did not demonstrate either undetectable HCV RNA or at least a 2log₁₀ reduction from baseline in HCV RNA titer by 12 weeks of PEGASYS and COPEGUS combination therapy, 2% (2/85) achieved an SVR.

In CHC patients with HIV coinfection who received 48 weeks of PEGASYS alone or in combination with COPEGUS treatment, mean and median HIV RNA titers did not increase above baseline during treatment or 24 weeks post-treatment.

Chronic Hepatitis B Studies 7 and 8: PEGASYS Monotherapy

The safety and effectiveness of PEGASYS for the treatment of chronic hepatitis B were assessed in controlled clinical trials in HBeAg positive (Study 7) and HBeAg negative (Study 8) patients with chronic hepatitis B.

PEGASYS® (peginterferon alfa-2a)

247 Patients were randomized to PEGASYS 180 µg sc once weekly (qw), PEGASYS 180 µg
248 sc qw combined with lamivudine 100 mg once daily po or lamivudine 100 mg once daily
249 po. All patients received 48 weeks of their assigned therapy followed by 24 weeks of
250 treatment-free follow-up. Assignment to receipt of PEGASYS or no PEGASYS was not
251 masked.

252 All patients were adults with compensated liver disease, had chronic hepatitis B virus
253 (HBV) infection, and evidence of HBV replication (serum HBV >500,000 copies/mL for
254 Study 7 and >100,000 copies/mL for Study 8) as measured by PCR (COBAS
255 AMPLICOR® HBV Assay). All patients had serum alanine aminotransferase (ALT)
256 between 1 and 10 times the upper limit of normal (ULN) and liver biopsy findings
257 compatible with the diagnosis of chronic hepatitis.

258 The results observed in the PEGASYS and lamivudine monotherapy groups are shown in
259 Table 5.

260 **Table 5 Percentage of Patients with Serological, Virological,**
261 **Biochemical, and Histological Response**

| | Study 7 HBeAg positive | | | Study 8 HBeAg negative | | |
|---|---------------------------|------------------|--------------------|---------------------------|------------------|--------------------|
| | Lamivudine N = 272 | | PEGASYS N = 271 | Lamivudine N = 181 | | PEGASYS N = 177 |
| | EOT ¹ | EOF ² | EOF ² | EOT ¹ | EOF ² | EOF ² |
| HBeAg Seroconversion (%) | 20 | 19* | 32* | NA | NA | NA |
| HBV DNA Response (%) ³ | 62 | 22*** | 32*** | 85 | 29** | 43** |
| ALT Normalization (%) | 62 | 28 | 41 | 73 | 44** | 59** |
| HBsAg Seroconversion (%) | 0 | 0 | 3 | 1 | 0 | 3 |
| | N = 184 | | N = 207 | N = 125 | | N = 143 |
| Histological Improvement (%) ⁴ | ND | 40 | 41 | ND | 41 | 48 |
| Changes in Ishak fibrosis score compared to baseline (%): | | | | | | |
| - Improved ⁵ | ND | 32 | 25 | ND | 31 | 32 |
| - Unchanged | | 20 | 25 | | 23 | 30 |
| - Worsened ⁵ | | 16 | 26 | | 15 | 19 |

262 ¹End of Treatment (week 48)

263 ²End of follow-up – 24 weeks post-treatment (week 72)

PEGASYS® (peginterferon alfa-2a)

264 ³<100,000 copies/mL for HBeAg positive and <20,000 copies/mL for HBeAg negative patients

265 ⁴≥2 point decrease in Ishak necro-inflammatory score from baseline with no worsening of the Ishak fibrosis
266 score. Not all patients provided both initial and end of follow-up biopsies (missing biopsy rates: 19% to
267 24% in the PEGASYS and 31% to 32% in the Lamivudine arms)

268 ⁵Change of 1 point or more in Ishak fibrosis score

269 *p<0.001; **p<0.01; ***p=0.012 (primary efficacy endpoints Cochran-Mantel-Haenszel test comparisons
270 of PEGASYS to Lamivudine)

271

272 PEGASYS co-administered with lamivudine did not result in any additional sustained
273 response when compared to PEGASYS monotherapy.

274 Conclusions regarding comparative efficacy of PEGASYS and lamivudine treatment
275 based upon the end of follow-up results are limited by the different mechanisms of action
276 of the two compounds. Most treatment effects of lamivudine are unlikely to persist 24
277 weeks after therapy is withdrawn.

278 INDICATIONS AND USAGE

279 PEGASYS, peginterferon alfa-2a, alone or in combination with COPEGUS, is indicated
280 for the treatment of adults with chronic hepatitis C virus infection who have compensated
281 liver disease and have not been previously treated with interferon alpha. Patients in whom
282 efficacy was demonstrated included patients with compensated liver disease and
283 histological evidence of cirrhosis (Child-Pugh class A) and patients with HIV disease that
284 is clinically stable (e.g., antiretroviral therapy not required or receiving stable
285 antiretroviral therapy).

286 PEGASYS is indicated for the treatment of adult patients with HBeAg positive and
287 HBeAg negative chronic hepatitis B who have compensated liver disease and evidence of
288 viral replication and liver inflammation.

289 CONTRAINDICATIONS

290 PEGASYS is contraindicated in patients with:

- 291 • Hypersensitivity to PEGASYS or any of its components
- 292 • Autoimmune hepatitis
- 293 • Hepatic decompensation (Child-Pugh score greater than 6 [class B and C]) in
294 cirrhotic patients before or during treatment
- 295 • Hepatic decompensation with Child-Pugh score greater than or equal to 6 in cirrhotic
296 CHC patients coinfectd with HIV before or during treatment

297 PEGASYS is contraindicated in neonates and infants because it contains benzyl alcohol.
298 Benzyl alcohol is associated with an increased incidence of neurologic and other
299 complications in neonates and infants, which are sometimes fatal.

300 PEGASYS and COPEGUS combination therapy is additionally contraindicated in:

- 301 • Patients with known hypersensitivity to COPEGUS or to any component of the tablet

PEGASYS® (peginterferon alfa-2a)

- 302 • Women who are pregnant
- 303 • Men whose female partners are pregnant
- 304 • Patients with hemoglobinopathies (e.g., thalassemia major, sickle-cell anemia)

305 **WARNINGS**

306 **General**

307 Patients should be monitored for the following serious conditions, some of which may
308 become life threatening. Patients with persistently severe or worsening signs or
309 symptoms should have their therapy withdrawn (see **BOXED WARNING**).

310 **Neuropsychiatric**

311 Life-threatening or fatal neuropsychiatric reactions may manifest in patients receiving
312 therapy with PEGASYS and include suicide, suicidal ideation, homicidal ideation,
313 depression, relapse of drug addiction, and drug overdose. These reactions may occur in
314 patients with and without previous psychiatric illness.

315 PEGASYS should be used with extreme caution in patients who report a history of
316 depression. Neuropsychiatric adverse events observed with alpha interferon treatment
317 include aggressive behavior, psychoses, hallucinations, bipolar disorders, and mania.
318 Physicians should monitor all patients for evidence of depression and other psychiatric
319 symptoms. Patients should be advised to report any sign or symptom of depression or
320 suicidal ideation to their prescribing physicians. In severe cases, therapy should be
321 stopped immediately and psychiatric intervention instituted (see **ADVERSE**
322 **REACTIONS** and **DOSAGE AND ADMINISTRATION**).

323 **Infections**

324 While fever may be associated with the flu-like syndrome reported commonly during
325 interferon therapy, other causes of high or persistent fever must be ruled out, particularly
326 in patients with neutropenia. Serious and severe infections (bacterial, viral, fungal), some
327 fatal, have been reported during treatment with alpha interferons including PEGASYS.
328 Appropriate anti-infective therapy should be started immediately and discontinuation of
329 therapy should be considered.

330 **Bone Marrow Toxicity**

331 PEGASYS suppresses bone marrow function and may result in severe cytopenias.
332 Ribavirin may potentiate the neutropenia and lymphopenia induced by alpha interferons
333 including PEGASYS. Very rarely alpha interferons may be associated with aplastic
334 anemia. It is advised that complete blood counts (CBC) be obtained pre-treatment and
335 monitored routinely during therapy (see **PRECAUTIONS: Laboratory Tests**).

336 PEGASYS and COPEGUS should be used with caution in patients with baseline
337 neutrophil counts <1500 cells/mm³, with baseline platelet counts $<90,000$ cells/mm³ or
338 baseline hemoglobin <10 g/dL. PEGASYS therapy should be discontinued, at least
339 temporarily, in patients who develop severe decreases in neutrophil and/or platelet counts
340 (see **DOSAGE AND ADMINISTRATION: Dose Modifications**).

PEGASYS® (peginterferon alfa-2a)

341 Severe neutropenia and thrombocytopenia occur with a greater incidence in HIV
342 coinfecting patients than monoinfected patients and may result in serious infections or
343 bleeding (see **ADVERSE REACTIONS**).

Cardiovascular Disorders

345 Hypertension, supraventricular arrhythmias, chest pain, and myocardial infarction have
346 been observed in patients treated with PEGASYS.

347 PEGASYS should be administered with caution to patients with pre-existing cardiac
348 disease. Because cardiac disease may be worsened by ribavirin-induced anemia, patients
349 with a history of significant or unstable cardiac disease should not use COPEGUS (see
350 **WARNINGS: Anemia and COPEGUS Package Insert**).

Cerebrovascular Disorders

352 Ischemic and hemorrhagic cerebrovascular events have been observed in patients treated
353 with interferon alfa-based therapies, including PEGASYS. Events occurred in patients
354 with few or no reported risk factors for stroke, including patients less than 45 years of
355 age. Because these are spontaneous reports, estimates of frequency cannot be made and a
356 causal relationship between interferon alfa-based therapies and these events is difficult to
357 establish.

Hepatic Failure and Hepatitis Exacerbations

359 Chronic hepatitis C (CHC) patients with cirrhosis may be at risk of hepatic
360 decompensation and death when treated with alpha interferons, including PEGASYS.
361 Cirrhotic CHC patients coinfecting with HIV receiving highly active antiretroviral therapy
362 (HAART) and interferon alfa-2a with or without ribavirin appear to be at increased risk
363 for the development of hepatic decompensation compared to patients not receiving
364 HAART. In Study 6, among 129 CHC/HIV cirrhotic patients receiving HAART, 14
365 (11%) of these patients across all treatment arms developed hepatic decompensation
366 resulting in 6 deaths. All 14 patients were on NRTIs, including stavudine, didanosine,
367 abacavir, zidovudine, and lamivudine. These small numbers of patients do not permit
368 discrimination between specific NRTIs for the associated risk. During treatment,
369 patients' clinical status and hepatic function should be closely monitored, and PEGASYS
370 treatment should be immediately discontinued if decompensation (Child-Pugh score ≥ 6)
371 is observed (see **CONTRAINDICATIONS**).

372 Exacerbations of hepatitis during hepatitis B therapy are not uncommon and are
373 characterized by transient and potentially severe increases in serum ALT. Chronic
374 hepatitis B patients experienced transient acute exacerbations (flares) of hepatitis B (ALT
375 elevation >10 -fold higher than the upper limit of normal) during PEGASYS treatment
376 (12% and 18%) and post-treatment (7% and 12%) in HBeAg negative and HBeAg
377 positive patients, respectively. Marked transaminase flares while on PEGASYS therapy
378 have been accompanied by other liver test abnormalities. Patients experiencing ALT
379 flares should receive more frequent monitoring of liver function. PEGASYS dose
380 reduction should be considered in patients experiencing transaminase flares. If ALT
381 increases are progressive despite reduction of PEGASYS dose or are accompanied by
382 increased bilirubin or evidence of hepatic decompensation, PEGASYS should be

PEGASYS® (peginterferon alfa-2a)

383 immediately discontinued (see **ADVERSE REACTIONS: Chronic Hepatitis B** and
384 **DOSAGE AND ADMINISTRATION: Dose Modifications**).

385 Hypersensitivity

386 Severe acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction,
387 and anaphylaxis) have been rarely observed during alpha interferon and ribavirin therapy.
388 If such reaction occurs, therapy with PEGASYS and COPEGUS should be discontinued
389 and appropriate medical therapy immediately instituted. Serious skin reactions including
390 vesiculobullous eruptions, reactions in the spectrum of Stevens Johnson Syndrome
391 (erythema multiforme major) with varying degrees of skin and mucosal involvement and
392 exfoliative dermatitis (erythroderma) have been rarely reported in patients receiving
393 PEGASYS with and without ribavirin. Patients developing signs or symptoms of severe
394 skin reactions must discontinue therapy (see **ADVERSE REACTIONS: Postmarketing**
395 **Experience**).

396 Endocrine Disorders

397 PEGASYS causes or aggravates hypothyroidism and hyperthyroidism. Hyperglycemia,
398 hypoglycemia, and diabetes mellitus have been observed to develop in patients treated
399 with PEGASYS. Patients with these conditions at baseline who cannot be effectively
400 treated by medication should not begin PEGASYS therapy. Patients who develop these
401 conditions during treatment and cannot be controlled with medication may require
402 discontinuation of PEGASYS therapy.

403 Autoimmune Disorders

404 Development or exacerbation of autoimmune disorders including myositis, hepatitis,
405 thrombotic thrombocytopenic purpura, idiopathic thrombocytopenic purpura, psoriasis,
406 rheumatoid arthritis, interstitial nephritis, thyroiditis, and systemic lupus erythematosus
407 have been reported in patients receiving alpha interferon. PEGASYS should be used with
408 caution in patients with autoimmune disorders.

409 Pulmonary Disorders

410 Dyspnea, pulmonary infiltrates, pneumonia, bronchiolitis obliterans, interstitial
411 pneumonitis and sarcoidosis, some resulting in respiratory failure and/or patient deaths,
412 may be induced or aggravated by PEGASYS or alpha interferon therapy. Patients who
413 develop persistent or unexplained pulmonary infiltrates or pulmonary function
414 impairment should discontinue treatment with PEGASYS.

415 Colitis

416 Ulcerative and hemorrhagic/ischemic colitis, sometimes fatal, have been observed within
417 12 weeks of starting alpha interferon treatment. Abdominal pain, bloody diarrhea, and
418 fever are the typical manifestations of colitis. PEGASYS should be discontinued
419 immediately if these symptoms develop. The colitis usually resolves within 1 to 3 weeks
420 of discontinuation of alpha interferon.

421 Pancreatitis

422 Pancreatitis, sometimes fatal, has occurred during alpha interferon and ribavirin
423 treatment. PEGASYS and COPEGUS should be suspended if symptoms or signs

PEGASYS® (peginterferon alfa-2a)

424 suggestive of pancreatitis are observed. PEGASYS and COPEGUS should be
425 discontinued in patients diagnosed with pancreatitis.

Ophthalmologic Disorders

427 Decrease or loss of vision, retinopathy including macular edema, retinal artery or vein
428 thrombosis, retinal hemorrhages and cotton wool spots, optic neuritis, papilledema and
429 serous retinal detachment are induced or aggravated by treatment with PEGASYS or
430 other alpha interferons. All patients should receive an eye examination at baseline.
431 Patients with pre-existing ophthalmologic disorders (e.g., diabetic or hypertensive
432 retinopathy) should receive periodic ophthalmologic exams during interferon alpha
433 treatment. Any patient who develops ocular symptoms should receive a prompt and
434 complete eye examination. PEGASYS treatment should be discontinued in patients who
435 develop new or worsening ophthalmologic disorders.

Pregnancy: Use with Ribavirin (also, see COPEGUS Package Insert)

437 **Ribavirin may cause birth defects and/or death of the exposed fetus. Extreme care**
438 **must be taken to avoid pregnancy in female patients and in female partners of male**
439 **patients taking PEGASYS and COPEGUS combination therapy. COPEGUS**
440 **THERAPY SHOULD NOT BE STARTED UNLESS A REPORT OF A**
441 **NEGATIVE PREGNANCY TEST HAS BEEN OBTAINED IMMEDIATELY**
442 **PRIOR TO INITIATION OF THERAPY. Women of childbearing potential and**
443 **men must use two forms of effective contraception during treatment and for at least**
444 **6 months after treatment has concluded. Routine monthly pregnancy tests must be**
445 **performed during this time (see BOXED WARNING, CONTRAINDICATIONS,**
446 **PRECAUTIONS: Information for Patients, and COPEGUS Package Insert).**

Anemia

448 The primary toxicity of ribavirin is hemolytic anemia. Hemoglobin <10 g/dL was
449 observed in approximately 13% of COPEGUS and PEGASYS treated patients in chronic
450 hepatitis C clinical trials (see **PRECAUTIONS: Laboratory Tests**). The anemia
451 associated with COPEGUS occurs within 1 to 2 weeks of initiation of therapy with
452 maximum drop in hemoglobin observed during the first eight weeks. BECAUSE THE
453 INITIAL DROP IN HEMOGLOBIN MAY BE SIGNIFICANT, IT IS ADVISED THAT
454 HEMOGLOBIN OR HEMATOCRIT BE OBTAINED PRE-TREATMENT AND AT
455 WEEK 2 AND WEEK 4 OF THERAPY OR MORE FREQUENTLY IF CLINICALLY
456 INDICATED. Patients should then be followed as clinically appropriate.

457 Fatal and nonfatal myocardial infarctions have been reported in patients with anemia
458 caused by ribavirin. Patients should be assessed for underlying cardiac disease before
459 initiation of ribavirin therapy. Patients with pre-existing cardiac disease should have
460 electrocardiograms administered before treatment, and should be appropriately monitored
461 during therapy. If there is any deterioration of cardiovascular status, therapy should be
462 suspended or discontinued (see **DOSAGE AND ADMINISTRATION: COPEGUS**
463 **Dosage Modification Guidelines**). Because cardiac disease may be worsened by drug-
464 induced anemia, patients with a history of significant or unstable cardiac disease should
465 not use COPEGUS (see **COPEGUS Package Insert**).

PEGASYS® (peginterferon alfa-2a)

Renal

It is recommended that renal function be evaluated in all patients started on COPEGUS. COPEGUS should not be administered to patients with creatinine clearance <50 mL/min (see **CLINICAL PHARMACOLOGY: Special Populations**).

PRECAUTIONS

General

The safety and efficacy of PEGASYS alone or in combination with COPEGUS have not been established in:

- Patients who have failed alpha interferon treatment with or without ribavirin
- Liver or other organ transplant recipients
- Hepatitis B patients coinfecting with HCV or HIV
- Hepatitis C patients coinfecting with HBV or coinfecting with HIV with a CD4+ cell count <100 cells/ μ L

Caution should be exercised in initiating treatment in any patient with baseline risk of severe anemia (e.g., spherocytosis, history of GI bleeding).

Renal Impairment

A 25% to 45% higher exposure to PEGASYS is seen in subjects undergoing hemodialysis. In patients with impaired renal function, signs and symptoms of interferon toxicity should be closely monitored. Doses of PEGASYS should be adjusted accordingly. PEGASYS should be used with caution in patients with creatinine clearance <50 mL/min (see **DOSAGE AND ADMINISTRATION: Dose Modifications**).

COPEGUS should not be used in patients with creatinine clearance <50 mL/min (see **COPEGUS Package Insert**).

Information for Patients

Patients receiving PEGASYS alone or in combination with COPEGUS should be directed in its appropriate use, informed of the benefits and risks associated with treatment, and referred to the PEGASYS and, if applicable, COPEGUS (ribavirin) MEDICATION GUIDES.

PEGASYS and COPEGUS combination therapy must not be used by women who are pregnant or by men whose female partners are pregnant. COPEGUS therapy should not be initiated until a report of a negative pregnancy test has been obtained immediately before starting therapy. Female patients of childbearing potential and male patients with female partners of childbearing potential must be advised of the teratogenic/embryocidal risks and must be instructed to practice effective contraception during COPEGUS therapy and for 6 months post-therapy. Patients should be advised to notify the healthcare provider immediately in the event of a pregnancy (see **CONTRAINDICATIONS** and **WARNINGS**).

PEGASYS® (peginterferon alfa-2a)

504 Women of childbearing potential and men must use two forms of effective contraception
505 during treatment and during the 6 months after treatment has been stopped; routine
506 monthly pregnancy tests must be performed during this time (see
507 **CONTRAINDICATIONS** and **COPEGUS Package Insert**).

508 To monitor maternal and fetal outcomes of pregnant women exposed to COPEGUS, the
509 Ribavirin Pregnancy Registry has been established. Patients should be encouraged to
510 register by calling 1-800-593-2214.

511 Patients should be advised that laboratory evaluations are required before starting therapy
512 and periodically thereafter (see **Laboratory Tests**). Patients should be instructed to
513 remain well hydrated, especially during the initial stages of treatment. Patients should be
514 advised to take COPEGUS with food.

515 Patients should be informed that it is not known if therapy with PEGASYS alone or in
516 combination with COPEGUS will prevent transmission of HCV or HBV infection to
517 others or prevent cirrhosis, liver failure or liver cancer that might result from HCV or
518 HBV infection. Patients who develop dizziness, confusion, somnolence, and fatigue
519 should be cautioned to avoid driving or operating machinery.

520 If home use is prescribed, a puncture-resistant container for the disposal of used needles
521 and syringes should be supplied to the patients. Patients should be thoroughly instructed
522 in the importance of proper disposal and cautioned against any reuse of any needles and
523 syringes. The full container should be disposed of according to the directions provided by
524 the physician (see **MEDICATION GUIDE**).

Laboratory Tests

526 Before beginning PEGASYS or PEGASYS and COPEGUS combination therapy,
527 standard hematological and biochemical laboratory tests are recommended for all
528 patients. Pregnancy screening for women of childbearing potential must be performed.

529 After initiation of therapy, hematological tests should be performed at 2 weeks and 4
530 weeks and biochemical tests should be performed at 4 weeks. Additional testing should
531 be performed periodically during therapy. In the clinical studies, the CBC (including
532 hemoglobin level and white blood cell and platelet counts) and chemistries (including
533 liver function tests and uric acid) were measured at 1, 2, 4, 6, and 8 weeks, and then
534 every 4 to 6 weeks or more frequently if abnormalities were found. Thyroid stimulating
535 hormone (TSH) was measured every 12 weeks. Monthly pregnancy testing should be
536 performed during combination therapy and for 6 months after discontinuing therapy.

537 The entrance criteria used for the clinical studies of PEGASYS may be considered as a
538 guideline to acceptable baseline values for initiation of treatment:

- 539 • Platelet count $\geq 90,000$ cells/mm³ (as low as 75,000 cells/mm³ in HCV patients with
540 cirrhosis or 70,000 cells/mm³ in patients with CHC and HIV)
- 541 • Absolute neutrophil count (ANC) ≥ 1500 cells/mm³
- 542 • Serum creatinine concentration $< 1.5 \times$ upper limit of normal

PEGASYS® (peginterferon alfa-2a)

- 543 • TSH and T₄ within normal limits or adequately controlled thyroid function
- 544 • CD4⁺ cell count ≥200 cells/μL or CD4⁺ cell count ≥100 cells/μL but <200 cells/μL
- 545 and HIV-1 RNA <5000 copies/mL in patients coinfecting with HIV
- 546 • Hemoglobin ≥12 g/dL for women and ≥13 g/dL for men in CHC monoinfected
- 547 patients
- 548 • Hemoglobin ≥11 g/dL for women and ≥12 g/dL for men in patients with CHC and
- 549 HIV

550 PEGASYS treatment was associated with decreases in WBC, ANC, lymphocytes, and
551 platelet counts often starting within the first 2 weeks of treatment (see **ADVERSE**
552 **REACTIONS**). Dose reduction is recommended in patients with hematologic
553 abnormalities (see **DOSAGE AND ADMINISTRATION: Dose Modifications**).

554 While fever is commonly caused by PEGASYS therapy, other causes of persistent fever
555 must be ruled out, particularly in patients with neutropenia (see **WARNINGS:**
556 **Infections**).

557 In chronic hepatitis C, transient elevations in ALT (2-fold to 5-fold above baseline) were
558 observed in some patients receiving PEGASYS, and were not associated with
559 deterioration of other liver function tests. When the increase in ALT levels is progressive
560 despite dose reduction or is accompanied by increased bilirubin, PEGASYS therapy
561 should be discontinued (see **DOSAGE AND ADMINISTRATION: Dose**
562 **Modifications**).

563 Unlike hepatitis C, during hepatitis B therapy and follow up, transient elevations in ALT
564 of 5 to 10 x ULN were observed in 25% and 27% and of >10 x ULN were observed in
565 12% and 18%, of HBeAg negative and HBeAg positive patients, respectively. These
566 ALT elevations have been accompanied by other liver test abnormalities (see
567 **WARNINGS: Hepatic Failure and Hepatitis Exacerbations** and **DOSAGE AND**
568 **ADMINISTRATION: Dose Modifications**).

569 **Drug Interactions**

570 Theophylline

571 Treatment with PEGASYS once weekly for 4 weeks in healthy subjects was associated
572 with an inhibition of P450 1A2 and a 25% increase in theophylline AUC. Theophylline
573 serum levels should be monitored and appropriate dose adjustments considered for
574 patients given both theophylline and PEGASYS (see **CLINICAL PHARMACOLOGY:**
575 **Drug Interactions**).

576 Methadone

577 In a PK study of HCV patients concomitantly receiving methadone, treatment with
578 PEGASYS once weekly for 4 weeks was associated with methadone levels that were
579 10% to 15% higher than at baseline (see **CLINICAL PHARMACOLOGY: Drug**
580 **Interactions**). The clinical significance of this finding is unknown; however, patients
581 should be monitored for the signs and symptoms of methadone toxicity.

PEGASYS® (peginterferon alfa-2a)

582 Nucleoside Analogues

583 *NRTIs*

584 In Study 6 among the CHC/HIV coinfecting cirrhotic patients receiving NRTIs cases of
585 hepatic decompensation (some fatal) were observed (see **WARNINGS: Hepatic Failure**
586 **and Hepatitis Exacerbations**).

587 Patients receiving PEGASYS/COPEGUS and NRTIs should be closely monitored for
588 treatment associated toxicities. Physicians should refer to prescribing information for the
589 respective NRTIs for guidance regarding toxicity management. In addition, dose
590 reduction or discontinuation of PEGASYS, COPEGUS or both should also be considered
591 if worsening toxicities are observed (see **WARNINGS, PRECAUTIONS, DOSAGE**
592 **AND ADMINISTRATION: Dose Modifications**).

593 *Didanosine*

594 Co-administration of COPEGUS and didanosine is not recommended. Reports of fatal
595 hepatic failure, as well as peripheral neuropathy, pancreatitis, and symptomatic
596 hyperlactatemia/lactic acidosis have been reported in clinical trials (see **CLINICAL**
597 **PHARMACOLOGY: Drug Interactions**).

598 *Zidovudine*

599 In Study 6, patients who were administered zidovudine in combination with
600 PEGASYS/COPEGUS developed severe neutropenia (ANC <500) and severe anemia
601 (hemoglobin <8 g/dL) more frequently than similar patients not receiving zidovudine
602 (neutropenia 15% vs. 9%) (anemia 5% vs. 1%). Discontinuation of zidovudine should be
603 considered as medically appropriate. Dose reduction or discontinuation of PEGASYS,
604 COPEGUS or both should also be considered if worsening clinical toxicities are
605 observed, including hepatic decompensation (e.g., Childs Pugh > 6).

606 *Lamivudine, Stavudine, and Zidovudine*

607 In vitro studies have shown ribavirin can reduce the phosphorylation of pyrimidine
608 nucleoside analogs such as lamivudine, stavudine, and zidovudine. No evidence of a
609 pharmacokinetic or pharmacodynamic interaction was seen when ribavirin was co-
610 administered with lamivudine, stavudine, and/or zidovudine in HIV/HCV coinfecting
611 patients (see **CLINICAL PHARMACOLOGY: Drug Interactions**).

612 **Carcinogenesis, Mutagenesis, Impairment of Fertility**

613 Carcinogenesis

614 PEGASYS has not been tested for its carcinogenic potential.

615 Mutagenesis

616 PEGASYS did not cause DNA damage when tested in the Ames bacterial mutagenicity
617 assay and in the in vitro chromosomal aberration assay in human lymphocytes, either in
618 the presence or absence of metabolic activation.

PEGASYS® (peginterferon alfa-2a)

Use with Ribavirin

Ribavirin is genotoxic and mutagenic. The carcinogenic potential of ribavirin has not been fully determined. In a p53 (+/-) mouse carcinogenicity study at doses up to the maximum tolerated dose of 100 mg/kg/day ribavirin was not oncogenic. However, on a body surface area basis, this dose was 0.5 times maximum recommended human 24-hour dose of ribavirin. A study in rats to assess the carcinogenic potential of ribavirin is ongoing (see **COPEGUS Package Insert**).

Impairment of Fertility

PEGASYS may impair fertility in women. Prolonged menstrual cycles and/or amenorrhea were observed in female cynomolgus monkeys given sc injections of 600 µg/kg/dose (7200 µg/m²/dose) of PEGASYS every other day for one month, at approximately 180 times the recommended weekly human dose for a 60 kg person (based on body surface area). Menstrual cycle irregularities were accompanied by both a decrease and delay in the peak 17β-estradiol and progesterone levels following administration of PEGASYS to female monkeys. A return to normal menstrual rhythm followed cessation of treatment. Every other day dosing with 100 µg/kg (1200 µg/m²) PEGASYS (equivalent to approximately 30 times the recommended human dose) had no effects on cycle duration or reproductive hormone status.

The effects of PEGASYS on male fertility have not been studied. However, no adverse effects on fertility were observed in male Rhesus monkeys treated with non-pegylated interferon alfa-2a for 5 months at doses up to 25 x 10⁶ IU/kg/day.

Use with Ribavirin

Ribavirin has shown reversible toxicity in animal studies of male fertility (see **COPEGUS Package Insert**).

Pregnancy

Pregnancy: Category C

PEGASYS has not been studied for its teratogenic effect. Non-pegylated interferon alfa-2a treatment of pregnant Rhesus monkeys at approximately 20 to 500 times the human weekly dose resulted in a statistically significant increase in abortions. No teratogenic effects were seen in the offspring delivered at term. PEGASYS should be assumed to have abortifacient potential. There are no adequate and well-controlled studies of PEGASYS in pregnant women. PEGASYS is to be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. PEGASYS is recommended for use in women of childbearing potential only when they are using effective contraception during therapy.

Pregnancy: Category X: Use With Ribavirin (see CONTRAINDICATIONS)

Significant teratogenic and/or embryocidal effects have been demonstrated in all animal species exposed to ribavirin. COPEGUS therapy is contraindicated in women who are pregnant and in the male partners of women who are pregnant (see **CONTRAINDICATIONS, WARNINGS, and COPEGUS Package Insert**).

PEGASYS® (peginterferon alfa-2a)

659 *Ribavirin Pregnancy Registry*

660 A Ribavirin Pregnancy Registry has been established to monitor maternal and fetal
661 outcomes of pregnancies of female patients and female partners of male patients exposed
662 to ribavirin during treatment and for 6 months following cessation of treatment.
663 Healthcare providers and patients are encouraged to report such cases by calling 1-800-
664 593-2214.

665 *Nursing Mothers*

666 It is not known whether peginterferon or ribavirin or its components are excreted in
667 human milk. The effect of orally ingested peginterferon or ribavirin from breast milk on
668 the nursing infant has not been evaluated. Because of the potential for adverse reactions
669 from the drugs in nursing infants, a decision must be made whether to discontinue
670 nursing or discontinue PEGASYS and COPEGUS treatment.

671 *Pediatric Use*

672 The safety and effectiveness of PEGASYS, alone or in combination with COPEGUS in
673 patients below the age of 18 years have not been established.

674 PEGASYS contains benzyl alcohol. Benzyl alcohol has been reported to be associated
675 with an increased incidence of neurological and other complications in neonates and
676 infants, which are sometimes fatal (see **CONTRAINDICATIONS**).

677 *Geriatric Use*

678 Younger patients have higher virologic response rates than older patients. Clinical studies
679 of PEGASYS alone or in combination with COPEGUS did not include sufficient
680 numbers of subjects aged 65 or over to determine whether they respond differently from
681 younger subjects. Adverse reactions related to alpha interferons, such as CNS, cardiac,
682 and systemic (e.g., flu-like) effects may be more severe in the elderly and caution should
683 be exercised in the use of PEGASYS in this population. PEGASYS and COPEGUS are
684 excreted by the kidney, and the risk of toxic reactions to this therapy may be greater in
685 patients with impaired renal function. Because elderly patients are more likely to have
686 decreased renal function, care should be taken in dose selection and it may be useful to
687 monitor renal function. PEGASYS should be used with caution in patients with creatinine
688 clearance <50 mL/min and COPEGUS should not be administered to patients with
689 creatinine clearance <50 mL/min.

690 *ADVERSE REACTIONS*

691 PEGASYS alone or in combination with COPEGUS causes a broad variety of serious
692 adverse reactions (see **BOXED WARNING** and **WARNINGS**). The most common life-
693 threatening or fatal events induced or aggravated by PEGASYS and COPEGUS were
694 depression, suicide, relapse of drug abuse/overdose, and bacterial infections, each
695 occurring at a frequency of <1%. Hepatic decompensation occurred in 2% (10/574) of
696 CHC/HIV patients (see **WARNINGS: Hepatic Failure and Hepatitis Exacerbations**).

697 In all hepatitis C studies, one or more serious adverse reactions occurred in 10% of CHC
698 monoinfected patients and in 19% of CHC/HIV patients receiving PEGASYS alone or in
699 combination with COPEGUS. The most common serious adverse event (3% in CHC and

PEGASYS® (peginterferon alfa-2a)

5% in CHC/HIV) was bacterial infection (e.g., sepsis, osteomyelitis, endocarditis, pyelonephritis, pneumonia). Other SAEs occurred at a frequency of <1% and included: suicide, suicidal ideation, psychosis, aggression, anxiety, drug abuse and drug overdose, angina, hepatic dysfunction, fatty liver, cholangitis, arrhythmia, diabetes mellitus, autoimmune phenomena (e.g., hyperthyroidism, hypothyroidism, sarcoidosis, systemic lupus erythematosus, rheumatoid arthritis), peripheral neuropathy, aplastic anemia, peptic ulcer, gastrointestinal bleeding, pancreatitis, colitis, corneal ulcer, pulmonary embolism, coma, myositis, cerebral hemorrhage, thrombotic thrombocytopenic purpura, psychotic disorder, and hallucination.

Nearly all patients in clinical trials experienced one or more adverse events. For hepatitis C patients, the most commonly reported adverse reactions were psychiatric reactions, including depression, insomnia, irritability, anxiety, and flu-like symptoms such as fatigue, pyrexia, myalgia, headache, and rigors. Other common reactions were anorexia, nausea and vomiting, diarrhea, arthralgias, injection site reactions, alopecia, and pruritus.

Overall 11% of CHC monoinfected patients receiving 48 weeks of therapy with PEGASYS either alone or in combination with COPEGUS discontinued therapy; 16% of CHC/HIV coinfecting patients discontinued therapy. The most common reasons for discontinuation of therapy were psychiatric, flu-like syndrome (e.g., lethargy, fatigue, headache), dermatologic, and gastrointestinal disorders and laboratory abnormalities (thrombocytopenia, neutropenia, and anemia).

Overall 39% of patients with CHC or CHC/HIV required modification of PEGASYS and/or COPEGUS therapy. The most common reason for dose modification of PEGASYS in CHC and CHC/HIV patients was for laboratory abnormalities, neutropenia (20% and 27%, respectively) and thrombocytopenia (4% and 6%, respectively). The most common reason for dose modification of COPEGUS in CHC and CHC/HIV patients was anemia (22% and 16%, respectively).

PEGASYS dose was reduced in 12% of patients receiving 1000 mg to 1200 mg COPEGUS for 48 weeks and in 7% of patients receiving 800 mg COPEGUS for 24 weeks. COPEGUS dose was reduced in 21% of patients receiving 1000 mg to 1200 mg COPEGUS for 48 weeks and in 12% of patients receiving 800 mg COPEGUS for 24 weeks.

Chronic hepatitis C monoinfected patients treated for 24 weeks with PEGASYS and 800 mg COPEGUS were observed to have lower incidence of serious adverse events (3% vs. 10%), Hgb <10 g/dL (3% vs. 15%), dose modification of PEGASYS (30% vs. 36%) and COPEGUS (19% vs. 38%) and of withdrawal from treatment (5% vs. 15%) compared to patients treated for 48 weeks with PEGASYS and 1000 mg or 1200 mg COPEGUS. On the other hand the overall incidence of adverse events appeared to be similar in the two treatment groups.

Because clinical trials are conducted under widely varying and controlled conditions, adverse reaction rates observed in clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug. Also, the adverse event rates listed here may not predict the rates observed in a broader patient population in clinical practice.

PEGASYS® (peginterferon alfa-2a)

743 **Table 6** **Adverse Reactions Occurring in ≥5% of Patients in Chronic**
744 **Hepatitis C Clinical Trials (Pooled Studies 1, 2, 3, and**
745 **Study 4)**

| | CHC Monotherapy (Pooled Studies 1-3) | | CHC Combination Therapy Study 4 | |
|--|---|-------------|--|---|
| Body System | PEGASYS 180 µg 48 week† | ROFERON-A*† | PEGASYS 180 µg + 1000 mg or 1200 mg COPEGUS 48 week** | Intron® A + 1000 mg or 1200 mg REBETOL® 48 week** |
| | N=559 | N=554 | N=451 | N=443 |
| | % | % | % | % |
| Application Site Disorders | | | | |
| Injection site reaction | 22 | 18 | 23 | 16 |
| Endocrine Disorders | | | | |
| Hypothyroidism | 3 | 2 | 4 | 5 |
| Flu-like Symptoms and Signs | | | | |
| Fatigue/Asthenia | 56 | 57 | 65 | 68 |
| Pyrexia | 37 | 41 | 41 | 55 |
| Rigors | 35 | 44 | 25 | 37 |
| Pain | 11 | 12 | 10 | 9 |
| Gastrointestinal | | | | |
| Nausea/Vomiting | 24 | 33 | 25 | 29 |
| Diarrhea | 16 | 16 | 11 | 10 |
| Abdominal pain | 15 | 15 | 8 | 9 |
| Dry mouth | 6 | 3 | 4 | 7 |
| Dyspepsia | <1 | 1 | 6 | 5 |
| Hematologic‡ | | | | |
| Lymphopenia | 3 | 5 | 14 | 12 |
| Anemia | 2 | 1 | 11 | 11 |
| Neutropenia | 21 | 8 | 27 | 8 |
| Thrombocytopenia | 5 | 2 | 5 | <1 |
| Metabolic and Nutritional | | | | |
| Anorexia | 17 | 17 | 24 | 26 |
| Weight decrease | 4 | 3 | 10 | 10 |

PEGASYS® (peginterferon alfa-2a)

| | CHC Monotherapy (Pooled Studies 1-3) | | CHC Combination Therapy Study 4 | |
|--|---|--------------------|--|--|
| Body System | PEGASYS 180 µg 48 week† | ROFERON-A*† | PEGASYS 180 µg + 1000 mg or 1200 mg COPEGUS 48 week** | Intron® A + 1000 mg or 1200 mg REBETOL® 48 week** |
| | N=559 | N=554 | N=451 | N=443 |
| | % | % | % | % |
| Musculoskeletal, Connective Tissue and Bone | | | | |
| Myalgia | 37 | 38 | 40 | 49 |
| Arthralgia | 28 | 29 | 22 | 23 |
| Back pain | 9 | 10 | 5 | 5 |
| Neurological | | | | |
| Headache | 54 | 58 | 43 | 49 |
| Dizziness (excluding vertigo) | 16 | 12 | 14 | 14 |
| Memory impairment | 5 | 4 | 6 | 5 |
| Resistance Mechanism Disorders | | | | |
| Overall | 10 | 6 | 12 | 10 |
| Psychiatric | | | | |
| Irritability/Anxiety/ Nervousness | 19 | 22 | 33 | 38 |
| Insomnia | 19 | 23 | 30 | 37 |
| Depression | 18 | 19 | 20 | 28 |
| Concentration impairment | 8 | 10 | 10 | 13 |
| Mood alteration | 3 | 2 | 5 | 6 |
| Respiratory, Thoracic and Mediastinal | | | | |
| Dyspnea | 4 | 2 | 13 | 14 |
| Cough | 4 | 3 | 10 | 7 |
| Dyspnea exertional | <1 | <1 | 4 | 7 |

PEGASYS® (peginterferon alfa-2a)

| | CHC Monotherapy (Pooled Studies 1-3) | | CHC Combination Therapy Study 4 | |
|-------------------------------------|--------------------------------------|--------------|--|---|
| Body System | PEGASYS 180 µg 48 week† | ROFERON-A**† | PEGASYS 180 µg + 1000 mg or 1200 mg COPEGUS 48 week** | Intron® A + 1000 mg or 1200 mg REBETOL® 48 week** |
| | N=559 | N=554 | N=451 | N=443 |
| | % | % | % | % |
| Skin and Subcutaneous Tissue | | | | |
| Alopecia | 23 | 30 | 28 | 33 |
| Pruritus | 12 | 8 | 19 | 18 |
| Dermatitis | 8 | 3 | 16 | 13 |
| Dry skin | 4 | 3 | 10 | 13 |
| Rash | 5 | 4 | 8 | 5 |
| Sweating increased | 6 | 7 | 6 | 5 |
| Eczema | 1 | 1 | 5 | 4 |
| Visual Disorders | | | | |
| Vision blurred | 4 | 2 | 5 | 2 |

† Pooled studies 1, 2, and 3

* Either 3 MIU or 6/3 MIU of ROFERON-A

**Study 4

‡ Severe hematologic abnormalities (lymphocyte <0.5 x 10⁹/L; hemoglobin <10 g/dL; neutrophil <0.75 x 10⁹/L; platelet <50 x 10⁹/L).

CHC With HIV Coinfection

The adverse event profile of coinfecting patients treated with PEGASYS and COPEGUS in Study 6 was generally similar to that shown for mono-infected patients in Study 4 (Table 6). Events occurring more frequently in coinfecting patients were neutropenia (40%), anemia (14%), thrombocytopenia (8%), weight decrease (16%), and mood alteration (9%).

Chronic Hepatitis B

In clinical trials of 48 week treatment duration, the adverse event profile of PEGASYS in chronic hepatitis B was similar to that seen in chronic hepatitis C PEGASYS monotherapy use, except for exacerbations of hepatitis (see **WARNINGS: Hepatic Failure and Hepatitis Exacerbations**). Six percent of PEGASYS treated patients in the hepatitis B studies experienced one or more serious adverse events.

PEGASYS® (peginterferon alfa-2a)

764 The most common or important serious adverse events in the hepatitis B studies were
765 infections (sepsis, appendicitis, tuberculosis, influenza), hepatitis B flares, anaphylactic
766 shock, thrombotic thrombocytopenic purpura.

767 The most commonly observed adverse reactions were pyrexia (54% vs. 4%), headache
768 (27% vs. 9%), fatigue (24% vs. 10%), myalgia (26% vs. 4%), alopecia (18% vs. 2%), and
769 anorexia (16% vs. 3%) in the PEGASYS and lamivudine groups respectively.

770 Overall 5% of hepatitis B patients discontinued PEGASYS therapy and 40% of patients
771 required modification of PEGASYS dose. The most common reason for dose
772 modification in patients receiving PEGASYS therapy was for laboratory abnormalities
773 including neutropenia (20%), thrombocytopenia (13%), and ALT disorders (11%).

774 Laboratory Test Values

775 The laboratory test values observed in the hepatitis B trials (except where noted below)
776 were similar to those seen in the PEGASYS monotherapy hepatitis C trials.

777 Neutrophils

778 In the hepatitis C studies, decreases in neutrophil count below normal were observed in
779 95% of all patients treated with PEGASYS either alone or in combination with
780 COPEGUS. Severe potentially life-threatening neutropenia ($ANC < 0.5 \times 10^9/L$) occurred
781 in 5% of CHC patients and 12% of CHC/HIV patients receiving PEGASYS either alone
782 or in combination with COPEGUS. Modification of PEGASYS dose for neutropenia
783 occurred in 17% of patients receiving PEGASYS monotherapy and 22% of patients
784 receiving PEGASYS/COPEGUS combination therapy. In the CHC/HIV patients 27%
785 required modification of interferon dosage for neutropenia. Two percent of patients with
786 CHC and 10% of patients with CHC/HIV required permanent reductions of PEGASYS
787 dosage and <1% required permanent discontinuation. Median neutrophil counts return to
788 pre-treatment levels 4 weeks after cessation of therapy (see **DOSAGE AND**
789 **ADMINISTRATION: Dose Modifications**).

790 Lymphocytes

791 Decreases in lymphocyte count are induced by interferon alpha therapy. PEGASYS plus
792 COPEGUS combination therapy induced decreases in median total lymphocyte counts
793 (56% in CHC and 40% in CHC/HIV, with median decrease of 1170 cells/mm^3 in CHC
794 and 800 cells/mm^3 in CHC/HIV). In the hepatitis C studies, lymphopenia was observed
795 during both monotherapy (81%) and combination therapy with PEGASYS and
796 COPEGUS (91%). Severe lymphopenia ($< 0.5 \times 10^9/L$) occurred in approximately 5% of
797 all monotherapy patients and 14% of all combination PEGASYS and COPEGUS therapy
798 recipients. Dose adjustments were not required by protocol. The clinical significance of
799 the lymphopenia is not known.

800 In CHC with HIV coinfection, CD4 counts decreased by 29% from baseline (median
801 decrease of 137 cells/mm^3) and CD8 counts decreased by 44% from baseline (median
802 decrease of 389 cells/mm^3) in the PEGASYS plus COPEGUS combination therapy arm.
803 Median lymphocyte CD4 and CD8 counts return to pre-treatment levels after 4 to 12
804 weeks of the cessation of therapy. CD4% did not decrease during treatment.

PEGASYS® (peginterferon alfa-2a)

805 Platelets

806 In the hepatitis C studies, platelet counts decreased in 52% of CHC patients and 51% of
807 CHC/HIV patients treated with PEGASYS alone (respectively median decrease of 41%
808 and 35% from baseline), and in 33% of CHC patients and 47% of CHC/HIV patients
809 receiving combination therapy with COPEGUS (median decrease of 30% from baseline).
810 Moderate to severe thrombocytopenia ($<50,000/\text{mm}^3$) was observed in 4% of CHC and
811 8% of CHC/HIV patients. Median platelet counts return to pre-treatment levels 4 weeks
812 after the cessation of therapy.

813 Hemoglobin

814 In the hepatitis C studies, the hemoglobin concentration decreased below 12 g/dL in 17%
815 (median Hgb reduction of 2.2 g/dL) of monotherapy and 52% (median Hgb reduction of
816 3.7 g/dL) of combination therapy patients. Severe anemia (Hgb <10 g/dL) was
817 encountered in 13% of all patients receiving combination therapy and in 2% of CHC
818 patients and 8% of CHC/HIV patients receiving PEGASYS monotherapy. Dose
819 modification for anemia in COPEGUS recipients treated for 48 weeks occurred in 22% of
820 CHC patients and 16% of CHC/HIV patients (see **DOSAGE AND**
821 **ADMINISTRATION: Dose Modifications**).

822 Triglycerides

823 Triglyceride levels are elevated in patients receiving alfa interferon therapy and were
824 elevated in the majority of patients participating in clinical studies receiving either
825 PEGASYS alone or in combination with COPEGUS. Random levels ≥ 400 mg/dL were
826 observed in about 20% of CHC patients. Severe elevations of triglycerides (>1000
827 mg/dL) occurred in 2% of CHC monoinfected patients.

828 In HCV/HIV coinfecting patients, fasting levels ≥ 400 mg/dL were observed in up to 36%
829 of patients receiving either PEGASYS alone or in combination with COPEGUS. Severe
830 elevations of triglycerides (>1000 mg/dL) occurred in 7% of coinfecting patients.

831 ALT Elevations

832 *Chronic Hepatitis C*

833 One percent of patients in the hepatitis C trials experienced marked elevations (5- to 10-
834 fold above the upper limit of normal) in ALT levels during treatment and follow-up.
835 These transaminase elevations were on occasion associated with hyperbilirubinemia and
836 were managed by dose reduction or discontinuation of study treatment. Liver function
837 test abnormalities were generally transient. One case was attributed to autoimmune
838 hepatitis, which persisted beyond study medication discontinuation (see **DOSAGE AND**
839 **ADMINISTRATION: Dose Modifications**).

840 *Chronic Hepatitis B*

841 Transient ALT elevations are common during hepatitis B therapy with PEGASYS.
842 Twenty-five percent and 27% of patients experienced elevations of 5 to 10 x ULN and
843 12% and 18% had elevations of >10 x ULN during treatment of HBeAg negative and
844 HBeAg positive disease, respectively. Flares have been accompanied by elevations of
845 total bilirubin and alkaline phosphatase and less commonly with prolongation of PT and
846 reduced albumin levels. Eleven percent of patients had dose modifications due to ALT

PEGASYS® (peginterferon alfa-2a)

847 flares and <1% of patients were withdrawn from treatment (see **WARNINGS: Hepatic**
848 **Failure and Hepatitis Exacerbations** and **DOSAGE AND ADMINISTRATION:**
849 **Dose Modifications**).

850 ALT flares of 5 to 10 x ULN occurred in 13% and 16% of patients, while ALT flares of
851 >10 x ULN occurred in 7% and 12% of patients in HBeAg negative and HBeAg positive
852 disease, respectively, after discontinuation of PEGASYS therapy.

853 Thyroid Function

854 PEGASYS alone or in combination with COPEGUS was associated with the
855 development of abnormalities in thyroid laboratory values, some with associated clinical
856 manifestations. In the hepatitis C studies, hypothyroidism or hyperthyroidism requiring
857 treatment, dose modification or discontinuation occurred in 4% and 1% of PEGASYS
858 treated patients and 4% and 2% of PEGASYS and COPEGUS treated patients,
859 respectively. Approximately half of the patients, who developed thyroid abnormalities
860 during PEGASYS treatment, still had abnormalities during the follow-up period (see
861 **PRECAUTIONS: Laboratory Tests**).

862 Immunogenicity

863 *Chronic Hepatitis C*

864 Nine percent (71/834) of patients treated with PEGASYS with or without COPEGUS
865 developed binding antibodies to interferon alfa-2a, as assessed by an ELISA assay. Three
866 percent of patients (25/835) receiving PEGASYS with or without COPEGUS, developed
867 low-titer neutralizing antibodies (using an assay with a sensitivity of 100 INU/mL).

868 *Chronic Hepatitis B*

869 Twenty-nine percent (42/143) of hepatitis B patients treated with PEGASYS for 24
870 weeks developed binding antibodies to interferon alfa-2a, as assessed by an ELISA assay.
871 Thirteen percent of patients (19/143) receiving PEGASYS developed low-titer
872 neutralizing antibodies (using an assay with a sensitivity of 100 INU/mL).

873 The clinical and pathological significance of the appearance of serum neutralizing
874 antibodies is unknown. No apparent correlation of antibody development to clinical
875 response or adverse events was observed. The percentage of patients whose test results
876 were considered positive for antibodies is highly dependent on the sensitivity and
877 specificity of the assays.

878 Additionally, the observed incidence of antibody positivity in these assays may be
879 influenced by several factors including sample timing and handling, concomitant
880 medications, and underlying disease. For these reasons, comparison of the incidence of
881 antibodies to PEGASYS with the incidence of antibodies to other products may be
882 misleading.

883 Postmarketing Experience

884 The following adverse reactions have been identified and reported during post-approval
885 use of PEGASYS therapy: dehydration, hearing impairment, hearing loss, and serious
886 skin reactions (see **WARNINGS: Hypersensitivity**). Because these reactions are
887 reported voluntarily from a population of uncertain size, it is not always possible to

PEGASYS® (peginterferon alfa-2a)

888 reliably estimate their frequency or establish a causal relationship to drug exposure.
889 Decisions to include these reactions in labeling are typically based on one or more of the
890 following factors: (1) seriousness of the reaction, (2) frequency of reporting or (3)
891 strength of causal connection to PEGASYS.

892 OVERDOSAGE

893 There is limited experience with overdosage. The maximum dose received by any patient
894 was 7 times the intended dose of PEGASYS (180 µg/day for 7 days). There were no
895 serious reactions attributed to overdosages. Weekly doses of up to 630 µg have been
896 administered to patients with cancer. Dose-limiting toxicities were fatigue, elevated liver
897 enzymes, neutropenia, and thrombocytopenia. There is no specific antidote for
898 PEGASYS. Hemodialysis and peritoneal dialysis are not effective.

899 DOSAGE AND ADMINISTRATION

900 There are no safety and efficacy data on treatment of chronic hepatitis C or hepatitis B for
901 longer than 48 weeks. For patients with hepatitis C, consideration should be given to
902 discontinuing therapy after 12 to 24 weeks of therapy if the patient has failed to
903 demonstrate an early virologic response defined as undetectable HCV RNA or at least a
904 2log₁₀ reduction from baseline in HCV RNA titer by 12 weeks of therapy (see
905 **CLINICAL STUDIES**).

906 A patient should self-inject PEGASYS only if the physician determines that it is
907 appropriate and the patient agrees to medical follow-up as necessary and training in
908 proper injection technique has been provided to him/her (see illustrated PEGASYS
909 **MEDICATION GUIDE** for directions on injection site preparation and injection
910 instructions).

911 PEGASYS should be inspected visually for particulate matter and discoloration before
912 administration, and not used if particulate matter is visible or product is discolored. Vials
913 and prefilled syringes with particulate matter or discoloration should be returned to the
914 pharmacist.

915 Chronic Hepatitis C

916 PEGASYS Monotherapy

917 The recommended dose of PEGASYS monotherapy for chronic hepatitis C is 180 µg (1.0
918 mL vial or 0.5 mL prefilled syringe) once weekly for 48 weeks by subcutaneous
919 administration in the abdomen or thigh.

920 PEGASYS and COPEGUS Combination Therapy

921 The recommended dose of PEGASYS when used in combination with ribavirin for
922 chronic hepatitis C is 180 µg (1.0 mL vial or 0.5 mL prefilled syringe) once weekly. The
923 recommended dose of COPEGUS and duration for PEGASYS/COPEGUS therapy is
924 based on viral genotype (see **Table 7**).

925 The daily dose of COPEGUS is 800 mg to 1200 mg administered orally in two divided
926 doses. The dose should be individualized to the patient depending on baseline disease
927 characteristics (e.g., genotype), response to therapy, and tolerability of the regimen.

PEGASYS® (peginterferon alfa-2a)

928 Since COPEGUS absorption increases when administered with a meal, patients are
929 advised to take COPEGUS with food.

930 **Table 7 PEGASYS and COPEGUS Dosing Recommendations**

| Genotype | PEGASYS Dose | COPEGUS Dose | Duration |
|----------------|--------------|------------------|----------|
| Genotypes 1, 4 | 180 µg | <75 kg = 1000 mg | 48 weeks |
| | | ≥75 kg = 1200 mg | 48 weeks |
| Genotypes 2, 3 | 180 µg | 800 mg | 24 weeks |

931 Genotypes 2 and 3 showed no increased response to treatment beyond 24 weeks (see Table 3).

932 Data on genotypes 5 and 6 are insufficient for dosing recommendations.

933

934 **CHC with HIV Coinfection**

935 **PEGASYS Monotherapy**

936 The recommended dose of PEGASYS monotherapy for chronic hepatitis C in patients
937 coinfecting with HIV is 180 µg (1.0 mL vial or 0.5 mL prefilled syringe) once weekly for
938 48 weeks by subcutaneous administration in the abdomen or thigh.

939 **PEGASYS/COPEGUS Combination Therapy**

940 The recommended dose when used in combination with ribavirin is PEGASYS 180 µg sc
941 once weekly and COPEGUS 800 mg po daily given in two divided doses for a total of 48
942 weeks, regardless of genotype.

943 Since COPEGUS absorption increases when administered with a meal, patients are
944 advised to take COPEGUS with food.

945 **Chronic Hepatitis B**

946 **PEGASYS Monotherapy**

947 The recommended dose of PEGASYS monotherapy for hepatitis B is 180 µg (1.0 mL
948 vial or 0.5 mL prefilled syringe) once weekly for 48 weeks by subcutaneous
949 administration in the abdomen or thigh.

950 **Dose Modifications**

951 **If severe adverse reactions or laboratory abnormalities develop during combination**
952 **COPEGUS/PEGASYS therapy, the dose should be modified or discontinued, if**
953 **appropriate, until the adverse reactions abate. If intolerance persists after dose**
954 **adjustment, COPEGUS/PEGASYS therapy should be discontinued.**

955 **PEGASYS**

956 **General**

957 When dose modification is required for moderate to severe adverse reactions (clinical
958 and/or laboratory), initial dose reduction to 135 µg (which is 0.75 mL for the vials or
959 adjustment to the corresponding graduation mark for the syringes) is generally adequate.
960 However, in some cases, dose reduction to 90 µg (which is 0.5 mL for the vials or

PEGASYS® (peginterferon alfa-2a)

adjustment to the corresponding graduation mark for the syringes) may be needed. Following improvement of the adverse reaction, re-escalation of the dose may be considered (see **WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS**).

Hematological

Table 8 PEGASYS Hematological Dose Modification Guidelines

| Laboratory Values | Reduce PEGASYS Dose to: | Discontinue PEGASYS if: |
|---|---|--|
| ANC $\geq 750/\text{mm}^3$ ANC $< 750/\text{mm}^3$ | Maintain 180 μg Reduce to 135 μg | ANC $< 500/\text{mm}^3$, treatment should be suspended until ANC values return to more than $1000/\text{mm}^3$ Reinstitute at 90 μg and monitor ANC |
| Platelet $\geq 50,000/\text{mm}^3$ Platelet $< 50,000/\text{mm}^3$ | Maintain 180 μg Reduce to 90 μg | Platelet count $< 25,000/\text{mm}^3$ |

Psychiatric: Depression

Table 9 Guidelines for Modification or Discontinuation of PEGASYS and for Scheduling Visits for Patients with Depression

| Depression Severity | Initial Management (4-8 weeks) | | Depression | | |
|---------------------|---|---|--|--|-------------------------------------|
| | Dose modification | Visit schedule | Remains stable | Improves | Worsens |
| Mild | No change | Evaluate once weekly by visit and/or phone | Continue weekly visit schedule | Resume normal visit schedule | (See moderate or severe depression) |
| Moderate | Decrease PEGASYS dose to 135 μg (in some cases dose reduction to 90 μg may be needed) | Evaluate once weekly (office visit at least every other week) | Consider psychiatric consultation. Continue reduced dosing | If symptoms improve and are stable for 4 weeks, may resume normal visit schedule. Continue reduced dosing or return to normal dose | (See severe depression) |
| Severe | Discontinue PEGASYS permanently | Obtain immediate psychiatric | Psychiatric therapy necessary | | |

PEGASYS® (peginterferon alfa-2a)

| | | | |
|--|--|--------------|--|
| | | consultation | |
|--|--|--------------|--|

969 Renal Function

970 In patients with end-stage renal disease requiring hemodialysis, dose reduction to 135 µg
971 PEGASYS is recommended. Signs and symptoms of interferon toxicity should be closely
972 monitored.

973 Liver Function

974 If ALT increases are progressive despite dose reduction or accompanied by increased
975 bilirubin or evidence of hepatic decompensation, therapy should be immediately
976 discontinued.

977 In chronic hepatitis C patients with progressive ALT increases above baseline values, the
978 dose of PEGASYS should be reduced to 135 µg and more frequent monitoring of liver
979 function should be performed. After PEGASYS dose reduction or withholding, therapy
980 can be resumed after ALT flares subside.

981 In chronic hepatitis B patients with elevations in ALT (>5 x ULN), more frequent
982 monitoring of liver function should be performed and consideration should be given to
983 either reducing the dose of PEGASYS to 135 µg or temporarily discontinuing treatment.
984 After PEGASYS dose reduction or withholding, therapy can be resumed after ALT flares
985 subside.

986 In patients with persistent, severe (ALT >10 times above the upper limit of normal)
987 hepatitis B flares, consideration should be given to discontinuation of treatment.

988 COPEGUS

989 **Table 10 COPEGUS Dosage Modification Guidelines**

| Laboratory Values | Reduce Only COPEGUS Dose to 600 mg/day* if: | Discontinue COPEGUS if: |
|---|---|---|
| Hemoglobin in patients with no cardiac disease | <10 g/dL | <8.5 g/dL |
| Hemoglobin in patients with history of stable cardiac disease | ≥2 g/dL decrease in hemoglobin during any 4 week period treatment | <12 g/dL despite 4 weeks at reduced dose |

990 * One 200 mg tablet in the morning and two 200 mg tablets in the evening.
991

992 Once COPEGUS has been withheld due to a laboratory abnormality or clinical
993 manifestation, an attempt may be made to restart COPEGUS at 600 mg daily and further
994 increase the dose to 800 mg daily depending upon the physician's judgment. However, it
995 is not recommended that COPEGUS be increased to the original dose (1000 mg or
996 1200 mg).

PEGASYS® (peginterferon alfa-2a)

997 Renal Impairment

998 COPEGUS should not be used in patients with creatinine clearance <50 mL/min (see
999 **CLINICAL PHARMACOLOGY, WARNINGS and COPEGUS Package Insert**).

1000 HOW SUPPLIED

1001 Single Dose Vial

1002 Each PEGASYS (peginterferon alfa-2a) 180 µg single use, clear glass vial provides
1003 1.0 mL containing 180 µg peginterferon alfa-2a for sc injection. Each package contains 1
1004 vial (NDC 0004-0350-09).

1005 Prefilled Syringes Monthly Convenience Pack

1006 Four prefilled syringes of PEGASYS (peginterferon alfa-2a), 180 µg single use,
1007 graduated, clear glass prefilled syringes, in a box with 4 needles and 4 alcohol swabs
1008 (NDC 0004-0352-39). Each syringe is a 0.5 mL (½ cc) volume syringe supplied with a
1009 27-gauge, ½-inch needle with needle-stick protection device.

1010 Storage

1011 Store in the refrigerator at 2°C to 8°C (36°F to 46°F). Do not freeze or shake. Protect
1012 from light. Vials and prefilled syringes are for single use only. Discard any unused
1013 portion.

1014 REBETRON®, REBETROL®, and INTRON® are registered trademarks of Schering
1015 Corporation.

1016 PI Revised: April 2009

1017 MEDICATION GUIDE

1018 PEGASYS®

1019 (peginterferon alfa-2a)

1020 Before you start taking PEGASYS (PEG-ah-sis), alone or in combination with
1021 COPEGUS® (Co-PEG-UHS), please read this Medication Guide carefully. Read this
1022 Medication Guide each time you refill your prescription in case new information has
1023 been added and make sure the pharmacist has given you the medicine your healthcare
1024 provider prescribed for you. Reading the information in this Medication Guide does not
1025 take the place of talking with your healthcare provider.

1026 If you are taking PEGASYS in combination with COPEGUS, you should also read the
1027 Medication Guide for COPEGUS (ribavirin, USP) Tablets.

1028 What is the most important information I should know about PEGASYS 1029 therapy?

1030 PEGASYS, taken alone or in combination with COPEGUS, is a treatment for some
1031 people who are infected with hepatitis C virus. PEGASYS taken alone is a treatment for
1032 some people who are infected with the hepatitis B virus. However, PEGASYS and
1033 COPEGUS can have serious side effects that may cause death in rare cases. Before
1034 starting PEGASYS therapy, you should talk with your healthcare provider about the

PEGASYS® (peginterferon alfa-2a)

1035 possible benefits and the possible side effects of treatment, to decide if either of these
1036 treatments is right for you. If you begin treatment you will need to see your healthcare
1037 provider regularly for examinations and blood tests to make sure your treatment is
1038 working and to check for side effects.

1039 The most serious possible side effects of PEGASYS taken alone or in combination with
1040 COPEGUS include:

1041 Problems with Pregnancy:

1042 **Taking PEGASYS in combination with COPEGUS tablets can cause death, serious**
1043 **birth defects or other harm to your unborn child. Therefore, if you are pregnant or**
1044 **your partner is pregnant or plans to become pregnant, do not take**
1045 **PEGASYS/COPEGUS combination therapy. Female patients and female partners**
1046 **of male patients being treated with PEGASYS/COPEGUS combination therapy**
1047 **must not become pregnant during treatment and for 6 months after treatment has**
1048 **stopped. During this time, you must have pregnancy tests that show you are not**
1049 **pregnant. You must also use two effective forms of birth control during therapy and**
1050 **for 6 months after stopping therapy. Male patients should use a condom with**
1051 **spermicide as one of the two forms. You must use birth control even if you believe that**
1052 **you are not fertile or that your fertility is low. You should talk to your healthcare provider**
1053 **about birth control for you and your partner.**

1054 **If you are pregnant, you or your male partner must not take PEGASYS/COPEGUS**
1055 **combination therapy. If you or your partner are being treated and you become**
1056 **pregnant either during treatment or within 6 months of stopping treatment, call**
1057 **your healthcare provider right away.**

1058 If you or a female sexual partner becomes pregnant, you should tell your healthcare
1059 provider. There is a Ribavirin Pregnancy Registry that collects information about
1060 pregnancy outcomes of female patients and female partners of male patients exposed to
1061 ribavirin. You or your healthcare provider are encouraged to contact the Registry at 1-
1062 800-593-2214.

1063 Mental health problems and suicide:

1064 PEGASYS and PEGASYS/COPEGUS combination therapy may cause some patients to
1065 develop mood or behavioral problems. Signs of these problems include irritability
1066 (getting easily upset), depression (feeling low, feeling bad about yourself or feeling
1067 hopeless), and anxiety. Some patients may have aggressive behavior. Former drug addicts
1068 may fall back into drug addiction or overdose. Some patients think about hurting or
1069 killing themselves or other people and some have killed (suicide) or hurt themselves or
1070 hurt other people. You must tell your healthcare provider if you are being treated for a
1071 mental illness or have a history of mental illness, including depression and suicidal
1072 behavior or if you are or have ever been addicted to drugs or alcohol. Call your
1073 healthcare provider immediately if you develop any of these problems while on
1074 PEGASYS treatment.

PEGASYS® (peginterferon alfa-2a)

1075 Heart problems:

1076 Some patients taking PEGASYS or PEGASYS/COPEGUS therapies may develop
1077 problems with their heart, including low blood pressure, fast heart rate, and very rarely,
1078 heart attacks. Tell your healthcare provider if you have had any heart problems in the
1079 past.

1080 Blood problems:

1081 Many patients taking PEGASYS have had a drop in the number of their white blood cells
1082 and their platelets. If the numbers of these blood cells are too low, you could be at risk for
1083 serious infections or bleeding.

1084 COPEGUS causes a decrease in the number of your red blood cells (anemia). This can be
1085 dangerous, especially for patients who already have heart or circulatory (cardiovascular)
1086 problems. If you have or have ever had any cardiovascular problems, talk with your
1087 healthcare provider before taking the combination of PEGASYS and COPEGUS.

1088 Liver problems:

1089 Infrequently, some patients with hepatitis C and liver scarring can develop sudden severe
1090 worsening (failure) of their liver disease while taking PEGASYS. Patients infected with
1091 both the hepatitis C virus and HIV can have an increased chance of having liver failure
1092 during PEGASYS treatment.

1093 Some patients taking PEGASYS for hepatitis B have had a rise in a blood test that
1094 measures liver inflammation. If you have a rise in this blood test, your liver may need to
1095 be watched more closely with additional blood tests.

1096 Infections:

1097 Some patients taking interferon have had serious infections. Sometimes these infections
1098 have been fatal. If you develop a fever that does not go away or gets higher, call your
1099 healthcare provider right away. Your healthcare provider will need to examine you to rule
1100 out your having a serious infection.

1101 Eye problems:

1102 Changes in vision such as a decrease or loss of vision (blindness) may happen in some
1103 patients. You should have an eye exam before you take PEGASYS. If you have eye
1104 problems or have had them in the past you may need eye exams while you are taking
1105 PEGASYS. Tell your healthcare provider or eye doctor immediately if you have changes
1106 in your vision.

1107 Body organ problems:

1108 Some patients may experience lung problems (such as difficulty breathing or
1109 pneumonia). Certain symptoms like severe stomach pain may mean that your internal
1110 organs are being damaged. Cases of weakness, loss of coordination and numbness due to
1111 stroke have been reported in patients taking PEGASYS/COPEGUS, including patients
1112 with few or no reported risk factors for stroke.

PEGASYS® (peginterferon alfa-2a)

Call your healthcare provider immediately if you develop any of these conditions:

- You become very depressed, think about suicide or injuring/killing another person
- You have severe chest pain
- You have trouble breathing
- You have a change in your vision
- You become pregnant
- You notice unusual bleeding or bruising
- You have psoriasis (a skin disease) and it gets worse while taking PEGASYS
- You have weakness, loss of coordination, numbness or difficulty speaking
- High fever or a fever that does not go away
- You have severe stomach pain or lower back pain
- Bloody diarrhea
- Skin rash can occur in patients taking PEGASYS. In some patients a rash can be serious. If you develop a rash with fever, blisters, or sores in your mouth, nose or eyes or conjunctivitis (red or inflamed eyes, like “pink eye”), stop using PEGASYS and call your doctor right away

For more information on possible side effects with PEGASYS therapy, alone or in combination with COPEGUS, please read the section on “What are the possible side effects of PEGASYS, and PEGASYS taken with COPEGUS?” in this Medication Guide. You should also read the Medication Guide for COPEGUS tablets if you are taking that medicine with PEGASYS.

What is PEGASYS?

PEGASYS is a drug used to treat adults who have a lasting (chronic) infection with hepatitis C virus or hepatitis B virus and who show signs that the virus is damaging the liver. Patients with hepatitis have the virus in their blood and in their liver. PEGASYS reduces the amount of hepatitis C virus in the body and helps the body’s immune system fight the virus. The drug COPEGUS are tablets that may be taken with PEGASYS to help fight the virus infection. Do not take COPEGUS by itself.

In some patients that have received PEGASYS treatment for approximately one year to treat hepatitis C, the amount of the hepatitis virus in the body was decreased to a level so low that it could not be measured by blood tests. After 3 months of therapy, your healthcare provider may ask you to have a blood test to help determine how you are responding to your treatment.

It is not known if PEGASYS, used alone or in combination with COPEGUS, can cure hepatitis (permanently eliminate the virus) or if it can prevent liver failure or liver cancer that is caused by hepatitis infection.

It is also not known if PEGASYS, alone or in combination with COPEGUS, will prevent one infected person from infecting another person with hepatitis.

PEGASYS® (peginterferon alfa-2a)

1154 Who should not take PEGASYS, or PEGASYS with COPEGUS?

1155 Do not take PEGASYS or PEGASYS/COPEGUS therapy if you:

- 1156 • are pregnant, planning to get pregnant during treatment or during the 6 months after
- 1157 treatment or breast-feeding
- 1158 • are a male patient with a female sexual partner who is pregnant or plans to become
- 1159 pregnant at any time while you are being treated with COPEGUS or during the 6
- 1160 months after your treatment has ended
- 1161 • have hepatitis caused by your immune system attacking your liver (autoimmune
- 1162 hepatitis)
- 1163 • have unstable or severe liver disease
- 1164 • had an allergic reaction to another alpha interferon or are allergic to any of the
- 1165 ingredients in PEGASYS or COPEGUS tablets
- 1166 • Do not take PEGASYS, alone or in combination with COPEGUS, if you have
- 1167 abnormal red blood cells such as sickle-cell anemia or thalassemia major.
- 1168

1169 If you have ever had any of the following conditions or serious medical 1170 **problems, tell your healthcare provider before you start taking PEGASYS:**

- 1171 • History of or current severe mental illness (such as depression or anxiety)
- 1172 • History of drug or alcohol addiction or abuse
- 1173 • History of heart disease or previous heart attack
- 1174 • History of cancer
- 1175 • Autoimmune disease (where the body's immune system attacks the body's own
- 1176 cells), such as psoriasis (a skin disease), systemic lupus erythematosus, rheumatoid
- 1177 arthritis
- 1178 • Kidney problems
- 1179 • Blood disorders (bleeding problems)
- 1180 • Diabetes (high blood sugar)
- 1181 • Problems with the thyroid gland
- 1182 • Liver problems, other than hepatitis C or hepatitis B
- 1183 • Colitis (an inflammation of the bowels)
- 1184 • Eye problems
- 1185 • Sleep problems
- 1186 • HIV infection
- 1187 • Organ transplant and are taking medicine that keeps your body from rejecting your
- 1188 transplant (suppresses your immune system)
- 1189

1190 You should tell your healthcare provider if you are taking or planning to take other
1191 prescription or nonprescription medicines or vitamin and mineral supplements or herbal
1192 medicines.

1193 Also tell your healthcare provider if you are taking any of the following medicines:

- 1194 • Theophylline: Your healthcare provider may need to monitor the amount of
- 1195 theophylline in your body and make changes to your theophylline dose.

PEGASYS® (peginterferon alfa-2a)

- HIV medications called nucleoside reverse transcriptase inhibitors (abacavir, didanosine, emtricitabine, lamivudine, tenofovir, stavudine or zidovudine). Some patients developed serious liver problems including death.
- Didanosine: Do not take COPEGUS and didanosine.

If you have any questions about your health condition or about taking PEGASYS alone or in combination with COPEGUS, you should talk to your healthcare provider.

How should I take PEGASYS, or PEGASYS with COPEGUS?

PEGASYS is given by injection under the skin (subcutaneous injection). PEGASYS comes in two different forms (a liquid in a single use vial and a liquid in a prefilled syringe). Your healthcare provider will determine which is best for you. Your healthcare provider will also decide whether you will take PEGASYS alone or with COPEGUS. Your dose of PEGASYS is given as a single injection once per week. At some point, your healthcare provider may change your dose of PEGASYS or COPEGUS. Do not change your dose unless your healthcare provider tells you to change it. It is important that you take PEGASYS and COPEGUS exactly as your healthcare provider tells you. Once you start treatment with PEGASYS, do not switch to another brand of interferon without talking to your healthcare provider. Other interferons may not have the same effect on the treatment of your disease. Switching brands will also require a change in your dose.

Take your prescribed dose of PEGASYS once a week, on the same day of each week and at approximately the same time. Your total dose of COPEGUS tablets should be divided so you take it twice a day with food (breakfast and dinner). Taking half your dose of COPEGUS in the morning and the other half at night will keep the medicine in your body at a steady level. Do not take more than your prescribed dose of PEGASYS or COPEGUS. **Be sure to read the Medication Guide for COPEGUS (ribavirin, USP) for complete instructions on how to take the COPEGUS tablets.**

Your healthcare provider will train you and/or the person that will be giving you the PEGASYS injections on the proper way to give injections. Whether you give yourself the injection or another person gives the injection to you, it is important that you are comfortable with preparing and injecting a dose of PEGASYS, and you understand the instructions in "How do I inject PEGASYS?" **At the end of this guide there are detailed instructions on how to prepare and give yourself an injection of PEGASYS using the form your healthcare provider has prescribed for you.**

If you miss a dose and you remember **within 2 days** of when you should have taken PEGASYS, give yourself an injection of PEGASYS as soon as you remember. Take your next dose on the day you would usually take it. If **more than 2 days** have passed, ask your healthcare provider what you should do. If you miss a dose of COPEGUS, take the missed dose as soon as you remember during the same day. Do not take 2 doses too close together in time. If it is late in the day, wait until the next day and go back on schedule. **Do not double the next dose.**

If you take more than the prescribed amount of PEGASYS, call your healthcare provider right away. Your healthcare provider may want to examine you and take blood for testing.

PEGASYS® (peginterferon alfa-2a)

1239 You must get regular blood tests to help your healthcare provider check how the
1240 treatment is working and to check for side effects.

1241 **What should I avoid while taking PEGASYS, or PEGASYS with COPEGUS?**

- 1242 • If you are pregnant do not start taking or continue taking COPEGUS in combination
1243 with PEGASYS. (See “**What is the most important information I should know**
1244 **about PEGASYS therapy? Problems with Pregnancy**”.)
- 1245 • Avoid becoming pregnant while taking PEGASYS, alone or in combination with
1246 COPEGUS. PEGASYS, alone or in combination with COPEGUS, may harm your
1247 unborn child (death or serious birth defects) or cause you to lose your baby
1248 (miscarry). (See “**What is the most important information I should know about**
1249 **PEGASYS therapy? Problems with Pregnancy**”.)
- 1250 • Do not breast-feed your baby while on PEGASYS, alone or in combination with
1251 COPEGUS.
- 1252 • **Drinking alcohol**, including beer, wine and liquor. This may make your liver disease
1253 worse.
- 1254 • **Taking other medicines**. Take only medicines prescribed or approved by your
1255 healthcare provider. These include prescription and nonprescription medicines and
1256 herbal supplements.

1257 **What are the possible side effects of PEGASYS, and PEGASYS taken with** 1258 **COPEGUS?**

1259 *Also see “What is the most important information I should know about PEGASYS*
1260 *therapy?” in this Medication Guide.*

1261 The possible serious side effects of PEGASYS and PEGASYS/COPEGUS combination
1262 therapy are:

- 1263 • **Mental health problems including depression and suicidal thoughts**
- 1264 • **Blood problems including anemia**. Anemia is a reduction in the number of red
1265 blood cells you have which can be dangerous, especially if you have heart or
1266 breathing problems. Tell your health care provider right away if you feel tired, have
1267 chest pain or shortness of breath. These may be signs of low red blood cell counts.
- 1268 • **Serious infections**
- 1269 • **Stroke**. Some patients may experience weakness, loss of coordination, and numbness
1270 due to stroke.
- 1271 • **Autoimmune problems**: Some patients may develop a disease where the body’s own
1272 immune system begins to attack itself (autoimmune disease) while on PEGASYS
1273 therapy. These diseases can include rheumatoid arthritis, systemic lupus
1274 erythematosus, psoriasis or thyroid problems. In some patients who already have an
1275 autoimmune disease, the disease may worsen while on PEGASYS therapy.
- 1276 • **Heart problems**: PEGASYS may cause some patients to experience chest pain, and
1277 very rarely a heart attack. Patients who already have heart disease could be at greatest
1278 risk. Tell your healthcare provider if you have or have had a heart problem in the past.
- 1279 • **Liver problems**: Some patients may develop worsening of liver function. Some of
1280 the symptoms may include stomach bloating, confusion, brown urine, and yellow
1281 eyes. Tell your healthcare provider immediately if any of these symptoms occur.

PEGASYS® (peginterferon alfa-2a)

- 1282 • **Eye problems** including changes in vision.
- 1283 • **Harm to unborn children.** PEGASYS and PEGASYS/COPEGUS may cause birth
- 1284 defects or death of an unborn child. For more details, see *“What is the most important*
- 1285 *information I should know about PEGASYS therapy?”* in this Medication Guide.
- 1286

1287 Common, but less serious, side effects include:

- 1288 • **Flu-like symptoms:** Most patients who take PEGASYS have flu-like symptoms that
- 1289 usually lessen after the first few weeks of treatment. Flu-like symptoms may include
- 1290 fever, chills, muscle aches, joint pain, and headaches. Taking pain and fever reducers
- 1291 such as acetaminophen or ibuprofen before you take PEGASYS can help with these
- 1292 symptoms. You can also try taking PEGASYS at night. You may be able to sleep
- 1293 through the symptoms.
- 1294 • **Extreme fatigue (tiredness):** Many patients may become extremely tired while on
- 1295 PEGASYS therapy.
- 1296 • **Upset stomach:** Nausea, taste changes, diarrhea, and loss of appetite occur
- 1297 commonly.
- 1298 • **Blood sugar problems:** Some patients may develop a problem with the way their
- 1299 body controls their blood sugar and may develop diabetes.
- 1300 • **Thyroid problems:** Some patients develop changes in the function of their thyroid.
- 1301 Symptoms of thyroid changes include the inability to concentrate, feeling cold or hot
- 1302 all the time, a change in your weight, and changes to your skin.
- 1303 • **Skin reactions:** Some patients may develop redness, swelling, dry or itchy skin at the
- 1304 site of injection. If after several days these symptoms do not disappear, contact your
- 1305 health care provider. You may get a rash during therapy. If this occurs, your health
- 1306 care provider may recommend medicine to treat the rash.
- 1307 • **Hair thinning:** Temporary hair loss is not uncommon during treatment with
- 1308 PEGASYS.
- 1309 • **Trouble sleeping**

1310 These are not all of the side effects of PEGASYS, and PEGASYS taken with COPEGUS.
1311 Your healthcare provider or pharmacist can give you a more complete list. Call your
1312 doctor for medical advice about side effects. Call your doctor for medical advice about
1313 side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also
1314 report side effects to Roche at 1-800-526-6367.

1315 Talk to your healthcare provider if you are worried about side effects or find them very
1316 bothersome.

1317 **General advice about prescription medicines**

1318 Medicines are sometimes prescribed for purposes other than those listed in a Medication
1319 Guide. If you have any concerns or questions about PEGASYS, contact your healthcare
1320 provider. Do not use PEGASYS for a condition or person other than that for which it is
1321 prescribed. If you want to know more about PEGASYS, your healthcare provider or
1322 pharmacist will be able to provide you with detailed information that is written for health-
1323 care providers.

PEGASYS® (peginterferon alfa-2a)

1324 If you are taking COPEGUS (ribavirin, USP) in combination with PEGASYS, also read
1325 the Medication Guide supplied with that medicine.

1326 Keep this and all drugs out of the reach of children.

1327 This Medication Guide has been approved by the US Food and Drug Administration.

1328 MG Revised: April 2009

1329 Medication Guide Appendix: Instructions for Preparing and Giving a Dose with a 1330 PEGASYS® Prefilled Syringe

1331 How should I store PEGASYS Prefilled Syringes?

1332 PEGASYS must be stored in the refrigerator at a temperature of 2°C to 8°C (36°F to
1333 46°F). Do not leave PEGASYS outside of the refrigerator for more than 24 hours. Do not
1334 freeze PEGASYS. Keeping PEGASYS at temperatures outside the recommended range
1335 can destroy the medicine.

1336 Each PEGASYS prefilled syringe can only be used once. Discard after use.

1337 To avoid product foaming, do not shake the prefilled syringe of PEGASYS.

1338 Protect PEGASYS from light during storage.

1339 Keep this and all other medicines out of the reach of children.

1340 How do I prepare and inject PEGASYS?

1341 You should read through all of these directions and ask your healthcare provider for help
1342 if you have any questions before trying to give yourself an injection. It is important to
1343 follow these directions carefully. Talk to your healthcare provider if you have any
1344 questions about PEGASYS.

1345 Your healthcare provider may not want you to take all the medicine that comes in the
1346 prefilled syringe. To appropriately administer the dose that your healthcare provider tells
1347 you to take, you may have to get rid of some of the medicine before injecting the
1348 medicine.

1349 If you ever switch between using prefilled syringes and vials, talk to your healthcare
1350 provider about how much PEGASYS to use. Equal volumes of liquid from the prefilled
1351 syringes and the vials DO NOT contain the same amount of PEGASYS. If you switch
1352 between prefilled syringes and vials, you will have to adjust the volume of liquid that you
1353 use to give your injection. If you do not adjust this, you could accidentally take too much
1354 or too little of your medicine.


1355 If you are giving this injection to someone else, a healthcare provider must teach you how
1356 to avoid needle sticks. Being stuck by a used needle can pass diseases on to you.

1357 The prefilled syringes are used for injecting PEGASYS under the surface of the skin
1358 (subcutaneous).

1359
1360


PEGASYS® (peginterferon alfa-2a)

- 1361 1. Collect all the materials you will need before you start to give the injection:
- 1362 • One PEGASYS prefilled syringe Monthly Convenience Pack containing an
- 1363 inner carton holding the PEGASYS prefilled syringe
- 1364 • A puncture-resistant container for cleaning up when you are finished
- 1365
- 1366 2. Open the convenience pack and look at the contents.
- 1367 • Each convenience pack has everything you need for the PEGASYS injection.
- 1368 – 4 single use syringes filled with medicine (should be colorless to light
- 1369 yellow)
- 1370 – four 27-gauge, ½-inch needles with needle-stick protection device
- 1371 – 4 alcohol swabs
- 1372 3. Take the syringe out of the refrigerator. If there is foam in the solution, put it back in
- 1373 the refrigerator for use at a later time and use another syringe.
- 1374 4. Lay the syringe on a flat clean surface and wait a few minutes until it reaches room
- 1375 temperature. If you notice condensation water on the outside of the syringe, wait
- 1376 another few minutes until it disappears.
- 1377 5. Wash your hands with soap and warm water to prevent infection.
- 1378 6. After the syringe has warmed up, pick it up by the glass barrel and look at it carefully.
- 1379 • Do not use PEGASYS if:
- 1380 – the medicine is cloudy
- 1381 – the medicine has particles floating in it
- 1382 – the medicine is any color besides colorless to light yellow
- 1383 – the expiration date has passed




- 1384
- 1385 7. Attachment of the needle to the PEGASYS prefilled syringe:
- 1386 • Remove the needle from its package. Do not remove the needle shield yet.
- 1387 Keep the needle covered until just before you give the injection.

PEGASYS® (peginterferon alfa-2a)




1388
1389
1390

- Remove and discard the rubber cap from the tip of the syringe barrel.



1391
1392
1393
1394

- Hold the needle close to the hub where the green needle cover connects.
- Put the needle onto the syringe by using an easy twisting motion to tighten the needle onto the syringe.



1395


PEGASYS® (peginterferon alfa-2a)

1396

1397

1398

- Here is a picture of the assembled syringe:



1399

1400

1401

1402

1403

- Keep the syringe in a horizontal position until ready for use.
- If you need to set the syringe down, make sure the plastic needle shield covers the needle. Never let the needle touch any surface.

1404

8. Decide where you will give the injection.


1405

1406

1407

1408

- Pick a place on your stomach or thigh (see the picture below). Avoid your navel and waistline. You should use a different place each time you give yourself an injection.



1409

1410

1411

1412

1413


9. Prepare your skin for the injection.

- To minimize the discomfort from injections, you may want to gently tap the area where you plan to give yourself an injection.
- Clean the area using the alcohol pad. Let the skin dry for 10 seconds.


PEGASYS® (peginterferon alfa-2a)

10. Prepare the syringe for injection.

- Pull the green needle cover back from the needle toward the syringe barrel. The green needle cover will remain in the position you set, do not remove it. This is the needle-stick protection device.



- Hold the syringe-needle assembly tightly at the hub.



- Remove the clear plastic needle shield covering the needle by pulling it straight off.

11. Remove air bubbles from the syringe.


- Hold the syringe with the needle pointing up to the ceiling.
- If you see little bubbles, pull down slightly on the plunger rod.
- Using your thumb and finger, gently tap the syringe to bring air bubbles to the top (small air bubbles may remain on the glass surface).
- Press the plunger in slightly to push air bubbles out of the syringe.
- If you find that there is still small air bubbles on glass surface after you push the air bubbles out, you can still give yourself the injection, the small air bubbles will not hurt you.

12. Dose adjustment.

- Your healthcare provider may not want you to take all the medicine that comes in the prefilled syringe.
- To appropriately administer the dose that your healthcare provider tells you to take, you may have to get rid of some of the medicine before injecting the medicine.

PEGASYS® (peginterferon alfa-2a)


- The syringe has markings for 180 mcg, 135 mcg, and 90 mcg. Your healthcare provider will tell you which mark to use.




- Once you know which mark to use, slowly and carefully press on the plunger rod of the syringe to push out medicine from the syringe. Keep pressing until the edge of the plunger stopper reaches the right mark on the side of the syringe.
- Do not decrease or increase your dose of PEGASYS unless your healthcare provider tells you to.

13. Give the injection of PEGASYS.

- Position the point of the needle (the bevel) so it is facing up.



- Pinch a fold of skin on your stomach or thigh firmly with your thumb and forefinger.




- Hold the syringe like a pencil at a 45° to 90° angle to your skin. In one quick motion, insert the needle as far as it will go into the pinched area of skin. Pull the plunger of the syringe back very slightly. If blood comes into the syringe, the needle has entered a blood vessel. **Do not inject. Withdraw the needle and discard the syringe as outlined in step 11. Repeat the above steps with a new prefilled syringe and prepare a new site.**

PEGASYS® (peginterferon alfa-2a)

1467
1468

- If no blood appears, release your skin and slowly push the plunger all the way down so that you get all of your medicine.



1469

1470

- Pull out the needle at same angle you put it in.

1471

- Wipe the area with an alcohol swab.

1472


14. For safety reasons, before you dispose of the syringe and needle, push the green needle cover toward the needle. Then place the free end of the green cap on a flat surface and push down on it until it clicks and covers over the needle. Always place used syringes and needles in a puncture-resistant container immediately after use and never reuse them. Keep your disposal container out of the reach of children.

1473

1474

1475

1476



1477

1478

1479

How should I dispose of materials used to inject PEGASYS?

1480

There may be special state and local laws for disposal of used needles and syringes. Your healthcare provider or pharmacist should provide you with instructions on how to properly dispose of your used syringes and needles. Always follow these instructions.

1481

1482

1483

The instructions below should be used as a general guide for proper disposal:

1484

- The needles and syringes should never be reused.

PEGASYS® (peginterferon alfa-2a)

- 1485 • Place all used needles and syringes in a puncture-proof disposable container that is
- 1486 available through your pharmacy or healthcare provider (Sharp's container).
- 1487 • DO NOT use glass or clear plastic containers for disposal of needles and syringes.
- 1488 • Dispose of the full container as instructed by your healthcare provider or pharmacist.
- 1489

1490 **DO NOT throw the container in your household trash. DO NOT recycle. Keep the**

1491 **container out of the reach of children.**

1492 MG Appendix: Prefilled Syringe revision date: April 2009

Medication Guide Appendix: Instructions for Preparing and Giving a Dose with a PEGASYS® Vial

How should I store PEGASYS vials?

1496 PEGASYS must be stored in the refrigerator at a temperature of 2°C to 8°C (36°F to

1497 46°F). Do not leave PEGASYS outside of the refrigerator for more than 24 hours. Do not

1498 freeze PEGASYS. Keeping PEGASYS at temperatures outside the recommended range

1499 can destroy the medicine.

1500 Each PEGASYS vial can only be used once. Discard after use.

1501 Do not shake the vial of PEGASYS. If PEGASYS is shaken too hard, it will not work

1502 properly.

1503 Protect PEGASYS from light during storage.

1504 Keep this and all other medicines out of the reach of children.

How do I inject PEGASYS?

1506 The following instructions will help you learn how to measure your dose and give

1507 yourself an injection of PEGASYS. You should read through all of these directions and

1508 ask your healthcare provider for help if you have any questions before trying to give

1509 yourself an injection. It is important to follow these directions carefully. Talk to your

1510 healthcare provider if you have any questions about PEGASYS.

1511 If you are giving an injection to someone else, a healthcare provider must teach you how

1512 to avoid needle sticks. Being stuck by a used needle can pass diseases on to you.

1513 1. Collect all the materials you will need before you start to give the injection:

- 1514 • One vial of PEGASYS
- 1515 • One syringe and needle
- 1516 • Several alcohol pads
- 1517 • A puncture-resistant container to dispose of the needle and syringe when you are
- 1518 finished

1519 2. Check the date on the carton the PEGASYS comes in and make sure the expiration

1520 date has not passed, then remove a vial from the package and look at the medicine.

- 1521 • Do not use PEGASYS if:
- 1522 – the medicine is cloudy
- 1523 – the medicine has particles floating in it

PEGASYS® (peginterferon alfa-2a)

- 1524 – the medicine is any color besides colorless to light yellow
- 1525 – the expiration date has passed
- 1526 3. Warm the refrigerated medicine by gently rolling it in the palms of your hands for
- 1527 about one minute. Do not shake.
- 1528 4. Wash your hands with soap and warm water to prevent infection.
- 1529 5. Take the vial of PEGASYS and flip off the plastic top covering the vial opening, and
- 1530 clean the rubber stopper on the top of the vial with a different alcohol pad.




1531

1532 **If you are not sure how much medicine to use or which mark to use, STOP and call**


1533 **your healthcare provider right away.**

- 1534 6. Remove the needle and syringe from their packaging and attach the needle to the end
- 1535 of the syringe.
- 1536 • Pull the plunger back so the end of it is to the mark on the syringe barrel that
- 1537 matches the dose prescribed for you by your healthcare provider. This will pull air
- 1538 into the syringe barrel.



1539

- 1540 • Push the needle through the center of the stopper on the vial.
- 1541 • Slowly inject all the air from the syringe into the air space above the solution. Do
- 1542 not inject air into the fluid.



1543

- 1544 • Keep the needle inside the vial and turn both upside down. Hold the vial and
- 1545 syringe straight up. Slowly pull back on the plunger until the medicine is in the
- 1546 syringe up to the mark that matches your dose. Make sure the needle tip always
- 1547 stays in the medicine (not in the air space above it).

PEGASYS® (peginterferon alfa-2a)



1548


- 1549
- When the medicine is up to the right mark on the syringe barrel, take the syringe and needle out of the rubber stopper on the vial.
- 1550
- 1551
- Keep the syringe pointing up until you are ready to use it.
- 1552
- If you need to set the syringe down, make sure that you never let the needle touch any surface.
- 1553

1554 7. Remove air bubbles from the syringe.

- 1555
- Hold the syringe with the needle pointing up to the ceiling.
- 1556
- Using your thumb and finger, tap the syringe to bring air bubbles to the top.
- 1557
- Press the plunger in slightly to push air bubbles out of the syringe.

1558 8. Decide where you will give the injection.

- 1559
- Pick a place on your stomach or thigh (see the picture below). Avoid your navel and waistline. You should use a different place each time you give yourself an injection.
- 1560
- 1561




1562

1563 9. Prepare your skin for the injection.

- 1564
- To minimize the discomfort from injections, you may want to gently tap the area where you plan to give yourself an injection.
- 1565
- Clean the area using an alcohol pad. Let the skin dry for 10 seconds.
- 1566
- 1567

1568 10. Give the injection of PEGASYS.

- 1569
- Position the point of the needle (the bevel) so it is facing up.
- 1570




1571

- 1572
- Pinch a fold of skin on your stomach or thigh firmly between your thumb and forefinger.
- 1573


PEGASYS® (peginterferon alfa-2a)

1574



1575

- 1576
- 1577
- 1578
- 1579
- 1580
- 1581
- 1582
- 1583
- 1584
- Hold the syringe like a pencil at a 45° to 90° angle to your skin. In one quick motion, insert the needle as far as it will go into the pinched area of skin. Pull the plunger of the syringe back very slightly. If blood comes into the syringe, the needle has entered a blood vessel. **Do not inject. Withdraw the needle and discard the syringe as outlined in step 11. Repeat the above steps with a new vial and syringe and prepare a new site.**
 - If no blood appears, release your skin and slowly push the plunger all the way down so that you get all of your medicine.



1585

- 1586
- Pull out the needle at same angle you put it in. Wipe the area with an alcohol pad.
- 1587
- 1588
- 1589
- 1590
- 1591
11. For safety reasons, always place used syringes and needles in a puncture-resistant container immediately after use and never reuse them.
- If you are using a syringe with a needle-stick protection device, before you dispose of the syringe and needle, place the free end of the green cap on a flat surface and push down on it until it clicks and covers over the needle.

1592 **How should I dispose of materials used to inject PEGASYS?**

1593 There may be special state and local laws for disposal of used needles and syringes. Your

1594 healthcare provider or pharmacist should provide you with instructions on how to

1595 properly dispose of your used syringes and needles. Always follow these instructions.

1596 The instructions below should be used as a general guide for proper disposal:

- 1597
- 1598
- 1599
- 1600
- 1601
- 1602
- The needles and syringes should never be reused.
 - Place all used needles and syringes in a puncture-proof disposable container that is available through your pharmacy or healthcare provider (Sharp's container).
 - DO NOT use glass or clear plastic containers for disposal of needles and syringes.
 - Dispose of the full container as instructed by your healthcare provider or pharmacist.

1603 **DO NOT throw the container in your household trash. DO NOT recycle. Keep the**

1604 **container out of the reach of children.**

PEGASYS® (peginterferon alfa-2a)

1605 MG Appendix: Vial revision date: October 2008



Hoffmann-La Roche Inc.
340 Kingsland Street
Nutley, New Jersey 07110-1199
(US Govt. Lic. No. 0136)

1606

1607 XXXXXXXXX

1608 XXXXXXXXX

1609 Copyright © 2003-200X by Hoffmann-La Roche Inc. All rights reserved.