Gyne-Lotrimin 3 3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
Vaginal Cream
3 Disposable Applicators
Cures most vaginal yeast infections
Drug Facts
Active ingredients
Purpose
Clotrimazole 2% (100 mg in each applicatorful) ..... Vaginal antifungal
Clotrimazole 2% (external cream) ..... Vaginal antifungal
Uses ▪ treats vaginal yeast infections ▪ relieves external itching and irritation due to a vaginal yeast infection
Warnings
For vaginal use only
Do not use if you have never had a vaginal yeast infection diagnosed by a doctor
Ask a doctor before use if you have
▪ vaginal itching and discomfort for the first time
▪ lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.
▪ vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
▪ been exposed to the human immunodeficiency virus (HIV) that causes AIDS
When using this product
▪ do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
▪ do not have vaginal intercourse ▪ mild increase in vaginal burning, itching or irritation may occur
▪ if you do not get complete relief ask a doctor before using another product
Stop use and ask a doctor if
▪ symptoms do not get better in 3 days ▪ symptoms last more than 7 days
▪ you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions ▪ before using this product read the enclosed educational brochure for complete directions and information
▪ adults and children 12 years of age and over:
▪ vaginal cream: insert one applicatorful of cream into the vagina at bedtime for 3 days in a row. Throw applicator away after use.
▪ external cream: use the same tube of cream if you have itching and irritation on the skin outside the vagina. Squeeze a small amount of cream onto your fingertip. Apply to itchy, irritated skin outside the vagina. Use 2 times daily for up to 7 days as needed.
▪ children under 12 years of age: ask a doctor
© Copyright & Distributed by Schering-Plough HealthCare Products, Inc., Memphis, TN 38151 USA. All rights reserved. Product of Italy.
36414-01
Schering-Plough
recyclable carton
135
Drug Facts (continued)
Other information ▪ store between 20° to 25°C (68° to 77°F)
▪ see end panel of carton and tube crimp for lot number and expiration date
▪ tamper-evident: do not use if seal embossed with "S" over tube opening is missing, open or broken
Inactive ingredients benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-octyldodecanol, polysorbate 60, purified water, sorbitan monostearate
Questions or comments? If you have questions or need more information on this product, call toll free 1-800-317-2165 between 8:00 AM and 5:00 PM Central Standard Time, Monday through Friday.
Educational Brochure Enclosed
One 21g (0.74 oz) Tube of Clotrimazole Vaginal Cream (2%) and 3 Disposable Applicators
Gyne-Lotrimin 3 3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
Vaginal Cream
Cures most vaginal yeast infections
Gyne-Lotrimin 3 3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
3 Disposable Applicators
Vaginal Cream
Cures most vaginal yeast infections
Gyne-Lotrimin 3 3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
3 Disposable Applicators
Vaginal Cream
Gyne-Lotrimin 3 3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
3 Disposable Applicators
Vaginal Cream

Gyne-Lotrimin 3
Vaginal Cream
3-Day Treatment
Clotrimazole Vaginal Cream (2%)
Vaginal Antifungal
Net Weight 21g (0.74 oz)
Active Ingredients
Purpose
Clotrimazole 2% (100 mg in each applicatorful) ..... Vaginal antifungal
Clotrimazole 2% (external cream) ..... Vaginal antifungal
Uses = treats vaginal yeast infections = relieves external itching and irritation due to a vaginal yeast infection
Warnings For vaginal use only. Do not use if you have never had a vaginal yeast infection diagnosed by a doctor. Read the Warnings and Directions on the carton and in the educational brochure before using this product. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions: To Open: Unscrew the cap, use pointed end on cap to break seal. Fill the applicator and insert one applicatorful of cream into vagina at bedtime for 3 days in a row.
Other Information = store between 20° to 25°C (68° to 77°F) = see end panel of carton and tube crimp for lot number and expiration date = tamper-evident: do not use if seal embossed with "S" over tube opening is missing, open or broken
Questions or comments?
If you have questions or need more information on this product, call toll free 1-800-317-2165 between 8:00 AM and 5:00 PM Central Standard Time, Monday through Friday.
© Copyright & Distributed by Schering-Plough HealthCare Products, Inc., Memphis, TN 38151 USA.
All rights reserved. Product of Italy. LPK-4005-3-0508-2 147
23892-04
147
.9655"
.9315"
3.8125" Repeat Length

PAMPHLET PRINTS PMS 281

EDUCATIONAL BROCHURE

Read This Before Using

Gyne-Lotrimin 3®

Clotrimazole Vaginal Cream (2%) Vaginal Antifungal

Clotrimazole 2% (100 mg per applicator)

3-day treatment of vaginal yeast infections.

Schering-Plough

09679-99 LPK-9010-3, 0508-2.54

Use:

- For the treatment of vaginal yeast infections.

If you have symptoms such as vaginal itching, burning, or discharge, and, if at some time in the past your doctor has told you that these symptoms were due to a vaginal yeast infection, then Gyne-Lotrimin 3® Vaginal Cream should work for you. **If, however, you have never had these symptoms or a vaginal yeast infection before, you should see your doctor before using this product.**

What is a Vaginal Yeast Infection?

A yeast infection is a common type of vaginal infection. Your doctor may call it candidiasis. This condition is caused by an organism called Candida, which is a type of yeast. Healthy women usually have this yeast on the skin, in the mouth, in the digestive tract, and in the vagina. At times, the yeast can grow very quickly. In fact, the infection is sometimes called yeast (Candida) "overgrowth." Some women may have a yeast infection on the skin outside the vagina (vulva) associated with the vaginal infection.

A yeast infection can occur at any age. It is most common during the childbearing years. Women who are pregnant or diabetic, taking antibiotics, birth control pills or steroids, or who have a weakened immune system are more likely to get repeated yeast infections.

Various medical conditions can damage the body's normal defenses against infection. One of the most serious of these conditions is infection with the human immuno-deficiency virus (HIV-the virus that causes AIDS). The HIV virus causes the body to be more likely to get infections, including vaginal yeast infections. Therefore, women with HIV infection may have frequent vaginal yeast infections that do not clear up easily with proper treatment. If you may have been exposed to HIV and are experiencing repeated vaginal yeast infections, you should see your doctor right away. For additional information on HIV infection, please contact your doctor or the CDC National AIDS HOT-LINE. The CDC phone numbers are 1-800-CDC-INFO (1-800-232-4636) for English or Spanish or 1-800-232-6348 (hearing impaired, TDD/TTY).

Symptoms of a Vaginal Yeast Infection

There are many symptoms of a vaginal yeast infection. They can include:

- Vaginal itching (ranging from mild to severe).
- A clumpy, vaginal discharge that may look like cottage cheese.
- Vaginal soreness, irritation, or burning.
- Rash or redness on the skin outside the vagina (vulvar irritation).

Note: Vaginal discharge that is different from above, for example, a yellow/green discharge or a foul-smelling discharge may mean that you have a different infection. If this is the case, you should talk to your doctor before using Gyne-Lotrimin 3® Vaginal Cream.

Warnings

For vaginal use only.

Do not use if you have never had a vaginal yeast infection diagnosed by a doctor.

Ask a doctor before use if you have

- **vaginal itching and discomfort for the first time**
- **lower abdominal, back or shoulder pain, fever, chills, nausea, vomiting, or foul-smelling vaginal discharge. You may have a more serious condition.**
- vaginal yeast infections often (such as once a month or 3 in 6 months). You could be pregnant or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS.

When using this product

- do not use tampons, douches, spermicides, or other vaginal products. Condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).
- do not have vaginal intercourse
- mild increase in vaginal burning, itching or irritation may occur
- if you do not get complete relief ask a doctor before using another product

Stop use and ask a doctor if

- **symptoms do not get better in 3 days**
- **symptoms last more than 7 days**
- **you get a rash or hives, abdominal pain, fever, chills, nausea, vomiting, or a foul-smelling vaginal discharge**

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Tamper-evident: do not use if seal embossed with "3" over tube opening is missing, open or broken.

For Best Results

1. Be sure to use this product for the full 3 days, even if your symptoms go away sooner.
2. Use one applicatorful of Gyne-Lotrimin 3® Vaginal Cream at bedtime for most days in a row, even during your menstrual period.
3. Wear cotton underwear.
4. If your partner has itching, redness or discomfort at his penis, he should consult his doctor. He should tell the doctor that you are treating yourself for a vaginal yeast infection.
5. Dry the outside vaginal area thoroughly after a shower, bath or swim. Change out of wet bathing suit or damp clothes as soon as possible. A dry area is less likely to lead to the growth of yeast.
6. Wipe from front to rear after a bowel movement or urination.
7. Do not douche unless your doctor tells you to do so. Douching may wash the product out of the vagina.
8. Try not to scratch the skin outside the vagina. Scratching can cause more irritation and can spread the infection.
9. Discuss with your doctor any drugs you are now taking. You are more likely to get a vaginal yeast infection with certain drugs.
10. Do not have vaginal intercourse while using this product.
11. When using this product condoms and diaphragms may be damaged and fail to prevent pregnancy or sexually transmitted diseases (STDs).

If you have a Question

Questions or a medical nature should be taken up with your doctor. If you have any other questions, or need more information on this product, call our TOLL-FREE Number at 1-800-317-2165, between 8:00 AM and 5:00 PM Central Standard Time, Monday through Friday.

Active Ingredient:

Clotrimazole 2% (100 mg per applicator)

Inactive Ingredients:

Benzyl alcohol, cetostearyl alcohol, cetyl esters wax, 2-ethyldodecanol, polysorbate 60, purified water, sorbitan monostearate.

Storage

Store at 20° to 25°C (68° to 77°F). See end flap of carton and end of tube for lot number and expiration date.

How to use Gyne-Lotrimin 3® Vaginal Cream with 3 Disposable Applicators

Directions for Use

Use the cream at bedtime for 3 days in a row.

Read the full directions below before using.

1. To open the tube, unscrew the cap.

Press the sharp point of the cap into the sealed end of the tube. Push down firmly until the seal is broken. (see picture.)



2. Attach the applicator to the tube of cream by placing end of the applicator firmly onto tube of cream. (see picture.)

3. Gently squeeze opposite end of tube, forcing cream into barrel of applicator. The plunger will stop moving outward when the barrel is full. Separate the applicator from the tube. After each use, replace cap and roll up the tube from the bottom.

4. Hold the barrel as shown in the picture.

5. The Gyne-Lotrimin 3® Vaginal Cream applicator is inserted into the vagina in much the same way as a tampon. Gently ease the applicator into the vagina as far back as it will go comfortably. This can be done while lying on your back with your knees bent (as shown in the picture), or while standing with your feet apart and your knees bent.

6. With one hand holding the barrel, use the other hand to push the plunger into the barrel as far as it will go. Remove the applicator from the vagina.

7. Throw the applicator away after each use. Do not flush down the toilet.

8. Lie down as soon as possible to keep the cream from leaking out.

9. Repeat steps 2 through 8 before bedtime for the next 2 days.

To protect your clothing from leakage or discharge, you may want to use deodorant-free pads or panty shields while you are using Gyne-Lotrimin 3® Vaginal Cream. **Do not use tampons, douches, spermicides or other vaginal products** until after you have completed the treatment and your symptoms are gone.

© Copyright & Distributed by Schering-Plough HealthCare Products, Inc., Memphis, TN 38151 USA.
All rights reserved. Product of Italy

APO #: P0384-00
10.12.00