## This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda

540	MEDICATION GUIDE
541	
542	$\mathbf{VOTRIENT}^{^{\mathrm{TM}}}$ (VO-tree-ent)
543	(pazopanib)
544	tablets
545	Read the Medication Guide that comes with VOTRIENT before you start taking it and each time
546	you get a refill. There may be new information. This Medication Guide does not take the place of
547	talking with your healthcare provider about your medical condition or treatment.
548	
549	What is the most important information I should know about VOTRIENT?
550	• VOTRIENT can cause serious liver problems including death. Your healthcare provider
551	will do blood tests to check your liver before you start and while you take VOTRIENT.
552	Tell your healthcare provider right away if you have any of these signs of liver problems:
553	<ul> <li>yellowing of your skin or the whites of your eyes (jaundice)</li> </ul>
554	• dark urine
555	• tiredness
556	<ul> <li>nausea or vomiting</li> </ul>
557	• loss of appetite
558	<ul> <li>pain on the right side of your stomach</li> </ul>
559	• bruise easily
560	
561	What is VOTRIENT?
562	VOTRIENT is a prescription medicine used to treat advanced renal cell cancer in adults.
563	It is not known if VOTRIENT is safe or effective in children under 18 years of age.
564	
565	What should I tell my healthcare provider before taking VOTRIENT?
566	Before you take VOTRIENT, tell your healthcare provider if you:
567	<ul> <li>have or had liver problems</li> </ul>
568	<ul> <li>have high blood pressure</li> </ul>
569	have heart problems or an irregular heartbeat including QT prolongation
570	<ul> <li>have a history of a stroke</li> </ul>
571	• have coughed up blood in the last 6 months
572	• had bleeding of your stomach or intestines in the last 6 months
573	• have a history of a tear (perforation) in your stomach or intestine, or an abnormal connection
574	between two parts of your gastrointestinal tract (fistula)
575	have thyroid problems
576	had recent surgery (within the last 7 days) or are going to have surgery
577	have any other medical conditions  NOTPIENT
578	are pregnant or plan to become pregnant. VOTRIENT can harm your unborn baby. You
579	should not become pregnant while you are taking VOTRIENT.

## This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda

580	• are breast-feeding or plan to breast-feed. It is not known if VOTRIENT passes into your
581	breast milk. You and your healthcare provider should decide if you will take VOTRIENT or
582	breast-feed. You should not do both.
583	
584	Tell your healthcare provider about all the medicines you take including prescription and
585	non-prescription medicines, vitamins, and herbal supplements. VOTRIENT may affect the way
586	other medicines work and other medicines may affect how VOTRIENT works.
587	
588	Especially, tell your healthcare provider if you:
589	<ul> <li>take medicines that can affect enzyme metabolism such as:</li> </ul>
590	<ul> <li>some antibiotics (used to treat infections)</li> </ul>
591	<ul> <li>some drugs used to treat HIV</li> </ul>
592	<ul> <li>some drugs used to treat depression</li> </ul>
593	<ul> <li>drugs used to treat irregular heart beats</li> </ul>
594	drink grapefruit juice
595	
596	Ask your healthcare provider if you are not sure if your medicine is one that is listed above.
597	
598	Know the medicines you take. Keep a list of them and show it to your healthcare provider and
599	pharmacist when you get a new medicine.
600	
601	How should I take VOTRIENT?
602	• Take VOTRIENT exactly as your healthcare provider tells you. Your healthcare provider
603	will tell you how much VOTRIENT to take.
604	Your healthcare provider may change your dose.
605	• Take VOTRIENT on an empty stomach, at least 1 hour before or 2 hours after food.
606	• Do not crush tablets of VOTRIENT.
607	• Do not eat grapefruit or drink grapefruit juice while taking VOTRIENT. They may change
608	the amount of VOTRIENT in your body.
609	• If you miss a dose, take it as soon as you remember. Do not take it if it is close (within
610	12 hours) to your next dose. Just take the next dose at your regular time. Do not take more
611	than 1 dose of VOTRIENT at a time.
612	• Your healthcare provider will test your blood and heart before you start and while you take
613	VOTRIENT.
614	
615	What are the possible side effects of VOTRIENT?
616	VOTRIENT may cause serious side effects including:

• See "What is the most important information I should know about VOTRIENT?".

617

618

619 high blood pressure including a sudden and severe rise in blood pressure which may be 620 life-threatening 621 • irregular or fast heartbeat or fainting 622 • heart attack or stroke: chest pain or pressure, pain in your arms, back, neck or jaw, 623 shortness of breath, numbness or weakness on one side of your body, trouble talking, 624 headache, or dizziness. 625 • **bleeding problems:** unusual bleeding, bruising, wounds that do not heal. 626 • tear in your stomach or intestinal wall (perforation) or bleeding: pain, swelling in your 627 stomach-area, vomiting blood, and black sticky stools. 628 thyroid problems 629 630 Call your healthcare provider right away, if you have any of the symptoms listed above. 631 632 The most common side effects of VOTRIENT include: 633 diarrhea 634 • change in hair color 635 • nausea or vomiting 636 • loss of appetite • feeling tired 637 638 639 640 Tell your healthcare provider if you have any side effect that bothers you or that does not go 641 away. These are not all the possible side effects of VOTRIENT. For more information, ask your 642 healthcare provider or pharmacist. 643 644 Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-645 800-FDA-1088. 646 647 How should I store VOTRIENT tablets? 648 Store VOTRIENT between 59°F to 86°F (15°C to 30°C). 649 650 Keep VOTRIENT and all medicines out of the reach of children. 651 652 General information about the safe and effective use of VOTRIENT: Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. 653 654 Do not use VOTRIENT for a condition for which it was not prescribed. Do not give VOTRIENT 655 to other people even if they have the same symptoms that you have. It may harm them. 656 This Medication Guide summarizes the most important information about VOTRIENT. If you 657 658 would like more information, talk with your healthcare provider. You can ask your pharmacist or

## This label may not be the latest approved by FDA. For current labeling information, please visit https://www.fda.gov/drugsatfda

659	healthcare provider for information about VOTRIENT that is written for healthcare
660	professionals. For more information, go to www.VOTRIENT.com or call 1-888-825-5249.
661	
662	What are the ingredients in VOTRIENT?
663	Active Ingredient: pazopanib.
664	Inactive Ingredients: hypromellose, iron oxide black, macrogol/polyethylene glycol 400
665	(PEG 400), magnesium stearate, microcrystalline cellulose, polysorbate 80, povidone, sodium
666	starch glycolate, titanium dioxide.
667	
668	VOTRIENT is a trademark of GlaxoSmithKline.
669	
670	This Medication Guide has been approved by the U.S. Food and Drug Administration.
671	
	gSK GlaxoSmithKline
672	
673	GlaxoSmithKline
674	Research Triangle Park, NC 27709
675	
676	©YEAR, GlaxoSmithKline. All rights reserved.
677	
678	Revised: Month YEAR
679	VTR:XMG