

Salonpas® Pain Relief Patch LARGE

Menthol 3%, Methyl salicylate 10%

3 patches

Hisamitsu®

LARGE

NDC#46581-675-03

for temporary relief of mild to moderate pain

- Backache
- Sore Muscle
- Strains
- Bruises



Salonpas® Pain Relief Patch

APPLY FOR
8-12
HOURS

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

3 LARGE patches

3 1/16" X 5 1/2" (10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

Salonpas® Pain Relief Patch LARGE
Menthol 3%, Methyl Salicylate 10%

3 patches

press open

minty scent

Salonpas® Pain Relief Patch LARGE
Menthol 3%, Methyl salicylate 10%

Drug Facts

Active ingredients (in each patch)

Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding.

- The chance is small but higher if you:
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

- Do not use
- on the face or rashes
 - with a heating pad
 - any patch from a pouch that has been open for 14 or more days
 - if allergic to aspirin or other NSAIDs
 - when sweating (such as from exercise or heat)
 - right before or after heart surgery
- Ask a doctor before use if
- you are allergic to topical products
 - you are taking a diuretic
 - the stomach bleeding warning applies to you
 - you have high blood pressure, heart disease, or kidney disease
- When using this product
- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
 - the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

- Adults 18 years and older:
- only use one patch at a time
 - clean and dry affected area
 - apply one patch to the affected area and leave in place for up to 8 to 12 hours
 - if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 - do not use more than 2 patches per day
 - do not use for more than 5 days in a row
 - the used patch should be removed from the skin when a new one is applied
 - Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

aliphatic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene styrene block copolymer, syntretic, aluminum silicate

Questions or comments?

Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST) | www.salonpas.us

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN

3 patches
Salonpas®
Pain Relief Patch LARGE
Menthol 3%, Methyl salicylate 10%

3 46581 67503
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503
© Registered Trade Mark

LOT
EXP.

Salonpas® Pain Relief Patch LARGE
Menthol 3%, Methyl salicylate 10%

3 patches

IM289T-0

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



3 LARGE patches
3 1/8" x 5 1/2" (10cm x 14cm)
Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

✂ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®
Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA-TOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503

LOT
EXP
Package not child resistant.
Keep out of reach of children.
© : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Pain Relief Patch LARGE Menthol 3%, Methyl salicylate 10% mildly scented

Active ingredients (in each patch)
Menthol 3%
Methyl salicylate 10% (NSAID : nonsteroidal anti-inflammatory drug)
Purpose
Topical analgesic
Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings
For external use only
Stomach bleeding warning:
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery
Ask a doctor before use if
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes.
If eye contact occurs, rinse thoroughly with water.
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky sides together.

Directions
Adults 18 years and older:
■ clean and dry affected area ■ only use one patch at a time
■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ remove patch from backing film and apply to skin (see illustration)
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information
■ avoid storing product in direct sunlight
■ store at 20°-25°C (68°-77°F)
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients alkylic, saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-877-SALONPAS, weekdays, 9AM to 5PM (PST) ■ www.salonpas.us ■ 1-7721-0

Salonpas[®] Pain Relief Patch **LARGE**

Menthol 3%, Methyl Salicylate 10%


2 patches

NDC#46581-675-02

Hisamitsu[®]
LARGE

for temporary relief of mild to moderate pain

- Backache
- Sore Muscle
- Strains
- Bruises



Salonpas[®] Pain Relief Patch

APPLY FOR
8-12
HOURS

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

2 **LARGE** patches

3 15/16" X 5 1/2" (10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

Salonpas[®] Pain Relief Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%

Salonpas[®] Pain Relief Patch **LARGE** patches
Menthol 3%, Methyl Salicylate 10%



Salonpas[®] Pain Relief Patch **LARGE**
Menthol 3%, Methyl Salicylate 10%

press
open

minty scent

Drug Facts

Active ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ strains ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use:
■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
■ with a heating pad ■ when sweating (such as from exercise or heat) ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery
Ask a doctor before use if:
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product:
■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
■ symptoms clear, up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
■ only use one patch at a time
■ clean and dry affected area ■ remove patch from backing film and apply to skin (see illustration)
■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied
Children under 18 years of age: do not use

Other information

■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST) www.salonpas.us

Manufactured by Hisamitsu Pharmaceutical Co., Inc. TORRANCE, CA 90503
Distributed by Hisamitsu America, Inc. TORRANCE, CA 90503
Menthol 3%, Methyl Salicylate 10%
JAPAN SAGA TOSU
MADE IN JAPAN
© Registered Trade Mark
3 46581 67502 3



LOT
EXP

2 patches

IM288T-0

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



Menthol 3%
Methyl salicylate 10%
Topical Analgesic

2 LARGE Patches
3 1/16" X 5 1/2"
(10cm X 14cm)

✂ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503
EXP. LOT
Package not child resistant.
Keep out of reach of children.
© : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Pain Relief Patch LARGE Menthol 3%, Methyl salicylate 10% minty scent

Drug Facts
Active ingredients (in each patch)
Menthol 3% Topical analgesic
Methyl salicylate 10% (NSAID) Nonsteroidal anti-inflammatory drug
..... Topical analgesic

Warnings
For external use only
Stomach bleeding warning:
This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ on the face or rashes ■ on wounds or damaged skin ■ If allergic to aspirin or other NSAIDs
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease
When using this product, ■ wash hands after applying or removing patch. Avoid contact with eyes.
If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
■ clean and dry affected area ■ only use one patch at a time
■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ remove patch from backing film and apply to skin (see illustration)
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied
Children under 16 years of age: do not use


Other information
■ avoid storing product in direct sunlight
■ store at 20-25°C (68-77°F)
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients allylicyclic saturated hydrocarbon resin, backing cloth film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST) www.salonpas.us
1L7711-0

Salonpas®

Chisamitsu®



Salonpas®

Salonpas®

Pain Relief Patch

APPLY FOR
8-12
HOURS



STRETCHABLE & FLEXIBLE

1 LARGE patch
3 1/4" X 5 1/2" (10cm X 14cm)

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

Salonpas®

Salonpas®

Pain Relief P

1 LARGE patch X **12** sachets
3 1/4" X 5 1/2" (10cm X 14cm)

303 46581 6/507 1

Hisamitsu®
LARGE

for temporary relief of mild to moderate pain

Backache
Sore Muscle
Sprains
Bruises

minly scent

NDC # 46581-075-01

1 Peel the center film.
2 Place on affected area.
3 Slide a film.
4 Slide the other film.

Salonpas®
Pain Relief Patch

**Menthyl 3%
Methyl Salicylate 10%
Topical Analgesic**

APPLY FOR 8-12 HOURS

1 LARGE patch
3 1/8" x 5 1/2" (10 cm x 14 cm)

DO NOT USE MORE THAN ONE SALONPAS PAIN RELIEF PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant. Keep out of reach of children.

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

Manufactured by Hisamitsu Pharmaceutical Co., Inc. Hisamitsu America, Inc. JAPAN SAGA TOYO Torrance, CA 90503
Distributed by Hisamitsu America, Inc. Torrance, CA 90503
MADE IN JAPAN

3 46581 07501 6

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Drug Facts

Active ingredients (in each patch)

Menthyl 3% Topical Analgesic
Methyl Salicylate 10% (NSAID) : nonsteroidal anti-inflammatory drug..... Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
 ■ sprains ■ simple backache ■ arthritis ■ bruises

Warnings
For external use only
 Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
 ■ have 3 or more alcoholic drinks every day while using this product
 ■ take more or for a longer time than directed

Do not use
 ■ on the face or rashes ■ on wounds or damaged skin ■ if allergic to aspirin or other NSAIDs
 ■ with a heating pad ■ when sweating (such as from exercise or heat)
 ■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
 ■ you are allergic to topical products ■ the stomach bleeding warning applies to you
 ■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product ■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 ■ stomach pain or upset gets worse or lasts ■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 ■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
 ■ symptoms clear up and occur again within a few days

If pregnant or try breast-feeding ■ ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If not in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
 ■ clean and dry affected area ■ only use one patch at a time
 ■ apply one patch to the affected area and leave in place for up to 8 to 12 hours ■ remove patch from backing film and apply to skin (see illustration)
 ■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
 ■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
 ■ the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information ■ store at 20-25°C (68-77°F) ■ avoid storing product in direct sunlight
 ■ protect product from excessive moisture
 ■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients
 alkylic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
 Toll free 1-877-SALONPAS, weekdays 9AM to 5PM (PST)
 www.salonpas.us

LOT
EXP

© : Registered Trade Mark 1L770T-0

Hisamitsu
LARGE

FREE SAMPLE NOT FOR RE-SALE

minty scent

NDC#45981-07-1-99

for temporary relief of mild to moderate pain

Backache
Sore Muscle
Bruises

APPLY FOR 8-12 HOURS

1 LARGE patch
3 1/8" x 5 1/8" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Salonpas

Pain Relief Patch

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

1 Peel the center film.

2 Place on affected area.

3 Slide a film.

4 Slide the other film.

Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAKA TOSU MADE IN JAPAN

Distributed by Hisamitsu America, Inc. Torrance, CA 90503

Package not child resistant. Keep out of reach of children.

X CUT OPEN HERE ----- CUT OPEN HERE X

Drug Facts

Active ingredients (in each patch)

Menthol 3% Topical analgesic
Methyl salicylate 10% (NSAID) nonsteroidal anti-inflammatory drug
Methyl salicylate 10% (NSAID) nonsteroidal anti-inflammatory drug
Methyl salicylate 10% (NSAID) nonsteroidal anti-inflammatory drug

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- strains
- sprains
- simple backache
- arthritis
- bruises

Purpose

Topical analgesic
Topical analgesic

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on wounds or damaged skin
- with a healing pad
- when sweating (such as from exercise or heat)
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery

Ask a doctor before use if

- you are allergic to topical products
- the stomach bleeding warning applies to you
- you are taking a diuretic
- you have high blood pressure, heart disease, or kidney disease

When using this product

- wash hands after applying or removing patch. Avoid contact with eyes.
- if eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- stomach pain or upset gets worse or lasts
- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:

- only use one patch at a time
- clean and dry affected area
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-877-SALONPAS. Weekdays 9AM to 5PM (PST)
www.salonpas.us

LOT

EXP.

© : Registered Trade Mark 1L789T-0

Salonpas[®] Arthritis Pain Patch LARGE

Menthol 3%, Methyl Salicylate 10%

3 patches

Hisamitsu[®]

LARGE

for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains

NDC#46581-685-03

3 Salonpas[®] Arthritis Pain Patch LARGE
Menthol 3%, Methyl Salicylate 10%

Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAGA TOSU
MADE IN JAPAN

Distributed by Hisamitsu America, Inc. Torrance, CA 90503
© : Registered Trade Mark

3 46581 68503 9

3 Salonpas[®] Arthritis Pain Patch LARGE
Menthol 3%, Methyl Salicylate 10%

3 patches

press open

minty scent

Hisamitsu[®]

LARGE

for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains

NDC#46581-685-03

3 Salonpas[®] Arthritis Pain Patch LARGE
Menthol 3%, Methyl Salicylate 10%

Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAGA TOSU
MADE IN JAPAN

Distributed by Hisamitsu America, Inc. Torrance, CA 90503
© : Registered Trade Mark

3 46581 68503 9

Salonpas[®] Arthritis Pain Patch

APPLY FOR 8-12 HOURS

Menthol 3%
Methyl Salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

3 LARGE patches

3 1/8" x 5 1/2" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant.
Keep out of reach of children.

LOT EXP

Salonpas[®] Arthritis Pain Patch LARGE
Menthol 3%, Methyl Salicylate 10%

3 patches



1 Peel the center film.



2 Place on affected area.



3 Slide the film upward.



4 Slide the other film downward.



Drug Facts

Active ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10% (NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses

Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

Warnings

For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on the face or rashes
- on wounds or damaged skin
- with a heating pad
- any patch from a pouch that has been open for 14 or more days
- if allergic to aspirin or other NSAIDs
- when sweating (such as from exercise or heat)
- right before or after heart surgery

When using this product

- wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:

- only use one patch at a time
- remove patch from backing film and apply to skin (see illustration)
- clean and dry affected area
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

- avoid storing product in direct sunlight
- store at 20-25°C (68-77°F)
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, poly(isobutylene), poly(isobutylene 1,200,000), styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-87-SALONPAS, Weekdays, 9AM to 5PM (PST) www.salonpas.us

1M2937-0

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



3 LARGE patches
3 1/4" x 5 1/2"
(10cm x 14cm)

**Menthol 3%
Methyl salicylate 10%
Topical Analgesic**

FDA APPROVED OVER THE COUNTER PAIN RELIEVING PATCH.
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

✂ DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Hisamitsu®

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
JAPAN SAGA TOSU
MADE IN JAPAN
Distributed by
Hisamitsu America, Inc.
Torrance, CA 90503

LOT
EXP

Package not child resistant.
Keep out of reach of children.

® : Registered Trade Mark

✂ CUT OPEN HERE ----- CUT OPEN HERE ✂

Salonpas® Arthritis Pain Patch LARGE Menthol 3%, Methyl salicylate 10% mintily scented

Drug Facts

Active ingredients (in each patch)

Menthol 3% Topical analgesic
Methyl salicylate 10% (NSAID : nonsteroidal anti-inflammatory drug) Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on the face or rashes
- with a healing pad
- any patch from a pouch that has been open for 14 or more days
- if allergic to aspirin or other NSAIDs
- when sweating (such as from exercise or heat)
- right before or after heart surgery

Ask a doctor before use if

- you are allergic to topical products
- you are taking a diuretic
- you are taking a diuretic
- you have high blood pressure, heart disease, or kidney disease
- you are taking a diuretic

When using this product

- wash hands after applying or removing patch. Avoid contact with eyes.
- if eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- stomach pain or upset gets worse or lasts
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days

Stop use and ask a doctor if

- symptoms clear up, and occur again within 14 days
- symptoms clear up, and occur again within 14 days
- symptoms clear up, and occur again within 14 days
- symptoms clear up, and occur again within 14 days

Directions

Adults 18 years and older:

- only use one patch at a time
- clean and dry affected area
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- protect in product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polybutylacrylate, polybutylacrylate 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments? www.salonpas.us

Toll free 1-877-SALONPAS, Weekdays, 9AM to 5PM (PST) 1-7767-0

Salonpas[®] Arthritis Pain Patch **LARGE** 2 patches

Menthol 3%, Methyl Salicylate 10%

NDC#46581-685-02



for temporary relief of mild to moderate pain

- Arthritis
- Joint Pain
- Sprains
- Strains

Salonpas[®] Arthritis Pain Patch

APPLY FOR **8-12 HOURS**

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

2 LARGE patches
3 9/16" X 5 1/2" (10cm X 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS


Package not child resistant.
Keep out of reach of children.

LOT
EXP
Salonpas[®] Arthritis Pain Patch **LARGE 2 patches**
Menthol 3%, Methyl salicylate 10%

Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAGA TOSU
MADE IN JAPAN
Distributed by Hisamitsu America, Inc. Torrance, CA 90503
© : Registered Trade Mark
3 46581 68502 2

Salonpas[®] 2 patches

Arthritis Pain Patch **LARGE**
Menthol 3%, Methyl Salicylate 10%



press open
minty scent

Salonpas[®] Arthritis Pain Patch **LARGE** 2 patches

Menthol 3%, Methyl Salicylate 10%

Drug Facts

Active ingredients (in each patch)
Menthol 3%..... Topical analgesic
Methyl salicylate 10%(NSAID : nonsteroidal anti-inflammatory drug)..... Topical analgesic

Uses
Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
■ arthritis ■ sprains ■ strains ■ bruises ■ simple headache

Warnings
For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding.
The chance is small but higher if you:
■ are age 60 or older ■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed

Do not use
■ on the face or rashes ■ on wounds or damaged skin
■ with a heating pad ■ when sweating (such as from exercise or heat)
■ any patch from a pouch that has been open for 14 or more days ■ right before or after heart surgery

Ask a doctor before use if
■ you are allergic to topical products ■ the stomach bleeding warning applies to you
■ you are taking a diuretic ■ you have high blood pressure, heart disease, or kidney disease

When using this product
■ wash hands after applying or removing patch. Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
■ you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
■ rash, itching or skin irritation develops ■ condition worsens ■ symptoms last for more than 3 days
■ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
■ only use one patch at a time
■ clean and dry affected area
■ remove patch from backing film and apply to skin (see illustration)
■ apply one patch to the affected area and leave in place for up to 8 to 12 hours
■ if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
■ do not use more than 2 patches per day ■ do not use for more than 3 days in a row
■ the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

■ avoid storing product in direct sunlight
■ store at 20-25°C (68-77°F)
■ protect product from excessive moisture
■ some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?

Toll free 1-87-SALONPAS, Weekdays, 9AM to 5PM (PST) www.salonpas.us

CUT OPEN HERE ----- CUT OPEN HERE >>

FOLD TWICE AFTER OPENING
Discard unused patches 14 days after opening
FOLD TWICE AFTER OPENING



2 LARGE patches
3 1/8" x 5 1/2"
(10cm x 14cm)
FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

⊗ **DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS**

Hisamitsu®
Manufactured by Hisamitsu Pharmaceutical Co., Inc. JAPAN SAGA TOSU
Distributed by Hisamitsu America, Inc. Torrance, CA 90503
LOT
EXP.
Package not child resistant. Keep out of reach of children.
⊗ : Registered Trade Mark

CUT OPEN HERE ----- CUT OPEN HERE >>

Salonpas® Arthritis Pain Patch LARGE Menthol 3%, Methyl salicylate 10% minty scent

Drug Facts
Active ingredients (in each patch):
Menthol 3% Topical analgesic
Methyl salicylate 10% NSAID: nonsteroidal anti-inflammatory drug Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
arthritis sprains strains bruises simple backache

Warnings
For external use only
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
are age 60 or older have had stomach ulcers or bleeding problems
take a blood thinning (anticoagulant) or steroid drug
take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Do not use:
on the face or rashes on wounds or damaged skin
with a healing pad when sweating (such as from exercise or heat)
any patch from a pouch that has been open for 14 or more days
right before or after heart surgery
Ask a doctor before use if:
you are allergic to topical products the stomach bleeding warning applies to you
you are taking a diuretic you have high blood pressure, heart disease, or kidney disease

When using this product:
wash hands after applying or removing patch. Avoid contact with eyes.
If eye contact occurs, rinse thoroughly with water.
If the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if:
you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
rash, itching or skin irritation develops condition worsens symptoms last for more than 3 days
symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions
Adults 18 years and older:
clean and dry affected area
apply one patch to the affected area and leave in place for up to 8 to 12 hours
if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
do not use more than 2 patches per day do not use for more than 3 days in a row
the used patch should be removed from the skin when a new one is applied


Children under 18 years of age, do not use
Other information
store at 20-25°C (68-77°F) avoid storing product in direct sunlight
protect product from excessive moisture
some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients allicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-87-SALONPAS. Weekdays, 9AM to 5PM (PST) www.salonpas.us
1L775T-0

Salonpas®

Salonpas®



Salonpas®

Salonpas®

Salonpas®

Arthritis Pain Patch

APPLY FOR
8-12
HOURS



STRETCHABLE & FLEXIBLE

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

1 LARGE patch
3 1/8" X 5 1/2" (10cm X 14cm)

Salonpas®

Arthritis Pain Patch

1 LARGE patch X **12** sachets
3 1/8" X 5 1/2" (10cm X 14cm)

503 46587 68507 0

© 2011 Johnson & Johnson Consumer Inc. All rights reserved. For information on this product, visit us at www.Salonpas.com. Patch is a registered trademark of Johnson & Johnson Consumer Inc. Menthol, Methyl salicylate, and Topical Analgesic are trademarks of Johnson & Johnson Consumer Inc. FDA Approved Non-Prescription Pain Relieving Patch. Effectiveness Confirmed in Clinical Trial. Patch is a registered trademark of Johnson & Johnson Consumer Inc. Menthol, Methyl salicylate, and Topical Analgesic are trademarks of Johnson & Johnson Consumer Inc. FDA Approved Non-Prescription Pain Relieving Patch. Effectiveness Confirmed in Clinical Trial.

Hisamitsu® **LARGE** **minty scent**
NDC#45581-655-01

for temporary relief of mild to moderate pain

Arthritis
Joint Pain
Sprains
Strains



- 1 Peel the center film.
- 2 Place on affected area.
- 3 Slide the film upward.
- 4 Slide the other film downward.

APPLY FOR 8-12 HOURS

SALONIPAS®
Arthritis Pain Patch

**Menthol 3%
Methyl salicylate 10%
Topical Analgesic**

**FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL**

1 LARGE patch
3 1/8" x 5 1/4" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONIPAS PATCH AT A TIME. SEE DIRECTIONS

**Packaging not child resistant.
Keep out of reach of children.**

Manufactured by
Hisamitsu Pharmaceutical Co., Inc. Hisamitsu America, Inc.
10000 Wilshire Blvd., Suite 1000
Beverly Hills, CA 90210

Describe your pain
at www.salonipas.us

3 4658168501 5

>> CUT OPEN HERE ----- CUT OPEN HERE **>>**

Drug Facts

Active ingredients (in each patch)

Menthol 3%	Topical analgesic
Methyl salicylate 10% (NSAID) : nonsteroidal anti-inflammatory drug	Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:

- arthritis
- sprains
- strains
- bruises
- simple backache

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- on the face or rashes
- with a heating pad
- any patch from a pouch that has been open for 14 or more days
- right before or after heart surgery
- Ask a doctor before use if
 - you are allergic to topical products
 - the stomach bleeding warning applies to you
 - you are taking a diuretic
 - you have high blood pressure, heart disease, or kidney disease
 - you are taking a diuretic

When using this product

- wash hands after applying or removing patch. Avoid contact with eyes.
- if eye contact occurs, rinse thoroughly with water.
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
- Stop use and ask a doctor if
 - you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
 - stomach pain or upset gets worse or lasts
- rash, itching or skin irritation develops
- condition worsens
- symptoms last for more than 3 days
- symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:

- only use one patch at a time
- remove patch from backing film and apply to skin (see illustration)
- apply one patch to the affected area and leave in place for up to 8 to 12 hours
- if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- do not use more than 2 patches per day
- do not use for more than 3 days in a row
- the used patch should be removed from the skin when a new one is applied

Children under 18 years of age: do not use

Other information

- store at 20-25°C (68-77°F)
- avoid storing product in direct sunlight
- protect product from excessive moisture
- some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients

alicyclic saturated hydrocarbon resin, backing skin, film, mineral oil, polyisobutylene, polyisobutylene 1,200,000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-877-SALONIPAS, weekdays, 9AM to 5PM (PST)
www.salonipas.us

LOT
EXP.

© - Registered Trade Mark 1L7747-0

Hisamitsu
LARGE

FREE SAMPLE NOT FOR RE-SALE

minty scent

NDC+45681-085-99

for temporary relief of mild to moderate pain

Arthritis
Joint Pain
Sprains
Strains

- 1 Peel the center film.
- 2 Place on affected area.
- 3 Slide the film upward.
- 4 Slide the other film downward.

Salonpas

Arthritis Pain Patch

Menthol 3%
Methyl salicylate 10%
Topical Analgesic

APPLY FOR 8-12 HOURS

1 LARGE patch
3 1/4" x 5 1/2" (10cm x 14cm)

DO NOT USE MORE THAN ONE SALONPAS PATCH AT A TIME. SEE DIRECTIONS

Package not child resistant. Keep out of reach of children.

FDA APPROVED NON-PRESCRIPTION PAIN RELIEVING PATCH
EFFECTIVENESS CONFIRMED IN CLINICAL TRIAL

Manufactured by
Hisamitsu Pharmaceutical Co., Inc.
MADE IN JAPAN

On a shelf at
Hisamitsu America, Inc.
Pompano Beach, FL 33063

> CUT OPEN HERE ----- CUT OPEN HERE <

Drug Facts

Active ingredients (in each patch)

Menthol 3% Topical analgesic
Methyl salicylate 10% NSAID: nonsteroidal anti-inflammatory drug
Topical analgesic

Uses Temporarily relieves mild to moderate aches & pains of muscles & joints associated with:
arthritis sprains strains bruises simple backache

Warnings

For external use only

Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you:
are age 60 or older
have had stomach ulcers or bleeding problems
take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product
take more or for a longer time than directed

Do not use
on the face or rashes
with a heating pad
any pain from a pouch that has been open for 14 or more days
if allergic to aspirin or other NSAIDs

Ask a doctor before use if
you are allergic to topical products
you have high blood pressure, heart disease, or kidney disease
you are taking a diuretic
wash hands after applying or removing patch. Avoid contact with eyes.

When using this product
wash thoroughly with water.
the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
rash, itching or skin irritation develops
symptoms clear up and occur again within a few days
symptoms worsen
symptoms last for more than 3 days

If pregnant or breast-feeding
ask a doctor before use while breast-feeding and during the first 6 months of pregnancy. Do not use during the last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. If put in mouth, get medical help or contact a Poison Control Center right away. Package not child resistant. Dispose of the used patches after folding sticky ends together.

Directions

Adults 18 years and older:
only use one patch at a time
clean and dry affected area
remove patch from backing film and apply to skin (see illustration)
apply one patch to the affected area and leave in place for up to 8 to 12 hours
if pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
do not use more than 2 patches per day
do not use for more than 3 days in a row
the used patch should be removed from the skin when a new one is applied
children under 18 years of age: do not use

Other information
store at 20-25°C (68-77°F)
avoid storing product in direct sunlight
protect product from excessive moisture
some individuals may not experience pain relief until several hours after applying the patch

Inactive ingredients
alicyclic saturated hydrocarbon resin, backing cloth, film, mineral oil, polyisobutylene, polyisobutylene 1:200:1000, styrene-isoprene-styrene block copolymer, synthetic aluminum silicate

Questions or comments?
Toll free 1-877-SALONPAS Weekdays, 9AM to 5PM (PST)
www.salonpas.us

LOT
EXP

© : Registered Trade Mark 1L773T-0

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

JOEL SCHIFFENBAUER
11/05/2012