

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DORAL safely and effectively. See full prescribing information for DORAL.

DORAL (quazepam) Tablets for oral use C-IV
Initial U.S. Approval: 1985

RECENT MAJOR CHANGES

Dosage and Administration (2) 4/2013
Warnings and Precautions (5) 4/2013

INDICATIONS AND USAGE

DORAL, a gamma-aminobutyric (GABA_A) agonist, is indicated for the treatment of insomnia characterized by difficulty falling asleep, frequent nocturnal awakenings, and/or early morning awakenings. (1)

DOSAGE AND ADMINISTRATION

- Use the lowest dose effective for the patient
- Recommended initial dose is 7.5 mg (2)
- Split the 15 mg tablet along the score line to achieve 7.5 mg dose (2)
- The elderly and debilitated may be more sensitive to benzodiazepines (2)

DOSAGE FORMS AND STRENGTHS

- 15 mg functionally scored tablet, oral (3)

CONTRAINDICATIONS

- Hypersensitivity to quazepam or other benzodiazepines (4)
- Established or suspected sleep apnea, or chronic pulmonary insufficiency (4)

WARNINGS AND PRECAUTIONS

- CNS depressant effects: Impaired alertness and motor coordination, including risk of daytime impairment. Caution patients against driving and other activities requiring complete mental alertness (5.1)

- Benzodiazepine withdrawal syndrome: avoid abrupt discontinuation in at-risk patients (5.2)
- The failure of insomnia to remit after 7 to 10 days of treatment may indicate the presence of a primary psychiatric and/or medical illness that should be evaluated. (5.3)
- Severe anaphylactic/anaphylactoid reactions: Angioedema and anaphylaxis have been reported. Do not rechallenge if such reactions occur. (5.4)
- Sleep driving and other complex behaviors while not fully awake. Risk increases with dose and concomitant CNS depressants and alcohol. Immediately evaluate any new onset behavioral changes (5.5)
- Worsening of depression or suicidal thinking may occur: Prescribe the least number of tablets feasible to avoid intentional overdose (5.6)

ADVERSE REACTIONS

Most common adverse reactions (>1%): drowsiness, headache, fatigue, dizziness, dry mouth, dyspepsia (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Questcor Pharmaceuticals, Inc. at 1-800-465-9217 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- CNS Depressants: downward dose adjustment may be necessary due to additive effects (7)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm (8.1)
- Nursing Mothers: Administration of DORAL Tablets to nursing mothers is not recommended as quazepam and its metabolites are excreted in human milk. (8.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved Medication Guide.

Revised: 04/2013

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 CNS-Depressant Effects and Daytime Impairment
- 5.2 Benzodiazepine Withdrawal Syndrome
- 5.3 Need to Evaluate for Co-morbid Disorders
- 5.4 Severe Anaphylactic or Anaphylactoid Reactions
- 5.5 Abnormal Thinking and Behavior Changes
- 5.6 Worsening of Depression

6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use

8.5 Geriatric Use

9 DRUG ABUSE AND DEPENDENCE

- 9.1 Controlled Substance
- 9.2 Abuse and Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics
- 12.4 Drug Interactions

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

DORAL[®] (quazepam) is indicated for the treatment of insomnia characterized by difficulty in falling asleep, frequent nocturnal awakenings, and/or early morning awakenings. The effectiveness of DORAL has been established in placebo-controlled clinical studies of 5 nights duration in acute and chronic insomnia. The sustained effectiveness of DORAL has been established in chronic insomnia in a sleep lab (polysomnographic) study of 28 nights duration. Because insomnia is often transient and intermittent, the prolonged administration of DORAL Tablets is generally not necessary or recommended. Since insomnia may be a symptom of several other disorders, the possibility that the complaint may be related to a condition for which there is a more specific treatment should be considered.

2 DOSAGE AND ADMINISTRATION

Use the lowest dose effective for the patient, as important adverse effects of Doral are dose related.

The recommended initial dose is 7.5 mg. The 7.5 mg dose can be increased to 15 mg if necessary for efficacy.

The 7.5 mg dose can be achieved by splitting the 15 mg tablet along the score line.

2.2 Special populations

Elderly and debilitated patients may be more sensitive to benzodiazepines.

3 DOSAGE FORMS AND STRENGTHS

Tablets, 15 mg, functionally scored, capsule-shaped, light orange, slightly white speckled tablets, impressed with the product identification number 15 on one side of the tablet, and the product name (DORAL) on the other.

4 CONTRAINDICATIONS

DORAL is contraindicated in patients with known hypersensitivity to quazepam or other benzodiazepines. Rare cases of angioedema involving the tongue, glottis or larynx have been reported in patients after taking the first or subsequent doses of DORAL. Some patients have had additional symptoms such as dyspnea, throat closing, or nausea and vomiting that suggest anaphylaxis. Patients who develop such reactions should not be rechallenged with Doral.

Contraindicated in patients with established or suspected sleep apnea, or with pulmonary insufficiency.

5 WARNINGS AND PRECAUTIONS

5.1 CNS-depressant effects and Daytime impairment

Doral is a central nervous system (CNS) depressant and can impair daytime function in some patients even when used as prescribed. Prescribers should monitor for excess depressant effects, but impairment can occur in the absence of subjective symptoms, and may not be reliably detected by ordinary clinical exam (i.e. less than formal psychomotor testing). While pharmacodynamic tolerance or adaptation to some adverse depressant effects of Doral may develop, patients using Doral should be cautioned against driving or engaging in other hazardous activities or activities requiring complete mental alertness.

Additive effects occur with concomitant use of other CNS depressants (e.g., other benzodiazepines, opioids, tricyclic antidepressants, alcohol), including daytime use. Downward dose adjustment of Doral and concomitant CNS depressants should be considered. The potential for adverse drug interactions continues for several days following discontinuation of Doral, until serum levels of both active parent drug and psychoactive metabolites decline.

Use of Doral with other sedative-hypnotics is not recommended. Alcohol generally should not be used during treatment with Doral. The risk of next-day psychomotor impairment is increased if Doral is taken with less than a full night of sleep remaining (7- to 8 hours); if higher than the recommended dose is taken; if co-administered with other CNS depressants [*see Dosage and Administration (2.3)*].

5.2 Benzodiazepine Withdrawal Syndrome

A withdrawal syndrome similar to that from alcohol (e.g., convulsions, tremor, abdominal and muscle cramps, vomiting, and sweating) can occur following abrupt discontinuation of Doral. The more severe withdrawal effects are usually limited to patients taking higher than recommended doses over an extended time. Abrupt discontinuation should be avoided in such patients, and the dose gradually tapered. Prescribers should monitor patients for tolerance, abuse, and dependence.

Milder withdrawal symptoms (e.g., dysphoria and insomnia) can occur following abrupt discontinuation of benzodiazepines taken at therapeutic levels for short periods [*See Drug Abuse and Dependence (9)*].

5.3 Need to Evaluate for Co-morbid Diagnoses

Because sleep disturbances may be the presenting manifestation of a physical and/or psychiatric disorder, symptomatic treatment of insomnia should be initiated only after a careful evaluation of the patient. The failure of insomnia to remit after 7 to 10 days of treatment may indicate the presence of a primary psychiatric and/or medical illness that should be evaluated. Worsening of insomnia or the emergence of new thinking or behavior abnormalities may be the consequence of an unrecognized psychiatric or physical disorder. Such findings have emerged during the course of treatment with sedative-hypnotic drugs.

5.4 Severe Anaphylactic and Anaphylactoid Reactions

Rare cases of angioedema involving the tongue, glottis or larynx have been reported in patients after taking the first or subsequent doses of sedative-hypnotics, including DORAL. Some patients have had additional symptoms such as dyspnea, throat closing, or nausea and vomiting that suggest anaphylaxis.

Some patients have required medical therapy in the emergency department. If angioedema involves the tongue, glottis or larynx, airway obstruction may occur and be fatal. Patients who develop angioedema after treatment with DORAL should not be rechallenged with the drug.

5.5 Abnormal Thinking and Behavior Changes

Abnormal thinking and behavior changes have been reported in patients treated with sedative-hypnotics including Doral. Some of these changes include decreased inhibition (e.g., aggressiveness and extroversion that seemed out of character), bizarre behavior, and depersonalization. Visual and auditory hallucinations have also been reported. Amnesia, and other neuro-psychiatric symptoms may occur.

Paradoxical reactions such as stimulation, agitation, increased muscle spasticity, and sleep disturbances may occur unpredictably.

Complex behaviors such as "sleep-driving" (i.e., driving while not fully awake, with amnesia for the event) have been reported with use of sedative-hypnotics. These behaviors can occur with initial treatment or in patients previously tolerant of Doral or other sedative-hypnotics. Although these behaviors can occur with use at therapeutic doses, risk is increased by higher doses or concomitant use of alcohol or other CNS depressants. Due to risk to the patient and community, Doral should be discontinued if "sleep-driving" occurs.

Other complex behaviors (e.g., preparing and eating food, making phone calls, or having sex) have been reported in patients who are not fully awake after taking a sedative-hypnotic. As with sleep-driving, patients usually do not remember these events.

5.6 Worsening of Depression

Benzodiazepines may worsen depression. Consequently, appropriate precautions (e.g., limiting the total prescription size and increased monitoring for suicidal ideation) should be considered.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the label:

- CNS-depressant effects and next-day impairment [*see Warnings and Precautions (5.1)*]
- Benzodiazepine withdrawal syndrome [*see Warnings and Precautions (5.2)*]
- Abnormal thinking and behavior changes, and complex behaviors [*see Warnings and Precautions (5.5)*]
- Worsening of depression [*see Warnings and Precautions (5.6)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

The table shows adverse reactions occurring at an incidence of 1% or greater in relatively short-duration, placebo-controlled clinical trials of Doral. Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in actual practice.

| | DORAL 15 mg | PLACEBO |
|--------------------------|-------------------------|---------|
| NUMBER OF PATIENTS | 267 | 268 |
| | % OF PATIENTS REPORTING | |
| Central Nervous System | | |
| Daytime Drowsiness | 12 | 3 |
| Headache | 5 | 2 |
| Fatigue | 2 | 0 |
| Dizziness | 2 | <1 |
| Autonomic Nervous System | | |
| Dry Mouth | 2 | <1 |
| Gastrointestinal System | | |
| Dyspepsia | 1 | <1 |

A double-blind, controlled sleep laboratory study (N=30) in elderly patients compared the effects of quazepam 7.5 mg and 15 mg to that of placebo over a period of 7 days. Both the 7.5 mg and 15 mg doses appeared to be well tolerated. Caution must be used in interpreting this data due to the small size of the study.

7 DRUG INTERACTIONS

Benzodiazepines, including DORAL, produce additive CNS depressant effects when co-administered with ethanol or other CNS depressants (e.g. psychotropic medications, anticonvulsants, antihistamines). Downward dose adjustment of Doral and/or concomitant CNS depressants may be necessary because of additive effects.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

There are no adequate and well-controlled studies in pregnant women. Administration of benzodiazepines immediately prior to or during childbirth can result in a syndrome of hypothermia, hypotonia, respiratory depression, and difficulty feeding. In addition, infants born to mothers who have taken benzodiazepines during the later stages of pregnancy can

development dependence, and subsequently withdrawal, during the postnatal period. Although administration of quazepam to pregnant animals did not indicate a risk for adverse effects on morphological development at clinically relevant doses, data for other benzodiazepines suggest the possibility of adverse developmental effects (long-term effects on neurobehavioral and immunological function) in animals following prenatal exposure to benzodiazepines. DORAL should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Developmental toxicity studies of quazepam in mice at doses up to 400 times the human dose (15 mg) revealed no major drug-related malformations. Minor fetal skeletal variations that occurred were delayed ossification of the sternum, vertebrae, distal phalanges and supraoccipital bones, at doses approximately 70 and 400 times the human dose. A developmental toxicity study of quazepam in New Zealand rabbits at doses up to approximately 130 times the human dose demonstrated no effect on fetal morphology or development of offspring.

8.3 Nursing Mothers

Quazepam and its metabolites are excreted in human milk. Caution should be exercised when administering DORAL to a nursing woman.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

8.5 Geriatric Use

Doral may cause confusion and over-sedation in the elderly. Elderly patients generally should be started on a low dose of Doral and observed closely.

Elderly and debilitated patients may be more sensitive to benzodiazepines, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

A double-blind controlled sleep laboratory study (N=30) compared the effects of quazepam 7.5 mg and 15 mg to that of placebo over a period of 7 days. Both the 7.5 mg and 15 mg doses appeared to be well tolerated. Caution must be used in interpreting this data due to the small size of the study.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Quazepam is classified as a Schedule IV controlled substance by federal regulation.

9.2 Abuse and Dependence

Addiction-prone individuals (e.g. history of drug addiction or alcoholism) should be under careful surveillance when receiving Doral because of increased risk of abuse and dependence. Benzodiazepine withdrawal symptoms can occur following discontinuation of Doral [*see Warnings and Precautions (5.2)*].

Abuse and addiction are separate and distinct from physical dependence and tolerance. Abuse is characterized by misuse of the drug for non-medical purposes, often in combination with other psychoactive substances. Physical dependence is a state of adaptation that is manifested by a specific withdrawal syndrome that can be produced by abrupt cessation, rapid dose reduction, decreasing blood level of the drug and/or administration of an antagonist. Tolerance is a state of adaptation in which exposure to a drug induces changes that result in a diminution of one or more of the drug's effects over time. Tolerance may occur to both the desired and undesired effects of drugs and may develop at different rates for different effects.

Addiction is a primary, chronic, neurobiological disease with genetic, psychosocial, and environmental factors influencing its development and manifestations. It is characterized by behaviors that include one or more of the following: impaired control over drug use, compulsive use, continued use despite harm, and craving. Drug addiction is a treatable disease, utilizing a multidisciplinary approach, but relapse is common.


10 OVERDOSAGE

Contact a poison control center for up-to-date information on the management of benzodiazepine overdose.

Manifestations of Doral overdose include somnolence, confusion, and coma. General supportive measures should be employed, along with immediate gastric lavage. Dialysis is of limited value. Flumazenil may be useful, but can contribute to the appearance of neurological symptoms including convulsions. Hypotension may be treated by appropriate medical intervention. Animal experiments suggest that forced diuresis or hemodialysis are of little value in treating Doral overdose. As with the management of intentional overdose with any drug, the possibility of multiple drug ingestion should be considered.

11 DESCRIPTION

DORAL contains quazepam, a trifluoroethyl benzodiazepine hypnotic agent, having the chemical name 7-chloro-5-(o-fluoro-phenyl)-1,3-dihydro-1-(2,2,2-trifluoroethyl)-2H-1,4-benzodiazepine-2-thione and the following structural formula:



Quazepam has the empirical formula $C_{17}H_{11}ClF_4N_2S$, and a molecular weight of 386.8. It is a white crystalline compound, soluble in ethanol and insoluble in water. Each DORAL Tablet contains 15 mg of quazepam. The inactive ingredients for DORAL Tablets include cellulose, corn starch, FD&C Yellow No. 6 Al Lake, lactose, magnesium stearate, silicon dioxide, and sodium lauryl sulfate.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Quazepam, like other central nervous system agents of the 1,4-benzodiazepine class, presumably exerts its effects by binding to stereo-specific receptors at several sites within the central nervous system (CNS). The exact mechanism of action is unknown.

12.3 Pharmacokinetics

Absorption: Quazepam is rapidly (absorption half-life of about 30 minutes) and well absorbed from the gastrointestinal tract. The peak plasma concentration of quazepam is approximately 20 ng/mL after a 15 mg dose and occurs at about 2 hours.

Metabolism: Quazepam, the active parent compound, is extensively metabolized in the liver; two of the plasma metabolites are 2-oxoquazepam and N-desalkyl-2-oxoquazepam. All three compounds show CNS depressant activity.

Distribution: The degree of plasma protein binding for quazepam and its two major metabolites is greater than 95%.

Elimination: Following administration of ^{14}C -quazepam, 31% of the dose appeared in the urine and 23% in the feces over five days; only trace amounts of unchanged drug were present in the urine.

The mean elimination half-life of quazepam and 2-oxoquazepam is 39 hours and that of N-desalkyl-2-oxoquazepam is 73 hours. Steady-state levels of quazepam and 2-oxoquazepam are attained by the seventh daily dose and that of N-desalkyl-2-oxoquazepam by the thirteenth daily dose.

Special Populations:

Geriatrics: The pharmacokinetics of quazepam and 2-oxoquazepam in geriatric subjects are comparable to those seen in young adults; as with desalkyl metabolites of other benzodiazepines, the elimination half-life of N-desalkyl-2-oxoquazepam in geriatric patients is about twice that of young adults.

12.4 Drug Interactions

Bupropion (a CYP2B6 substrate): Co-administration of a single dose of 150 mg Bupropion Hydrochloride XL with steady state quazepam did not significantly affect the AUC and Cmax of bupropion or its primary metabolite, hydroxybupropion.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Quazepam showed no evidence of carcinogenicity in oral carcinogenicity studies in mice and hamsters.

Mutagenesis

Quazepam was negative in the bacterial reverse mutation (Ames) assay and equivocal in the mouse lymphoma *tk* assay.

Impairment of Fertility

Reproduction studies in mice conducted with quazepam at doses equal to 60 and 180 times the human dose of 15 mg produced slight reductions in fertility rate. Similar reductions in fertility rate have been reported in mice dosed with other benzodiazepines, and is believed to be related to the sedative effects of these drugs at high doses.

14 CLINICAL STUDIES

The effectiveness of DORAL was established in placebo-controlled clinical studies of 5 nights duration in acute and chronic insomnia. The sustained effectiveness of DORAL was established in chronic insomnia in a sleep laboratory (polysomnographic) study of 28 nights duration.

In the sleep laboratory study, DORAL significantly decreased sleep latency and total wake time, and significantly increased total sleep time and percent sleep time, for one or more nights. Doral 15 mg was effective on the first night of administration. Sleep latency, total wake time and wake time after sleep onset were still decreased and percent sleep time was still increased for several nights after the drug was discontinued. Percent slow wave sleep was decreased, and REM sleep was essentially unchanged. No transient sleep disturbance, such as “rebound insomnia,” was observed after withdrawal of the drug in sleep laboratory studies in 12 patients using 15 mg doses.

In outpatient studies, DORAL Tablets improved all subjective measures of sleep including sleep latency, duration of sleep, number of awakenings, occurrence of early morning awakening, and sleep quality. Some effects were evident on the first night of administration of DORAL (sleep latency, number of awakenings, and duration of sleep).

A double-blind, controlled sleep laboratory study (N=30) in elderly patients compared the effects of quazepam 7.5 mg and 15 mg to that of placebo over a period of 7 days. Both the 7.5 mg and 15 mg doses appeared to be effective. Caution must be used in interpreting this data due to the small size of the study.

16 HOW SUPPLIED

DORAL Tablets, 15 mg, functionally scored, capsule-shaped, light orange, slightly white speckled tablets, impressed with the product identification number 15 on one side of the tablet, and the product name (DORAL) on the other.

15 mg Bottles of 100 NDC 63004-7734-1

Store DORAL[®] Tablets at controlled room temperature 20°-25°C (68°-77°F).

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (Medication Guide).

Inform patients about the benefits and risks of DORAL, stressing the importance of use as directed. Assist patients in understanding the Medication Guide and instruct them to read it with each prescription refill.

CNS depressant Effects and Next-Day Impairment

Tell patients that Doral can cause next-day impairment, even in the absence of symptoms. Caution patients against driving or engaging in other hazardous activities or activities requiring complete mental alertness when using Doral. Tell patients that daytime impairment may persist for several days following discontinuation of Doral.

Withdrawal

Instruct patients to contact you before stopping or decreasing the dose of Doral, because withdrawal symptoms can occur.

Abnormal thinking and behavior change

Instruct patients that sedative hypnotics can cause abnormal thinking and behavior change, including “sleep-driving” and other complex behaviors while not being fully awake (preparing and eating food, making phone calls, or having sex). Tell patients to call you immediately if they develop any of these symptoms.

Severe Allergic Reactions

Inform patients that severe allergic reactions can occur from Doral. Describe the signs/symptoms of these reactions and advise patients to seek medical attention immediately if these occur.

Suicide

Tell patients that Doral can worsen depression, and to immediately report any suicidal thoughts.

Alcohol and other drugs

Ask patients about alcohol consumption, medicines they are taking now, and drugs they may be taking without a prescription. Advise patients that alcohol generally should not be used during treatment with Doral.

Pregnancy

Instruct patients to inform you if they are nursing or pregnant, or may become pregnant while taking Doral.

Tolerance, Abuse, and Dependence

Tell patients not to increase the dose of Doral on their own, and to inform you if they believe the drug “does not work”.



Manufactured for:
Questcor Pharmaceuticals, Inc.
Hayward, CA 94545 USA
phone (800) 411-3065
(510) 400-0700
fax (510) 400-0799

Manufactured by:
Meda Pharmaceuticals, Inc.
Somerset, NJ 08873-4120

Under license from Baker Norton Pharmaceuticals, Inc.

Printed in USA.
IN-1500-06 Rev. 04/13

MEDICATION GUIDE

DORAL (DOR-aI) (quazepam) Tablets (C-IV)

Read the Medication Guide that comes with DORAL before you start taking it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about DORAL?

Doral may cause serious side effects that you may not know are happening to you. These side effects include:

- sleepiness during the day
- not thinking clearly
- act strangely, confused, or upset
- “sleep-walking” or doing other activities when you are asleep like:
 - eating
 - talking
 - having sex
 - driving a car

Call your healthcare provider right away if you find out that you have done any of the above activities after taking DORAL.

What is DORAL?

DORAL is a prescription medicine used to treat certain types of insomnia including difficulty falling asleep, waking up often during the night, or waking up early in the morning.

It is not known if DORAL is safe and effective in children.

DORAL is a federally controlled substance (C-IV) because it can be abused or lead to dependence. Keep DORAL in a safe place to prevent misuse and abuse. Selling or giving away DORAL may harm others, and is against the law. Tell your healthcare provider if you have ever abused or been dependent on alcohol, prescription medicines or street drugs.

Who should not take DORAL?

Do not take DORAL if you:

- are allergic to quazepam or any of the ingredients in DORAL. See the end of this Medication Guide for a complete list of ingredients in DORAL.

- have had an allergic reaction to other sleep medicines or sedatives such as benzodiazepines. Symptoms of a serious allergic reaction to quazepam can include:
 - swelling of your face, lips, and throat that may cause difficulty breathing or swallowing
 - nausea and vomiting
 - have sleep apnea, snoring, breathing or lung problems

Talk to your healthcare provider before taking this medicine if you have any of these conditions.

What should I tell my healthcare provider before taking DORAL?

DORAL may not be right for you. Before taking DORAL, tell your healthcare provider about all of your health conditions, including if you:

- have a history of depression, mental illness or, suicidal thoughts
- have a history of drug or alcohol abuse or addiction
- have lung disease or breathing problems
- are pregnant or plan to become pregnant. It is not known if DORAL can harm your unborn baby.
- are breastfeeding, or plan to breastfeed. DORAL can pass through your breast milk and may harm your baby. Talk to your healthcare provider about the best way to feed your baby if you take DORAL.

Tell your healthcare provider about all of the medicines you take, including prescription and nonprescription medicines, vitamins and herbal supplements.

Medicines can interact with each other, sometimes causing serious side effects. **Do not take DORAL with other medicines that can make you sleepy unless your healthcare provider tells you to.**

Know the medicines you take. Keep a list of your medicines with you to show your healthcare provider and pharmacist each time you get a new medicine.

How should I take DORAL?

- See **“What is the most important information I should know about DORAL?”**
- Take DORAL exactly as your healthcare providers tell you to take it.
- Do not stop taking DORAL without talking to your healthcare provider, drug withdrawal symptoms can happen.
- DORAL comes in 15 mg tablets. Your healthcare provider may start your DORAL dose at 7.5 mg which is half a tablet. Talk to your healthcare provider or pharmacist about your dose schedule.

- You should not drink alcohol while you are taking DORAL.
- Call your healthcare provider if your insomnia worsens or is not better within 7 to 10 days. This may mean that there is another condition causing your sleep problem.
- If you take too much DORAL or overdose, get emergency treatment right away.

What are the possible side effects of DORAL?

DORAL may cause serious side effects, including:

- **getting out of bed while not being fully awake and doing an activity that you do not know you are doing.** See “**What is the most important information I should know about DORAL?**”
- **abnormal thoughts and behavior.** Symptoms include more outgoing or aggressive behavior than normal, confusion, agitation, hallucinations, worsening of depression, and suicidal thoughts or actions.
- **memory loss**
- **severe allergic reactions.** Symptoms include swelling of the tongue or throat, and trouble breathing. Get emergency medical help right away if you have these symptoms after taking DORAL.

Call your healthcare provider right away if you have any of the above side effects or any other side effects that worry you while using DORAL.

Common side effects of DORAL include:

- drowsiness
- headache
- feeling very tired
- dizziness
- dry mouth
- upset stomach

After you stop taking a sleep medicine, you may have symptoms for the next 1 to 2 days such as:

- trouble sleeping
- nausea
- flushing
- lightheadedness
- uncontrolled crying
- vomiting
- stomach cramps
- panic attack
- nervousness
- stomach area pain

Tell your healthcare provider if you have any side effect that bothers you or that

does not go away.

These are not all the possible side effects of DORAL. Ask your healthcare provider or pharmacist for more information.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store DORAL?

- Store at room temperature between 68°F to 77° F (20°C to 25°C).
- **Keep DORAL and all medicines out of the reach of children**

General information about the safe and effective use of Doral.

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use DORAL for a condition for which it was not prescribed. Do not share DORAL with other people, even if they have the same symptoms that you have. It may harm them and it is against the law.

This Medication Guide summarizes the most important information about DORAL. If you would like more information about DORAL, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about DORAL that is written for healthcare professionals.

If you would like more information, go to <http://www.DORALforsleep.com> or call Questcor Pharmaceuticals at 1-800-411-3065DORAL.

What are the ingredients in DORAL?

Active Ingredient: quazepam

Inactive Ingredients: cellulose, corn starch, FD&C Yellow No. 6 Al Lake, lactose, magnesium stearate, silicon dioxide, and sodium lauryl sulfate.

Distributed by Questcor Pharmaceuticals, Inc.

Hayward, CA 94545 USA

This Medication Guide has been approved by the U.S. Food and Drug Administration.

(IS-1500-02 Rev. 00/13 Month and year MG is approved)