

Nasacort Allergy 24HR 120 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_120_CTN_r3



Drug Facts

Active ingredient (in each spray) **Purpose**
 Triamcinolone acetonide
 (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE
 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

■ **do not use more than directed**
 ■ if you forget a dose, do **not** double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France

000000-00

NDC xxxxx-xxxx-x

Multi-Symptom Nasal Allergy Spray
Nasacort Allergy 24HR
 triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR
 Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.57 fl. oz.
 (16.9 ml)

120
 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive Ingredients

benzalkonium chloride, benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

call toll-free 1-800-633-1610

Lot XXXXX
 Exp MM/YYYY

Nasacort Allergy 24HR 120 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_120_PC_r3

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose</p> <p>Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p>Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients</p> <p>Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts</p> <p>When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one</p> <p>Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle</p> <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <p>■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age ■ do not use</p> <p>do not use more than directed ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use</p> <p>Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°- 25°C (68°- 77°F)</p>	
--	--	---	--

0044404-01

Dist. by: Chattem, Inc.,
a Sanofi Company
Chattanooga, TN 37409-0219
© 2013 Origin France

UPC

LOT EXP

Original Prescription Strength

NDC 41167-5800-5

Multi-Symptom Nasal Allergy Spray

Nasacort® Allergy 24HR

triamcinolone acetonide (glucocorticoid)
55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

120

SPRAYS

0.57 fl. oz.
(16.9 ml)

Nasacort Allergy 24HR 120 Sprays Partial Cover Clamshell - Inside Layout

10.09.13

NAS_Allergy_24HR_120_PC_r3



Nasacort Allergy 24HR 120 Sprays Partial Cover Clamshell 3-D Rendering

10.09.13

NAS_Allergy_24HR_120_PC_r3



Nasacort Allergy 24HR 120 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_120_PC_CPN_r3

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p>Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients</p> <p>Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts</p> <p>When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one</p> <p>Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle</p> <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <p>■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age ■ do not use</p> <p>■ do not use more than directed ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use</p> <p>Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°- 25°C (68°-77°F)</p>	
<p>Drug Facts (continued)</p> <p>Inactive ingredients benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water</p> <p>Questions or comments? call toll-free 1-800-633-1610</p>		<p>Nasacort Allergy 24HR</p> <p>Original Prescription Strength</p> <p>NDC 41167-5800-5</p> <p>Multi-Symptom Nasal Allergy Spray</p> <p>Nasacort Allergy 24HR</p> <p>triamcinolone acetonide (glucocorticoid) 55 mcg per spray/Nasal Allergy Symptom Reliever</p> <p>24HR Relief of:</p> <ul style="list-style-type: none"> ▶ Nasal Congestion ▶ Sneezing ▶ Runny Nose ▶ Itchy Nose <p>120 SPRAYS</p> <p>0.57 fl. oz. (16.9 ml)</p>	
<p>0044308-01</p> <p>Dist. by: Chatterm, Inc., a Sanofi Company Chattanooga, TN 37409-0219 © 2013 Origin France</p> <p>UPC</p>		<p>LOT EXP </p>	

Nasacort Allergy 24HR 120 Sprays Partial Cover Clamshell - Inside Layout with Coupon

10.09.13

NAS_Allergy_24HR_120_PC_CPN_r3

Prescription strength allergy relief.
Without a prescription.

Nasacort® Allergy 24HR

24 hour symptom relief
from indoor & outdoor allergies

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XX

SAVE \$X.XX Nasacort®
Allergy 24HR

On your next purchase of any size Nasacort® Allergy 24HR Spray

READ THE PRODUCT LABELING TO DETERMINE IF THIS PRODUCT IS APPROPRIATE FOR YOU TO USE. USE ONLY AS DIRECTED. COUPON CANNOT BE COMBINED WITH OTHER COUPONS. Consumer: Limit one coupon per transaction. Good only in the U.S. and overseas commissaries on the purchase of the specified product size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1c. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chattem, Inc., PO Box 880445, El Paso, TX 88588-0445. Use only as directed.

FPO


0 4 1167 00000 7

**BEFORE FIRST USE,
A NEW BOTTLE
MUST BE PRIMED**

**SHAKE WELL
BEFORE EACH USE**

Read insert (inside package) for complete instructions on how to:

- ▶ Get a new bottle ready (primed) before first use
- ▶ Prime bottle again if not used for more than 2 weeks
- ▶ Use the spray
- ▶ Clean the spray nozzle



Nasacort Allergy 24HR 120 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_120_CTN_CPN_r3



Drug Facts

Active ingredient (in each spray) Purpose
 Triamcinolone acetonide (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE
 ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France
 000000-00

Original Prescription Strength

Nasacort Allergy 24HR

Original Prescription Strength

NDC xxxxx-xxxx-x

Multi-Symptom Nasal Allergy Spray

Nasacort Allergy 24HR

triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR
 Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.57 fl. oz. (16.9 ml)

120 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive ingredients
 benzalkonium chloride, butylparaben, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water


Questions or comments?
 call toll-free 1-800-633-1610

Lot XXXXX
 Exp MM/YYYY

Nasacort Allergy 24HR 120 Sprays Carton - Inside with On-Pack Coupon

10.09.13


NAS_Allergy_24HR_120_CTN_CPN_r3



NAS ALLERGY 24HR 2X120 CLUB - Front

10.9.13

NAS_ALG_2X120_CLUB_r3



NAS ALLERGY 24HR 2X120 CLUB - Back

10.9.13

NAS_ALG_2X120_CLUB_r3

LOT XXXXXXX

EXP MM/YYYY



Nasacort® Allergy 24HR

Drug Facts

Active ingredient (in each spray)
Triamcinolone acetonide
(glucocorticoid) 55 mcg.....symptom reliever

Purpose
Nasal allergy

Uses

temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings

Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product

■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

■ **if pregnant or breast-feeding**, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

■ **do not use more than directed**
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Other information

■ **do not use if sealed package is torn or opened**
 ■ keep package and insert. They contain important information.
 ■ store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

UPC

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219 © 2013
 Origin France 0044314-01

Nasacort Allergy 24HR 30 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_30_CTN_CPN_r3



Drug Facts

Active ingredient (in each spray) Purpose
 Triamcinolone acetonide
 (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

PROFESSIONAL SAMPLE - NOT FOR RETAIL SALE

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France

000000-00

NDC xxxxx-xxxx-x

Original Prescription Strength

Nasacort Allergy 24HR

Multi-Symptom Nasal Allergy Spray

triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.23 fl. oz.
 (6.7 ml)

30
 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive Ingredients

benzalkonium chloride, butylparaben, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

call toll-free 1-800-633-1610

Lot XXXXX

Exp MMYYYY

Nasacort Allergy 24HR 30 Sprays Carton - Inside with On-Pack Coupon

10.09.13

NAS_Allergy_24HR_30_CTN_CPN_r3

**VALUABLE
COUPON INSIDE!**

MANUFACTURER'S COUPON | EXPIRATION DATE XXXXXX

SAVE \$X.XX Nasacort[®]
Allergy 24HR

On your next purchase of any size Nasacort[®] Allergy 24HR Spray.

READ THE PRODUCT LABELING TO DETERMINE IF THIS PRODUCT IS APPROPRIATE FOR YOU TO USE. USE ONLY AS DIRECTED. COUPON CANNOT BE COMBINED WITH OTHER COUPONS. Consumer: Limit one coupon per transaction. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chatham, Inc., PO Box 880445, El Paso, TX 88588-0445.

Use only as directed. © 2013 Chatham, Inc.

Nasacort Allergy 24HR 30 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_30_CTN_r3



Drug Facts

Active ingredient (in each spray) **Purpose**
 Triamcinolone acetonide
 (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

PROFESSIONAL SAMPLE - NOT FOR RETAIL SALE

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France

000000-00

Original Prescription Strength

Nasacort[®] Allergy 24HR

Original Prescription Strength

NDC xxxxx-xxxx-x

Multi-Symptom Nasal Allergy Spray

Nasacort[®] Allergy 24HR

triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.23 fl. oz.
 (6.7 ml)

30
 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort[®] Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive Ingredients

benzalkonium chloride, butylparaben, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

call toll-free 1-800-633-1610

Lot XXXXX

Exp MMYYYY

Nasacort Allergy 24HR 30 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_30_PC_r3

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose</p> <p>Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses</p> <p>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> nasal congestion runny nose sneezing itchy nose <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> in children under 2 years of age if you have ever had an allergic reaction to any of the ingredients <p>Ask a doctor before use if you</p> <ul style="list-style-type: none"> have had recent nose ulcers or nose surgery have had a nose injury that has not healed are using a steroid medicine for asthma, allergies or skin rash have an eye infection have or had glaucoma or cataracts <p>When using this product</p> <ul style="list-style-type: none"> the growth rate of some children may be slower some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. do not share this bottle with anyone else as this may spread germs remember to tell your doctor about all the medicines you take, including this one <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have, or come into contact with someone who has, chickenpox, measles or tuberculosis you have or develop symptoms of an infection such as a persistent fever you have any change in vision you have severe or frequent nosebleeds <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Directions</p> <p>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> get a new bottle ready (primed) before first use prime bottle again if not used for more than 2 weeks use the spray clean the spray nozzle <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older</p> <ul style="list-style-type: none"> once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <ul style="list-style-type: none"> the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year. <p>children 6 to under 12 years of age</p> <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. <p>children 2 to under 6 years of age</p> <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently <p>children under 2 years of age</p> <ul style="list-style-type: none"> do not use <p>do not use more than directed</p> <ul style="list-style-type: none"> if you forget a dose, do not double the next dose do not spray into eyes or mouth if allergy symptoms do not improve after one week, stop using and talk to a doctor do not use for the common cold shake well before each use <p>Other information</p> <ul style="list-style-type: none"> do not use if sealed package is torn or opened keep package and insert. They contain important information. store between 20°- 25°C (68°- 77°F) 	
--	--	--	--

PROFESSIONAL SAMPLE- NOT FOR RETAIL SALE

Dist. by: Chattem, Inc.,
a Sanofi Company
Chattanooga, TN 37409-0219
© 2013 Origin France

0044400-01

UPC

LOT

EXP

Nasacort[®]
Allergy 24HR

Drug Facts (continued)

Inactive ingredients
benzalkonium chloride,
carboxymethylcellulose sodium,
dextrose, edetate disodium,
hydrochloric acid or sodium
hydroxide (for pH adjustment),
microcrystalline cellulose,
polysorbate 80, purified water

Questions or comments?
call toll-free 1-800-633-1610

NDC 41167-5800-1

Original Prescription Strength

Multi-Symptom Nasal Allergy Spray

Nasacort[®]

Allergy 24HR

triamcinolone acetonide (glucocorticoid)
55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- › Nasal Congestion
- › Sneezing
- › Runny Nose
- › Itchy Nose

30

SPRAYS

0.23 fl. oz.
(6.7 ml)

Nasacort Allergy 24HR 30 Sprays Partial Cover Clamshell - Inside Layout

10.09.13

NAS_Allergy_24HR_30_PC_r3



Nasacort Allergy 24HR 30 Sprays Partial Cover Clamshell 3-D Rendering

10.09.13

NAS_Allergy_24HR_30_PC_r3



Nasacort Allergy 24HR 30 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_30_PC_CPN_r3

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p>Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients</p> <p>Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts</p> <p>When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one</p> <p>Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle</p> <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <p>■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age ■ do not use</p> <p>do not use more than directed ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use</p> <p>Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°-25°C (68°-77°F)</p>			
<p>PROFESSIONAL SAMPLE- NOT FOR RETAIL SALE</p> <p>Dist. by: Chattem, Inc., a Sanofi Company Chattanooga, TN 37409-0219 © 2013 Origin France</p> <p>0044304-01</p>		<p>Nasacort Allergy 24HR</p> <p>Drug Facts (continued)</p> <p>Inactive ingredients benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water</p> <p>Questions or comments? call toll-free 1-800-633-1610</p> <p>LOT EXP</p>		<p>Original Prescription Strength</p> <p>NDC 41167-5800-1</p> <p>Multi-Symptom Nasal Allergy Spray</p> <p>Nasacort Allergy 24HR</p> <p>triamcinolone acetonide (glucocorticoid) 55 mcg per spray/Nasal Allergy Symptom Reliever</p> <p>24 HR Relief of:</p> <ul style="list-style-type: none"> ▶ Nasal Congestion ▶ Sneezing ▶ Runny Nose ▶ Itchy Nose <p>0.23 fl. oz. (6.7 ml)</p> <p>30 SPRAYS</p> <p>UPC</p>	

Nasacort Allergy 24HR 30 Sprays Partial Cover Clamshell - Inside Layout with Coupon

10.09.13

NAS_Allergy_24HR_30_PC_CPN_r3

Prescription strength allergy relief.
Without a prescription.

Nasacort[®] Allergy 24HR

24 hour symptom relief
from indoor & outdoor allergies

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XX

SAVE \$X.XX Nasacort[®] Allergy 24HR

On your next purchase of any size Nasacort[®] Allergy 24HR Spray

READ THE PRODUCT LABELING TO DETERMINE IF THIS PRODUCT IS APPROPRIATE FOR YOU TO USE. USE ONLY AS DIRECTED. COUPON CANNOT BE COMBINED WITH OTHER COUPONS. Consumer: Limit one coupon per transaction. Good only in the U.S. and overseas commissaries on the purchase of the specified product size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chattem, Inc., PO Box 880445, El Paso, TX 88588-0445. Use only as directed.

FPO


0 4 1167 00000 7

Read insert (inside package) for complete instructions on how to:

- ▶ Get a new bottle ready (primed) before first use
- ▶ Prime bottle again if not used for more than 2 weeks
- ▶ Use the spray
- ▶ Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED


SHAKE WELL BEFORE EACH USE



NAS ALLERGY 24HR 3X120 CLUB - Front

10.9.13

NAS_ALG_3X120_CLUB_r3



NAS ALLERGY 24HR 3X120 CLUB - Back

10.9.13

NAS_ALG_3X120_CLUB_r3

LOT XXXXXXX

EXP MM/YYYY



Nasacort[®] Allergy 24HR

Drug Facts

Active ingredient (in each spray)
Triamcinolone acetonide
(glucocorticoid) 55 mcg.....symptom reliever

Purpose
Nasal allergy

Uses

temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings

Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you

■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product

■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if

■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

■ do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Other information

■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

UPC

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219 © 2013
 Origin France 0044314-01

NAS ALLERGY 24HR 4X120 CLUB - Front

10.9.13

NAS_ALG_4X120_CLUB_r3



NAS ALLERGY 24HR 4X120 CLUB - Back

10.9.13

NAS_ALG_4X120_CLUB_r3

LOT XXXXXXXX

EXP MM/YYYY



Nasacort® Allergy 24HR

Drug Facts

Active ingredient (in each spray)
Triamcinolone acetonide
(glucocorticoid) 55 mcg.....symptom reliever

Purpose
Nasal allergy

Uses
temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
■ in children under 2 years of age
■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
■ have had recent nose ulcers or nose surgery
■ have had a nose injury that has not healed
■ are using a steroid medicine for asthma, allergies or skin rash
■ have an eye infection
■ have or had glaucoma or cataracts

When using this product
■ the growth rate of some children may be slower
■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
■ do not share this bottle with anyone else as this may spread germs
■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
■ you have or develop symptoms of an infection such as a persistent fever
■ you have any change in vision
■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

Read insert (inside package) on how to:
■ get a new bottle ready (primed) before first use
■ prime bottle again if not used for more than 2 weeks
■ use the spray
■ clean the spray nozzle

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
■ once daily, spray 2 times into each nostril while sniffing gently
■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
■ an adult should supervise use
■ once daily, spray 1 time into each nostril while sniffing gently
■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
■ an adult should supervise use
■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
■ do not use

■ do not use more than directed
■ if you forget a dose, do not double the next dose
■ do not spray into eyes or mouth
■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
■ do not use for the common cold
■ shake well before each use

Other information

■ do not use if sealed package is torn or opened
■ keep package and insert. They contain important information.
■ store between 20°-25°C (68°-77°F)

Inactive ingredients

benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments? call toll-free 1-800-633-1610

UPC

Dist. by: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 © 2013
Origin France 0044314-01

Nasacort Allergy 24HR 60 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_60_CTN_r3



Drug Facts

Active ingredient (in each spray) Purpose
 Triamcinolone acetonide
 (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

■ do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France

000000-00

NDC xxxxx-xxxx-x

Multi-Symptom Nasal Allergy Spray
Nasacort Allergy 24HR
 triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR
 Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.37 fl. oz.
 (10.8 ml)

60
 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive Ingredients

benzalkonium chloride, benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

Lot XXXXX
 Exp MM/YYYY

Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_60_PC_r3

<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose</p> <p>Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses</p> <p>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> nasal congestion runny nose sneezing itchy nose <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> in children under 2 years of age if you have ever had an allergic reaction to any of the ingredients <p>Ask a doctor before use if you</p> <ul style="list-style-type: none"> have had recent nose ulcers or nose surgery have had a nose injury that has not healed are using a steroid medicine for asthma, allergies or skin rash have an eye infection have or had glaucoma or cataracts <p>When using this product</p> <ul style="list-style-type: none"> the growth rate of some children may be slower some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. do not share this bottle with anyone else as this may spread germs remember to tell your doctor about all the medicines you take, including this one <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you have, or come into contact with someone who has, chickenpox, measles or tuberculosis you have or develop symptoms of an infection such as a persistent fever you have any change in vision you have severe or frequent nosebleeds <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>		<p>Drug Facts (continued)</p> <p>Directions</p> <p>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> get a new bottle ready (primed) before first use prime bottle again if not used for more than 2 weeks use the spray clean the spray nozzle <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older</p> <ul style="list-style-type: none"> once daily, spray 2 times into each nostril while sniffing gently once your allergy symptoms improve, reduce to 1 spray in each nostril per day <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <ul style="list-style-type: none"> the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year. an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. <p>children 6 to under 12 years of age</p> <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently <p>children 2 to under 6 years of age</p> <ul style="list-style-type: none"> an adult should supervise use once daily, spray 1 time into each nostril while sniffing gently <p>children under 2 years of age</p> <ul style="list-style-type: none"> do not use <p>do not use more than directed</p> <ul style="list-style-type: none"> if you forget a dose, do not double the next dose do not spray into eyes or mouth if allergy symptoms do not improve after one week, stop using and talk to a doctor do not use for the common cold shake well before each use <p>Other information</p> <ul style="list-style-type: none"> do not use if sealed package is torn or opened keep package and insert. They contain important information. store between 20°-25°C (68°-77°F) 	
--	--	--	--

0044402-01

Dist. by: Chattem, Inc.,
a Sanofi Company
Chattanooga, TN 37409-0219
© 2013 Origin France

UPC

LOT

EXP

Original Prescription Strength

NDC 41167-5800-3

Multi-Symptom Nasal Allergy Spray

Nasacort® Allergy 24HR

triamcinolone acetonide (glucocorticoid)
55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- › Nasal Congestion
- › Sneezing
- › Runny Nose
- › Itchy Nose

60

SPRAYS

0.37 fl. oz. (10.8 ml)

Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Inside Layout

10.09.13

NAS_Allergy_24HR_60_PC_r3



Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell 3-D Rendering

10.09.13

NAS_Allergy_24HR_60_PC_r3



Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Outside Layout

10.09.13

NAS_Allergy_24HR_60_PC_CPN_r3

<p>Drug Facts</p>	<p>Drug Facts (continued)</p>	<p>Nasacort[®] Allergy 24HR</p>	<p>Original Prescription Strength</p> <p>NDC 41167-5800-3</p> <p>Multi-Symptom Nasal Allergy Spray</p> <p>Nasacort[®] Allergy 24HR</p> <p>triamcinolone acetonide (glucocorticoid) 55 mcg per spray/Nasal Allergy Symptom Reliever</p> <p>24 HR Relief of:</p> <ul style="list-style-type: none"> Nasal Congestion Sneezing Runny Nose Itchy Nose <p>0.37 fl. oz. (10.8 ml)</p> <p>60 SPRAYS</p>
<p>Active ingredient (in each spray) Purpose Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses temporarily relieves these symptoms of hay fever or other upper respiratory allergies: ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose</p> <p>Warnings Do not use ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients</p> <p>Ask a doctor before use if you ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts</p> <p>When using this product ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one</p> <p>Stop use and ask a doctor if ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Directions Read insert (inside package) on how to: ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle</p> <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER adults and children 12 years of age and older ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day</p> <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.</p> <p>children 6 to under 12 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.</p> <p>children 2 to under 6 years of age ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently</p> <p>children under 2 years of age ■ do not use</p> <p>■ do not use more than directed ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use</p> <p>Other information ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°- 25°C (68°-77°F)</p>	<p>Drug Facts (continued)</p> <p>Inactive ingredients benzalkonium chloride, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water</p> <p>Questions or comments? call toll-free 1-800-633-1610</p>	
<p>0044306-01</p> <p>Dist. by: Chattem, Inc., a Sanofi Company Chattanooga, TN 37409-0219 © 2013 Origin France</p>	<p>UPC</p>	<p>LOT EXP</p>	

Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Inside Layout with Coupon

10.09.13

NAS_Allergy_24HR_60_PC_CPN_r3

Prescription strength allergy relief.
Without a prescription.

Nasacort® Allergy 24HR

24 hour symptom relief
from indoor & outdoor allergies

MANUFACTURER'S COUPON EXPIRATION DATE XX/XX/XX

SAVE \$X.XX Nasacort® Allergy 24HR

On your next purchase of any size Nasacort® Allergy 24HR Spray

READ THE PRODUCT LABELING TO DETERMINE IF THIS PRODUCT IS APPROPRIATE FOR YOU TO USE. USE ONLY AS DIRECTED. COUPON CANNOT BE COMBINED WITH OTHER COUPONS. Consumer: Limit one coupon per transaction. Good only in the U.S. and overseas commissaries on the purchase of the specified product size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chattem, Inc., PO Box 880445, El Paso, TX 88588-0445. Use only as directed.

FPO


0 4 1167 00000 7

Read insert (inside package) for complete instructions on how to:

- ▶ Get a new bottle ready (primed) before first use
- ▶ Prime bottle again if not used for more than 2 weeks
- ▶ Use the spray
- ▶ Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

SHAKE WELL BEFORE EACH USE



Nasacort Allergy 24HR 60 Sprays Carton - Outside Layout

10.09.13

NAS_Allergy_24HR_60_CTN_CPN_r3



Drug Facts

Active ingredient (in each spray) Purpose
 Triamcinolone acetonide
 (glucocorticoid) 55 mcg.....Nasal allergy symptom reliever

Uses
 temporarily relieves these symptoms of hay fever or other upper respiratory allergies:
 ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose

Warnings
Do not use
 ■ in children under 2 years of age
 ■ if you have ever had an allergic reaction to any of the ingredients

Ask a doctor before use if you
 ■ have had recent nose ulcers or nose surgery
 ■ have had a nose injury that has not healed
 ■ are using a steroid medicine for asthma, allergies or skin rash
 ■ have an eye infection
 ■ have or had glaucoma or cataracts

When using this product
 ■ the growth rate of some children may be slower
 ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief.
 ■ do not share this bottle with anyone else as this may spread germs
 ■ remember to tell your doctor about all the medicines you take, including this one

Stop use and ask a doctor if
 ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis
 ■ you have or develop symptoms of an infection such as a persistent fever
 ■ you have any change in vision
 ■ you have severe or frequent nosebleeds

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 Read insert (inside package) on how to:
 ■ get a new bottle ready (primed) before first use
 ■ prime bottle again if not used for more than 2 weeks
 ■ use the spray
 ■ clean the spray nozzle

Drug Facts (continued)

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

adults and children 12 years of age and older
 ■ once daily, spray 2 times into each nostril while sniffing gently
 ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day

CHILDREN 2 TO UNDER 12 YEARS OF AGE

■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

children 6 to under 12 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently
 ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day.

children 2 to under 6 years of age
 ■ an adult should supervise use
 ■ once daily, spray 1 time into each nostril while sniffing gently

children under 2 years of age
 ■ do not use

■ do not use more than directed
 ■ if you forget a dose, do not double the next dose
 ■ do not spray into eyes or mouth
 ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor
 ■ do not use for the common cold
 ■ shake well before each use

Dist. by: Chatterm, Inc., a Sanofi Company
 Chattanooga, TN 37409-0219
 © 2013 Origin France

000000-00

NDC xxxxx-xxxx-x

Original Prescription Strength

Multi-Symptom Nasal Allergy Spray

Nasacort Allergy 24HR

triamcinolone acetonide (glucocorticoid)
 55 mcg per spray/Nasal Allergy Symptom Reliever

24 HR

Relief of:

- Nasal Congestion
- Sneezing
- Runny Nose
- Itchy Nose

0.37 fl. oz.
 (10.8 ml)

60
 SPRAYS

Read insert (inside package) for complete instructions on how to:

- Get a new bottle ready (primed) before first use
- Prime bottle again if not used for more than 2 weeks
- Use the spray
- Clean the spray nozzle

BEFORE FIRST USE, A NEW BOTTLE MUST BE PRIMED

Nasacort Allergy 24HR

SHAKE WELL BEFORE EACH USE

UPC

Actual Size

Drug Facts (continued)

Other Information
 ■ do not use if sealed package is torn or opened
 ■ keep package and insert. They contain important information.
 ■ store between 20° - 25° C (68° - 77° F)

Inactive Ingredients

benzalkonium chloride, butylparaben, carboxymethylcellulose sodium, dextrose, edetate disodium, hydrochloric acid or sodium hydroxide (for pH adjustment), microcrystalline cellulose, polysorbate 80, purified water

Questions or comments?

call toll-free 1-800-633-1610

Lot XXXXX

Exp MM/YYYY

Nasacort Allergy 24HR 60 Sprays Carton - Inside with On-Pack Coupon

10.09.13

NAS_Allergy_24HR_60_CTN_CPN_r3

**VALUABLE
COUPON INSIDE!**

MANUFACTURER'S COUPON | EXPIRATION DATE XXXXXX

SAVE \$X.XX Nasacort® Allergy 24HR

On your next purchase of any size Nasacort® Allergy 24HR Spray.

READ THE PRODUCT LABELING TO DETERMINE IF THIS PRODUCT IS APPROPRIATE FOR YOU TO USE. USE ONLY AS DIRECTED. COUPON CANNOT BE COMBINED WITH OTHER COUPONS. Consumer: Limit one coupon per transaction. Good only in the U.S. and overseas commissaries on the purchase of the specified product/size. Void where prohibited or if altered, copied, transferred, auctioned, or sold. You must pay sales tax. Cash value: 1/100 of 1¢. Retailer: We will reimburse you for the face value of this coupon plus 8¢ if redeemed in compliance with our Coupon Reimbursement Policy (available at address below). Invoices proving purchase of sufficient stock to cover coupons presented for redemption must be provided upon request. Failure to follow the policy may constitute fraud and may void all coupons submitted. Mail to: Chatterm, Inc., PO Box 880445, El Paso, TX 88588-0445.

Use only as directed. © 2013 Chatterm, Inc.

Nasacort Allergy 24HR 120 LBL


10.9.13

NAS_Allergy_24HR_120_LBL_r3


PAGE 1 (Front Page)




PAGE 2 (Inside Front Page)




PAGE 3



PAGE 4




PAGE 5




PAGE 6



PAGE 7




PAGE 8



BASE LABEL



OVERALL VIEW





SANOFI FISIONS LTD

Number of Pages
8
Right Hand Open
28 x 86 mm

 Final cutter guide

 Keyline

 Maximum text area

 Glue area - no print

Nasacort Allergy 24HR 30 LBL


10.9.13

NAS_Allergy_24HR_30_LBL_r3


PAGE 1 (Front Page)




PAGE 2 (Inside Front Page)




PAGE 3




PAGE 4




PAGE 5




PAGE 6




PAGE 7




PAGE 8



BASE LABEL



OVERALL VIEW



SANOFI FISONS LTD

Number of Pages
8
 Right Hand Open
28 x 86 mm

Final cutter guide

Keyline

Maximum text area

Glue area - no print

Nasacort Allergy 24HR 60 LBL

10.9.13

NAS_Allergy_24HR_60_LBL_r3


PAGE 1 (Front Page)




PAGE 2 (Inside Front Page)




PAGE 3



PAGE 4



PAGE 5




PAGE 6




PAGE 7




PAGE 8



BASE LABEL



OVERALL VIEW





SANOFI FISIONS LTD

Number of Pages
8
Right Hand Open
28 x 86 mm

 Final cutter guide

 Keyline

 Maximum text area

 Glue area - no print

Nasacort Allergy 24HR Bottle Cap Rendering

09.24.13

NAS_Allergy_24HR_BottleCap_Logo



Nasacort Allergy 24HR Bottle Cap Rendering

09.24.13

NAS_Allergy_24HR_BottleCap_Logo_A

12 point
Meta Plus
Bold

36 point
Universe Black

63 point
ITC Officina Sans
Bold

45 point
ITC Officina Sans
Bold



Nasacort Allergy 24HR Bottle Cap Rendering

09.24.13

NAS_Allergy_24HR_BottleCap_No Logo



Nasacort Allergy 24HR Instructions - Front

10.9.13

NAS_ALLERGY_24HR_INS_r3

Product Information
0081000-01

Nasacort[®] Allergy 24HR

Multi-Symptom Nasal Allergy Spray

triamcinolone acetonide (glucocorticoid)
55 mcg per spray / Nasal Allergy Symptom Reliever


IMPORTANT INFORMATION

- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.

- a. Remove cap (Figure A).
- b. Shake bottle.
- c. Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.




PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

1. Blow nose gently to clear nostrils.
2. Remove cap, then shake bottle.
3. Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
4. Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
5. Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).



DO NOT spray toward nasal septum (the wall between the 2 nostrils) (as shown in Figure D).



Correct Position



Wrong Position

6. While sniffing gently, spray into nostril

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

7. Repeat steps "4" through "6" for the other nostril.
8. After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

Nasacort Allergy 24HR Instructions - Back

10.9.13

NAS_ALLERGY_24HR_INS_r3

DIRECTIONS FOR USE

- **BEFORE first use, you must get a new bottle ready (primed).**
- **If bottle is not used for more than 2 weeks, prime bottle again.**

Dosing

ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER

- | | |
|--|---|
| adults and children 12 years of age and older | <ul style="list-style-type: none">• once daily, spray 2 times into each nostril while sniffing gently• once your allergy symptoms improve, reduce to 1 spray in each nostril per day |
|--|---|

CHILDREN 2 TO UNDER 12 YEARS OF AGE

- the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year.

- | | |
|--|--|
| children 6 to under 12 years of age | <ul style="list-style-type: none">• an adult should supervise use• once daily, spray 1 time into each nostril while sniffing gently• if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. |
|--|--|

- | | |
|---|--|
| children 2 to under 6 years of age | <ul style="list-style-type: none">• an adult should supervise use• once daily, spray 1 time into each nostril while sniffing gently |
|---|--|


- | | |
|--------------------------------------|---|
| children under 2 years of age | <ul style="list-style-type: none">• do not use |
|--------------------------------------|---|

Other important information for use

- Do not use more than directed.
- If you forget a dose, do **not** double the next dose.
- Do not spray into eyes or mouth.
- If you get the spray in your eyes, rinse well with water.
- If allergy symptoms do not improve after one week, stop using and talk to a doctor.
- Do not share this bottle with anyone else as this may spread germs.


IF PUMP DOES NOT SPRAY PROPERLY, THE NOZZLE MAY BE BLOCKED

1. Never try to unblock nozzle with a pin or any other object (as shown in Figure E).
2. Clean the nozzle as shown below.



CLEANING INSTRUCTIONS

1. Gently pull spray nozzle away from bottle (as shown in Figure F).
2. Rinse **SPRAY NOZZLE ONLY** under warm water (as shown in Figure G).
3. Shake or tap to remove excess water.
4. Re-attach spray nozzle to bottle.
5. Press and release spray nozzle until a fine spray is produced, taking care not to spray in face.



Nasacort Allergy 24HR pump is now ready to use.

Where can I get more information?

1-800-633-1610

Store between 20°-25°C (68°-77°F)

Distributed by: Chattem, Inc., a Sanofi Company
Chattanooga, TN 37409-0219 0081000-01 © 2013 Chattem, Inc.

SIDE 2

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

SANDRA L KWEDER
10/11/2013

CURTIS J ROSEBRAUGH
10/11/2013