

MEDICATION GUIDE

Morphine Sulfate (mor-phen) (CII) Oral Solution

IMPORTANT: Keep Morphine Sulfate Oral Solution in a safe place away from children. Accidental use by a child is a medical emergency and can cause death. If a child accidentally takes Morphine Sulfate Oral Solution, get emergency help right away.

Read this Medication Guide that comes with Morphine Sulfate Oral Solution before you start taking it and each time you get a new prescription. There may be new information. This Medication Guide does not take the place of talking with your healthcare provider about your medical condition or your treatment.

What is the most important information I should know about Morphine Sulfate Oral Solution?

Morphine Sulfate Oral Solution can cause serious side effects, including death.

- **Take Morphine Sulfate Oral Solution exactly as prescribed by your healthcare provider.** If you take the wrong dose or strength of Morphine Sulfate Oral Solution, you could overdose and die.
- It is especially important when you take Morphine Sulfate Oral Solution that you know exactly what dose and strength to take, and the right way to measure your medicine. Your healthcare provider or pharmacist should show you the right way to measure your medicine.
- Always use the oral syringe provided with Morphine Sulfate Oral Solution, 100 mg per 5 mL (20 mg per mL) to help make sure you measure the right amount.
- **Do not drink alcohol.** Using alcohol with Morphine Sulfate Oral Solution may increase your risk of dangerous side effects, including death.
- **Opioid Tolerance. Do not take Morphine Sulfate Oral Solution, 100 mg per 5 mL (20 mg per mL), unless you are “opioid tolerant.”** Opioid tolerant means that you take another opioid medicine for your constant (around the clock) pain and your body is used to it.

What is Morphine Sulfate Oral Solution?

Morphine Sulfate Oral Solution is in a group of drugs called narcotic pain medicine. Morphine Sulfate Oral Solution is only for adults who have moderate to severe pain.

- a prescription medicine that is used to manage moderate to severe pain that is expected to last a short period of time (acute), and pain that continues around-the-clock and is expected to last for a long period of time (chronic).
- Morphine Sulfate Oral Solution is a federally controlled substance (CII) because it is a strong opioid pain medicine that can be abused by people who abuse prescription medicines or street drugs.
- Prevent theft, misuse or abuse. Keep Morphine Sulfate Oral Solution in a safe place to keep it from being stolen. Morphine Sulfate Oral Solution can be a target for people who misuse or abuse prescription medicines or street drugs.
- Never give Morphine Sulfate Oral Solution to anyone else, even if they have the same symptoms you have. It may harm them or even cause death.

- Selling or giving away this medicine is against the law.

- It is not known if Morphine Sulfate Oral Solution is safe and effective in children under age 18 years of age.

- Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL), is only for people with moderate to severe pain who are already using an opioid narcotic pain medicine and have been using this medicine continuously for several weeks or longer.

Who should not take Morphine Sulfate Oral Solution?

Do not take Morphine Sulfate Oral Solution if you:

- are allergic to morphine, morphine salts or any of the ingredients in Morphine Sulfate Oral Solution. See [the end of this Medication guide for a complete list of ingredients in Morphine Sulfate Oral Solution.](#)
- are having an asthma attack or have severe asthma, trouble breathing, or lung problems
- have a bowel blockage called paralytic ileus

What should I tell my healthcare provider before taking Morphine Sulfate Oral Solution?

Before taking Morphine Sulfate Oral Solution, tell your healthcare provider if you:

- have trouble breathing or lung problems
- have had a head injury
- have liver or kidney problems
- have had adrenal gland problems, such as Addison’s disease
- have severe scoliosis that affects your breathing
- have thyroid problems
- have problems urinating or enlargement of your prostate
- have had convulsions or seizures
- have a past or present drinking problem or alcoholism
- have severe mental problems or hallucinations (seeing or hearing things that are not really there).
- have constipation or other bowel problems
- have problems with your pancreas or gallbladder
- have past or present substance abuse or drug addiction
- plan to have surgery
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if Morphine Sulfate Oral Solution will harm your unborn baby. Talk to your healthcare provider if you are pregnant or plan to become pregnant.

If you take Morphine Sulfate Oral Solution regularly before your baby is born, your newborn baby may have withdrawal symptoms because their body has become used to the medicine. Symptoms of withdrawal in a newborn baby may include:

- o irritability
- o vomiting
- o being very active
- o diarrhea or more stools than normal
- o problems sleeping
- o weight loss
- o high pitched cry
- o shaking (tremors)

If you take Morphine Sulfate Oral Solution right before your baby is born, your baby could have breathing problems.

- are breast-feeding or plan to breast-feed. Some Morphine Sulfate Oral Solution passes into your breast milk. A nursing baby could become very sleepy and have problems breathing or feeding well. If you stop breast-feeding, your baby may have withdrawal symptoms. See the list of withdrawal symptoms above. You and your healthcare provider should decide if you will take Morphine Sulfate Oral Solution or breast-feed.

Tell your healthcare provider about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Sometimes the doses of medicines that you take with Morphine Sulfate Oral Solution may need to be changed if used together. Morphine Sulfate Oral Solution and other medicines may affect each other causing serious side effects. Be especially careful about taking other medicines that make you sleepy such as:

- sleeping pills
- other pain medicines
- anti-nausea medicines
- tranquilizers
- muscle relaxants
- Do not take Morphine Sulfate Oral Solution if you already take a monoamine oxidase inhibitor medicine (MAOI) or within 14 days after you stop taking an MAOI medicine.
- Do not take any new medicine while using Morphine Sulfate Oral Solution until you have talked with your healthcare provider or pharmacist. They will tell you if it is safe to take other medicines with Morphine Sulfate.

Ask your healthcare provider if you are not sure if your medicine is one listed above.

Know the medicines you take. Keep a list of them and show it to your healthcare provider and pharmacist when you get a new medicine.

How should I take Morphine Sulfate Oral Solution?

- See “What is the most important information I should know about Morphine Sulfate Oral Solution?”
- **Take Morphine Sulfate Oral Solution exactly as prescribed.** Do not change your dose unless your healthcare provider tells you to. Your healthcare provider may change your dose after seeing how the medicine affects you. Call your healthcare provider if your pain is not well controlled with your prescribed dose of Morphine Sulfate Oral Solution.
- **You can take Morphine Sulfate Oral Solution with or without food.**
- **Make sure you understand exactly how to measure your dose.** Make sure you understand exactly how to measure your dose. Always use the oral syringe provided with your Morphine Sulfate Oral Solution, 100 mg per 5 mL (20 mg per mL) to help make sure you measure the right amount. See the [Patient Instructions for Use](#) at the end of this Medication Guide for information about how to measure your dose the right way. Ask your healthcare provider or pharmacist if you are not sure what dose of Morphine Sulfate Oral Solution you should take or if you are not sure how to use the oral syringe.
- **Do not stop taking Morphine Sulfate Oral Solution suddenly.** If you have been taking Morphine Sulfate Oral Solution for more than a few weeks, stopping Morphine Sulfate Oral Solution suddenly can make you sick with withdrawal symptoms (for example, nausea, vomiting, diarrhea, anxiety, and shivering). If your healthcare provider decides you no longer need Morphine Sulfate Oral

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Morphine Sulfate Oral Solution safely and effectively. See full prescribing information for Morphine Sulfate Oral Solution.

Morphine Sulfate Oral Solution, CII
Initial U.S. Approval: 1984

WARNING: RISK OF MEDICATION ERRORS

- **Morphine Sulfate Oral Solution is available in multiple concentrations. The 100 mg per 5 mL (20 mg per mL) concentration is indicated for use in opioid-tolerant patients only.**
- **Take care to avoid dosing errors due to confusion between different concentrations and between mg and mL, which could result in accidental overdose and death. (5.1)**
- **Keep Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) out of the reach of children.**

INDICATIONS AND USAGE

Morphine sulfate is an opioid analgesic indicated for the management of moderate to severe acute and chronic pain where an opioid analgesic is appropriate. (1)

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is indicated for the management of moderate to severe acute and chronic pain in opioid-tolerant patients. (1)

DOSEAGE AND ADMINISTRATION

- Morphine Sulfate Oral Solution: 10 to 20 mg (0.5 mL to 1.0 mL) every 4 hours as needed. (2.2)

DOSEAGE FORMS AND STRENGTHS

Oral Solution: Each mL contains 20 mg of morphine sulfate USP. (3)

CONTRAINDICATIONS

- Known hypersensitivity to morphine. (4)
- Respiratory depression in the absence of resuscitative equipment. (4)
- Acute or severe bronchial asthma or hypercarbia. (4)
- Paralytic ileus. (4)

WARNINGS AND PRECAUTIONS

- Risk of Medication Errors: Use caution when prescribing, dispensing, and administering to avoid dosing errors due to confusion between different concentrations and between mg and mL, which could result in accidental overdose and death. (5.1)
- Respiratory depression: Increased risk in elderly, debilitated patients, those suffering from conditions accompanied by hypoxia, hypercapnia, or upper airway obstruction. (5.2)
- Controlled substance: Morphine sulfate is a Schedule II controlled substance with an abuse liability similar to other opioids. (5.3)

Patients should be assessed for their clinical risks for opioid abuse or addiction prior to being prescribed opioids (5.3)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: RISK OF MEDICATION ERRORS

1 INDICATIONS AND USAGE

2 DOSEAGE AND ADMINISTRATION

- Individualization of Dosage
- Initiation of Therapy in Opioid-Naive Patients
- Conversion to Oral Morphine Sulfate
- Maintenance of Therapy
- Cessation of Therapy

3 DOSEAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- Risk of Medication Errors
- Respiratory Depression
- Misuse, Abuse and Diversion of Opioids
- Interactions with Alcohol and Drugs of Abuse
- Use in Head Injury and Increased Intracranial Pressure
- Hypotensive Effect
- Gastrointestinal Effects
- Use in Pancreatic/Biliary Tract Disease
- Special Risk Groups
- Driving and Operating Machinery

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

- CNS Depressants
- Mixed Agonist/Antagonist Opioid Analgesics
- Cimetidine or cimetidine hydrochloride
- Monoamine Oxidase Inhibitors (MAOIs)
- Anticholinergics
- Glycoprotein (P-gp) Inhibitors

FULL PRESCRIBING INFORMATION

WARNING: RISK OF MEDICATION ERRORS

- **Morphine Sulfate Oral Solution is marketed in multiple concentrations. The 100 mg per 5 mL (20 mg per mL) concentration is indicated for use in opioid-tolerant patients only.**
- **Take care when prescribing and administering Morphine Sulfate Oral Solution to avoid dosing errors due to confusion between different concentrations and between mg and mL, which could result in accidental overdose and death. Take care to ensure the proper dose is communicated and dispensed.**
- **Keep Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) out of the reach of children. In case of accidental ingestion, seek emergency medical help immediately.**

1 INDICATIONS AND USAGE

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is an opioid analgesic indicated for the management of moderate to severe acute and chronic pain in opioid-tolerant patients.

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) may cause fatal respiratory depression when administered to patients not previously exposed to opioids. Patients considered to be opioid tolerant are those who are taking at least 60 mg oral morphine sulfate per day, or at least 30 mg of oral oxycodone hydrochloride per day, or at least 12 mg hydromorphone hydrochloride per day, or an equianalgesic dose of another opioid, for a week or longer.

2 DOSEAGE AND ADMINISTRATION

Morphine Sulfate Oral Solution is available in multiple concentrations. Take care when prescribing and administering Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) to avoid dosing errors due to confusion between different concentrations and between mg and mL, which could result in accidental overdose and death. Take care to ensure the proper dose is communicated and dispensed.

When writing prescriptions, include both the total dose in mg and the total dose in volume.

Always use the enclosed calibrated oral syringe when administering Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) to ensure the dose is measured and administered accurately.

Selection of patients for treatment with morphine sulfate should be governed by the same principles that apply to the use of similar opioid analgesics. Individualize treatment in every case, using non-opioid analgesics, opioids on an as needed basis and/or combination products, and chronic opioid therapy in a progressive plan of pain management such as outlined by the World Health Organization, the Agency for Healthcare Research and Quality, and the American Pain Society.

2.1 Individualization of Dosage

As with any opioid drug product, adjust the dosing regimen for each patient individually, taking into account the patient’s prior analgesic treatment experience. In the selection of the initial dose of morphine sulfate, give attention to the following:

- the total daily dose, potency and specific characteristics of the opioid the patient has been taking previously;
- the reliability of the relative potency estimate used to calculate the equivalent morphine sulfate dose needed;
- the patient’s degree of opioid tolerance;
- the general condition and medical status of the patient;
- concurrent medications;
- the type and severity of the patient’s pain;
- risk factors for abuse, addiction or diversion, including a prior history of abuse, addiction or diversion.

The following dosing recommendations, therefore, can only be considered suggested approaches to what is actually a series of clinical decisions over time in the management of the pain of each individual patient.

Continual re-evaluation of the patient receiving morphine sulfate is important, with special attention to the maintenance of pain control and the relative incidence of side effects associated with therapy. During chronic therapy, especially for non-cancer-related pain, periodically re-assess the continued need for the use of opioid analgesics.

During periods of changing analgesic requirements, including initial titration, frequent contact is recommended between physician, other members of the healthcare team, the patient, and the caregiver/family.

2.2 Initiation of Therapy in Opioid-Naive Patients

Do not initiate treatment with Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) in patients who are opioid naive. Select an alternate product with lower concentration.

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is for use in **opioid-tolerant patients only** who have already been receiving opioid therapy. Use this strength only for patients that have already been titrated to a stable analgesic regimen using lower strengths of morphine sulfate and who can benefit from use of a smaller volume of oral solution.

2.3 Conversion to Oral Morphine Sulfate

There is inter-patient variability in the potency of opioid drugs and opioid formulations. Therefore, a conservative approach is advised when determining the total daily dose of morphine sulfate. It is better to underestimate a patient’s 24-hour oral morphine sulfate dose and make available rescue medication than to overestimate the 24-hour oral morphine sulfate dose and manage an adverse experience of overdose.

Consider the following general points regarding opioid conversions.

Conversion from Parenteral Morphine to Oral Morphine Sulfate

For conversion from parenteral to oral morphine sulfate, anywhere from 3 to 6 mg of oral morphine sulfate may be required to provide pain relief equivalent to 1 mg of parenteral morphine.

Conversion from Parenteral Oral Non-Morphine Opioids to Oral Morphine Sulfate

In converting patients from other opioids to morphine sulfate, close observation and adjustment of dosage based upon the patient’s response to morphine sulfate is imperative. Physicians and other healthcare professionals are advised to refer to published relative potency information, keeping in mind that conversion ratios are only approximate.

Conversion from Controlled-Release Oral Morphine to Oral Morphine Sulfate

For a given dose, the same total amount of morphine sulfate is available from Morphine Sulfate Oral Solution, Morphine Sulfate Tablets, and controlled-release and extended-release morphine sulfate capsules. The extended duration of release of morphine sulfate from controlled-release tablets or extended-release tablets results in reduced maximum and increased minimum plasma morphine sulfate concentrations than with shorter acting morphine sulfate products. Conversion from oral solution or immediate-release tablets to the same total daily dose of controlled-release tablets or extended-release tablets could lead to excessive sedation at peak serum levels. Therefore, dosage adjustment with close observation is necessary.

2.4 Maintenance of Therapy

Continual re-evaluation of the patient receiving morphine sulfate is important, with special attention to the maintenance of pain control and the relative incidence of side effects associated with therapy. If the level of pain increases, effort should be made to identify the source of increased pain, while adjusting the dose as described above to decrease the level of pain. During chronic therapy, especially for non-cancer-related pain (or pain associated with other terminal illnesses), periodically reassess the continued need for the use of opioid analgesics.

2.5 Cessation of Therapy

When the patient no longer requires therapy with morphine sulfate, gradually taper the dose to prevent signs and symptoms of withdrawal in the physically dependent patient.

3 DOSEAGE FORMS AND STRENGTHS

Each mL of clear/colorless Morphine Sulfate Oral Solution contains 20 mg of morphine sulfate. USP. The concentration of this solution is 100 mg per 5 mL (20 mg per mL).

4 CONTRAINDICATIONS

- Morphine sulfate is contraindicated in patients with known hypersensitivity to morphine, morphine salts, or any components of the product.
- Morphine sulfate is contraindicated in patients with respiratory depression in the absence of resuscitative equipment.
- Morphine sulfate is contraindicated in patients with acute or severe bronchial asthma or hypercarbia.
- Morphine sulfate is contraindicated in any patient who has or is suspected of having paralytic ileus.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Medication Errors

Morphine Sulfate Oral Solution is marketed in lower concentrations (e.g. 10 mg per 5 mL and 20 mg

- Tolerance may develop. (9.3)

- CNS effects: Additive CNS depressive effects when used in conjunction with alcohol, other opioids, or illicit drugs. (5.4)

- Elevation of intracranial pressure: May be markedly exaggerated in the presence of head injury, other intracranial lesions. (5.5)

- Hypotensive effect: Increased risk with compromised ability to maintain blood pressure. (5.6)

- Prolonged gastric obstruction: In patients with gastrointestinal obstruction, especially paralytic ileus. (5.7)

- Spincter of Oddi spasm and diminished biliary/pancreatic secretions. Increased risk with biliary tract disease. (5.8)

- Special Risk Groups: Use with caution and in reduced dosages in patients with severe renal or hepatic impairment, Addison’s disease, hypothyroidism, prostatic hypertrophy, or urethral stricture, elderly, CNS depression, toxic psychosis, acute alcoholism and delirium tremens, may aggravate or induce seizures. (5.9)

- Impaired mental/physical abilities: Caution must be used with potentially hazardous activities. (5.10)

ADVERSE REACTIONS

Most common adverse reactions are constipation, nausea, somnolence, lightheadedness, dizziness, vomiting, sweating. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Lannett Company, Inc. at 1-800-325-9994 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](#)

DRUG INTERACTIONS

- CNS depressants: Increased risk of respiratory depression, hypotension, profound sedation, or coma. Use with caution in reduced dosages. (7.1)

- Muscle relaxants: Enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increased degree of respiratory depression. (7.2)

- Mixed agonist/antagonist opioid analgesics (i.e. pentazocine, nalbuphine, and butorphanol): May reduce the analgesic effect and/or may precipitate withdrawal symptoms. (7.3)

- Cimetidine or cimetidine hydrochloride: Precipitates apnea, confusion and muscle twitching. (7.4)

- Monoamine oxidase inhibitors (MAOIs): Potentiate the action of morphine sulfate. Morphine sulfate should not be used in patients taking MAOIs or within 14 days of stopping such treatment. (7.5)

USE IN SPECIFIC POPULATIONS

- Pregnancy: Based on animal data, may cause fetal harm. (8.1)
- Geriatric patients (8.5), Hepatic impairment (8.7), Renal impairment (8.8): Use caution during dose selection, starting at the low end of the dosing range while carefully monitoring for side effects.

See 17 for PATIENT COUNSELING INFORMATION

Revised: 10/2012

8 USE IN SPECIFIC POPULATIONS

- Pregnancy
- Labor and Delivery
- Nursing Mothers
- Pediatric Use
- Geriatric Use
- Gender
- Hepatic Impairment
- Renal Impairment

9 DRUG ABUSE AND DEPENDENCE

- Controlled Substance
- Abuse
- Dependence

10 OVERDOSAGE

- Symptoms
- Treatment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- Mechanism of Action
- Pharmacokinetics
- Pharmacodynamics

13 NONCLINICAL TOXICOLOGY

- Carcinogenesis, Mutagenesis, Impairment of Fertility

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- Information for Patients and Caregivers
- MEDICATION GUIDE
- INSTRUCTIONS FOR USE

*Sections or subsections omitted from the full prescribing information are not listed

per 5 mL). Use caution when prescribing, dispensing, and administering any Morphine Sulfate Oral Solution to avoid dosing errors due to confusion between different concentrations and between mg and mL, which could result in accidental overdose and death. Use caution to ensure the dose is communicated clearly and dispensed accurately. Always use the enclosed calibrated oral syringe when administering Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) to ensure the dose is measured and administered accurately.

5.2 Respiratory Depression

Respiratory depression is the primary risk of morphine sulfate. Respiratory depression occurs more frequently in elderly or debilitated patients and in those suffering from conditions accompanied by hypoxia, hypercapnia, or upper airway obstruction, in whom even moderate therapeutic doses may significantly decrease pulmonary ventilation.

Use morphine sulfate with extreme caution in patients with chronic obstructive pulmonary disease or cor pulmonale and in patients having a substantially decreased respiratory reserve (e.g., severe kyphoscoliosis), hypoxia, hypercapnia, or pre-existing respiratory depression. In such patients, even usual therapeutic doses of morphine sulfate may increase airway resistance and decrease respiratory drive to the point of apnea. Consider alternative non-opioid analgesics, and use morphine sulfate only under careful medical supervision at the lowest effective dose in such patients.

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is for use in **opioid-tolerant patients only**. Administration of this formulation may cause fatal respiratory depression when administered to patients who are not tolerant to the respiratory depressant effects of opioids.

5.3 Misuse, Abuse and Diversion of Opioids

Morphine sulfate is an opioid agonist and a Schedule II controlled substance. Such drugs are sought by drug abusers and people with addiction disorders. Diversion of Schedule II products is an act subject to criminal penalty.

Morphine sulfate can be abused in a manner similar to other opioid agonists, legal or illicit. This should be considered when prescribing or dispensing morphine sulfate in situations where the physician or pharmacist is concerned about an increased risk of misuse, abuse, or diversion.

Morphine sulfate may be abused by crushing, chewing, snorting or injecting the product. These practices pose a significant risk to the abuser that could result in overdose and death. [See [DRUG ABUSE AND DEPENDENCE \(9\)](#)]

Concerns about abuse, addiction, and diversion should not prevent the proper management of pain. Healthcare professionals should contact their State Professional Licensing Board or State Controlled Substances Authority for information on how to prevent and detect abuse or diversion of this product.

5.4 Interactions with Alcohol and Drugs of Abuse

Morphine sulfate may be expected to have additive effects when used in conjunction with alcohol, other opioids, or illicit drugs that cause central nervous system depression because respiratory depression, hypotension, profound sedation, coma or death may result.

5.5 Use in Head Injury and Increased Intracranial Pressure

In the presence of head injury, intracranial lesions or a preexisting increase in intracranial pressure, the possible respiratory depressant effects of morphine sulfate and its potential to elevate cerebrospinal fluid pressure (resulting from vasodilation following CO₂ retention) may be markedly exaggerated. Furthermore, morphine

7.3 Mixed Agonist/Antagonist Opioid Analgesics

Do not administer mixed agonist/antagonist analgesics (i.e., pentazocine, nalbuphine, and butorphanol) to patients who have received or are receiving a course of therapy with a pure opioid agonist analgesic such as morphine sulfate. In these patients, mixed agonist/antagonist analgesics may reduce the analgesic effect and/or may precipitate withdrawal symptoms.

7.4 Cimetidine or cimetidine hydrochloride

Concomitant administration of morphine sulfate and cimetidine or cimetidine hydrochloride has been reported to precipitate apnea, confusion, and muscle twitching in an isolated report. Monitor patients for increased respiratory and CNS depression when receiving cimetidine or cimetidine hydrochloride concomitantly with morphine sulfate.

7.5 Monoamine Oxidase Inhibitors (MAOIs)

MAOIs markedly potentiate the action of morphine sulfate. Allow at least 14 days after stopping treatment with MAOIs before initiating treatment with morphine sulfate.

7.6 Anticholinergics

Anticholinergics or other medications with anticholinergic activity when used concurrently with opioid analgesics may result in increased risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.

7.7 P-Glycoprotein (P-gp) Inhibitors

Based on published reports, P-gp Inhibitors (e.g. quinidine) may increase the absorption/exposure of morphine sulfate by about two fold. Therefore, exercise caution when morphine sulfate is co-administered with P-gp inhibitors.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects (Pregnancy Category C)

Animal reproduction studies have not been conducted with morphine sulfate. It is also not known whether morphine sulfate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Only give morphine sulfate to a pregnant woman if clearly needed.

In humans, the frequency of congenital anomalies has been reported to be no greater than expected among the children of 70 women who were treated with morphine sulfate during the first four months of pregnancy or in 448 women treated with this drug anytime during pregnancy. Furthermore, no malformations were observed in the infant of a woman who attempted suicide by taking an overdose of morphine sulfate and other medication during the first trimester of pregnancy.

Several literature reports indicate that morphine sulfate administered subcutaneously during the early gestational period in mice and hamsters produced neurological, soft tissue and skeletal abnormalities. With one exception, the effects that have been reported were following doses that were maternally toxic and the abnormalities noted were characteristic to those observed when maternal toxicity is present. In one study, following subcutaneous infusion of doses greater than or equal to 0.15 mg/kg to mice, exencephaly, hydrocephalus, intestinal hemorrhage, split supracollicular, malformed sternbrae, and malformed xiphoid were noted in the absence of maternal toxicity. In the hamster, morphine sulfate given subcutaneously on gestation day 8 produced exencephaly and cranioschisis. Morphine sulfate was not a significant teratogen in the rat at exposure levels significantly beyond that normally encountered in clinical practice. In one study however, decreased litter size and viability were observed in the offspring of male rats administered morphine intraperitoneally (25 mg/kg) 1 day prior to mating. In two studies performed in the rabbit, no evidence of teratogenicity was reported at subcutaneous doses up to 100 mg/kg.

Nonteratogenic Effects

Controlled studies of chronic *in utero* morphine exposure in pregnant women have not been conducted. Infants born to mothers who have taken opioids chronically may exhibit withdrawal symptoms following birth, reversible reduction in brain volume, small size, decreased ventilatory response to CO₂ and increased risk of sudden infant death syndrome.

Manifestations of the neonatal withdrawal syndrome include irritability, hyperactivity, abnormal sleep pattern, high-pitched cry, tremor, vomiting, diarrhea, weight loss, and failure to gain weight. The time and amount of the mother's last dose and the rate of elimination of the drug from the newborn may affect the onset, duration, and severity of the disorder. When severe symptoms occur, pharmacologic intervention may be required.

Published literature has reported that exposure to morphine sulfate during pregnancy is associated with reduction in growth and a host of behavioral abnormalities in the offspring of animals. Morphine sulfate treatment during gestational periods of organogenesis in rats, hamsters, guinea pigs and rabbits resulted in the following treatment-related embryotoxicity and neonatal toxicity in one or more studies: decreased litter size, embryo-fetal viability, fetal and neonatal body weights, absolute brain and cerebellar weights, lengths or widths at birth and during the neonatal period, delayed motor and sexual maturation, and increased neonatal mortality, cyanosis and hypothermia. Decreased fertility in female offspring, and decreased plasma and testicular levels of luteinizing hormone and testosterone, decreased testes weights, seminiferous tubule shrinkage, germinal cell aplasia, and decreased spermatogenesis in male offspring were also observed. Behavioral abnormalities resulting from chronic morphine sulfate exposure of fetal animals included altered reflex and motor skill development, mild withdrawal, and altered responsiveness to morphine sulfate persisting into adulthood.

8.2 Labor and Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. Morphine sulfate is not recommended for use in women during and immediately prior to labor. Occasionally, opioid analgesics may prolong labor through actions which temporarily reduce the strength, duration and frequency of uterine contractions. However this effect is not consistent and may be offset by an increased rate of cervical dilatation, which tends to shorten labor. Closely observe neonates whose mothers received opioid analgesics during labor for signs of respiratory depression. Have a specific opioid antagonist, such as naloxone or nalmefene, available for reversal of opioid-induced respiratory depression in the neonate.

8.3 Nursing Mothers

Low levels of morphine have been detected in maternal milk. The milk: plasma morphine AUC ratio is about 2.5:1. The amount of morphine delivered to the infant depends on the plasma concentration of the mother, the amount of milk ingested by the infant, and the extent of first-pass metabolism. Because of the potential for serious adverse reactions in nursing infants from morphine sulfate including respiratory depression, sedation and possibly withdrawal symptoms, upon cessation of morphine sulfate administration to the mother, decide whether to discontinue nursing or to discontinue the drug, taking into account the importance of the drug to the mother.

8.4 Pediatric Use

The safety and effectiveness and the pharmacokinetics of Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) in pediatric patients below the age of 18 have not been established.

8.5 Geriatric Use

Elderly patients (aged 65 years or older) may have increased sensitivity to morphine sulfate. In general, use caution when selecting a dose for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

8.6 Gender

While evidence of greater post-operative morphine sulfate consumption in men compared to women is present in the literature, clinically significant differences in analgesic outcomes and pharmacokinetic parameters have not been consistently demonstrated. Some studies have shown an increased sensitivity to the adverse effects of morphine, including respiratory depression, in women compared to men.

8.7 Hepatic Impairment

Morphine sulfate pharmacokinetics have been reported to be significantly altered in patients with cirrhosis. Clearance was found to decrease with a corresponding increase in half-life. The morphine-3-glucuronide (M3G) and morphine-6-glucuronide (M6G) to morphine AUC ratios also decreased in these subjects, indicating diminished metabolic activity. Start these patients cautiously with lower doses of morphine sulfate and titrate slowly while carefully monitoring for side effects.

8.8 Renal Impairment

Morphine sulfate pharmacokinetics are altered in patients with renal failure. Clearance is decreased and the metabolites, M3G and M6G, may accumulate to much higher plasma levels in patients with renal failure as compared to patients with normal renal function. Start these patients cautiously with lower doses of morphine sulfate and titrate slowly while carefully monitoring for side effects.

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Morphine sulfate is a mu-agonist opioid and is a Schedule II controlled substance. Morphine sulfate, like other opioids used in analgesia, can be abused and is subject to criminal diversion.

9.2 Abuse

Drug addiction is characterized by compulsive use, use for non-medical purposes, and continued use despite harm or risk of harm. Drug addiction is a treatable disease, utilizing a multi-disciplinary approach, but relapse is common.

"Drug-seeking" behavior is very common in addicts and drug abusers. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing or referral, repeated "loss" of prescriptions, tampering with prescriptions and reluctance to provide prior medical records or contact information for other treating physician(s). "Doctor shopping" to obtain additional prescriptions is common among drug abusers and people suffering from untreated addiction.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Physicians should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence. The converse is also true. In addition, abuse of opioids can occur in the absence of true addiction and is characterized by misuse for non-medical purposes, often in combination with other psychoactive substances. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests is strongly advised.

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is intended for oral use only. Abuse of morphine sulfate poses a risk of overdose and death. The risk is increased with concurrent abuse of alcohol and other substances. Parenteral drug abuse is commonly associated with transmission of infectious diseases such as hepatitis and HIV.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal symptoms. *[See USE IN SPECIFIC POPULATIONS (8.2)]*

9.3 Dependence

Tolerance to opioids is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). The first sign of tolerance is usually a reduced duration of effect. Tolerance to different effects of opioids may develop to varying degrees and at varying rates in a given individual. Physical dependence is manifested by withdrawal symptoms after abrupt discontinuation of a drug or upon administration of an antagonist. Physical dependence and tolerance are frequent during chronic opioid therapy.

The opioid abstinence or withdrawal syndrome is characterized by some or all of the following: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, myalgia, and mydriasis. Other symptoms also may develop, including irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, or increased blood pressure, respiratory rate, or heart rate.

In general, taper opioids rather than abruptly discontinue. *[See DOSAGE AND ADMINISTRATION (2.5)]*

10 OVERDOSAGE

10.1 Symptoms

Acute overdosage with morphine sulfate is manifested by respiratory depression (a decrease in respiratory rate and/or tidal volume, Cheyne-Stokes respiration, cyanosis), extreme somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases, pulmonary edema, bradycardia, hypotension, cardiac arrest and death.

Morphine may cause miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomonic (e.g., pontine lesions of hemorrhagic or ischemic origin may produce similar findings). Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations. *[See CLINICAL PHARMACOLOGY (12)]*

10.2 Treatment

Give primary attention to re-establishment of a patent airway and institution of assisted or controlled ventilation. Employ supportive measures (including oxygen and vasopressors) in the management of circulatory shock and pulmonary edema accompanying overdose as indicated. Cardiac arrest or arrhythmias may require cardiac massage or defibrillation.

The pure opioid antagonists, naloxone or nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. Since the duration of reversal is expected to be less than the duration of action of morphine sulfate, carefully monitor the patient until spontaneous respiration is reliably re-established. If the response to opioid antagonists is sub-optimal or only brief in nature, administer additional antagonist as directed by the manufacturer of the product.


Do not administer opioid antagonists in the absence of clinically significant respiratory or circulatory depression secondary to morphine sulfate overdose.

Administer such agents cautiously to persons who are known, or suspected to be physically dependent on morphine sulfate. In such cases, an abrupt or complete reversal of opioid effects may precipitate an acute abstinence syndrome.

In an individual physically dependent on opioids, administration of the usual dose of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the antagonist administered. Reserve use of an opioid antagonist for cases where such treatment is clearly needed. If it is necessary to treat serious respiratory depression in the physically dependent patient, initiate administration of the antagonist with care and titrate with smaller than usual doses.

11 DESCRIPTION

Chemically, morphine sulfate is 7,8-didehydro-4,5 alpha-epoxy-17 methyl-morphinan-3,6 alpha-diol sulfate (2:1) (salt) pentahydrate with a molecular mass of 758. Morphine sulfate occurs as white, feathery, silky crystals; cubical masses of crystal; or white crystalline powder. It is soluble in water and slightly soluble in alcohol, but is practically insoluble in chloroform or ether. The octanol:water partition coefficient of morphine is 1.42 at physiologic pH and the pK_a is 7.9 for the tertiary nitrogen (the majority is ionized at pH 7.4).



Each mL of oral solution contains 20 mg of morphine sulfate, USP and the following inactive ingredients: anhydrous citric acid, edetate disodium dihydrate, glycerin, methylparaben, propylparaben, sodium benzoate, sorbitol and purified water.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Morphine, a pure opioid agonist, is relatively selective for the mu receptor, although it can interact with other opioid receptors at higher doses. In addition to analgesia, the widely diverse effects of morphine sulfate include drowsiness, changes in mood, respiratory depression, decreased gastrointestinal motility, nausea, vomiting, and alterations of the endocrine and autonomic nervous system.

Effects on the Central Nervous System (CNS)

The principal therapeutic action of morphine is analgesia. Other therapeutic effects of morphine include anxiolysis, euphoria and feelings of relaxation. Although the precise mechanism of the analgesic action is unknown, specific CNS opiate receptors and endogenous compounds with morphine-like activity have been identified throughout the brain and spinal cord and are likely to play a role in the expression and perception of analgesic effects. In common with other opioids, morphine causes respiratory depression, in part by a direct effect on the brainstem respiratory centers. Morphine and related opioids depress the cough reflex by direct effect on the cough center in the medulla.

Morphine causes miosis, even in total darkness.

Effects on the Gastrointestinal Tract and on Other Smooth Muscle

Gastric, biliary and pancreatic secretions are decreased by morphine. Morphine causes a reduction in motility and is associated with an increase in tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone is increased to the point of spasm. The end result may be constipation. Morphine can cause a marked increase in biliary tract pressure as a result of spasm of the sphincter of Oddi. Morphine may also cause spasm of the sphincter of the urinary bladder.

Effects on the Cardiovascular System

In therapeutic doses, morphine does not usually exert major effects on the cardiovascular system. Morphine produces peripheral vasodilation which may result in orthostatic hypotension and fainting. Release of histamine can occur, which may play a role in opioid-induced hypotension. Manifestations of histamine release and/or peripheral vasodilation may include pruritus, flushing, red eyes and sweating.

Endocrine System

Opioid agonists have been shown to have a variety of effects on the secretion of hormones. Opioids inhibit the secretion of ACTH, cortisol, and luteinizing hormone (LH) in humans. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon in humans and other species, rats and dogs. Thyroid stimulating hormone (TSH) has been shown to be both inhibited and stimulated by opioids.

Immune System

Opioids have been shown to have a variety of effects on components of the immune system in *in vitro* and animal models. The clinical significance of these findings is unknown.

12.2 Pharmacodynamics

Morphine concentrations are not predictive of analgesic response, especially in patients previously treated with opioids. The minimum effective concentration varies widely and is influenced by a variety of factors, including the extent of previous opioid use, age and general medical condition. Effective doses in tolerant patients may be significantly higher than in opioid-naïve patients.

12.3 Pharmacokinetics

Absorption

Morphine, when administered as morphine sulfate, is about two-thirds absorbed from the gastrointestinal tract with the maximum analgesic effect occurring 60 minutes post-administration. The oral bioavailability of morphine, when administered as morphine sulfate, is less than 40% and shows large inter-individual variability due to extensive pre-systemic metabolism.

Food Effects

Presence of food is not expected to significantly impact absorption of morphine from Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL). The concurrent intake of a high-fat, high-calorie meal enhanced the extent of absorption (about 16% increase in AUC), reduced the rate of absorption (about 25% decrease in C_{max}), and delayed the T_{max} (from 0.58 to 0.75 h) of morphine from the Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL).

Steady-State

Steady-state levels of morphine with Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) are expected to be achieved within 48 hours.

Distribution

Once absorbed, morphine is distributed to skeletal muscle, kidneys, liver, intestinal tract, lungs, spleen and brain. Although the primary site of action is the CNS, only small quantities cross the blood-brain barrier. Morphine also crosses the placental membranes and has been found in breast milk. The volume of distribution of morphine is approximately 1 to 6 L/kg, and morphine is 20 to 35% reversibly bound to plasma proteins.

Metabolism

The major pathway of morphine detoxification is conjugation, either with D-glucuronic acid to produce glucuronides or with sulfuric acid to produce morphine-3-ether sulfate. While a small fraction (less than 5%) of morphine is demethylated, virtually all morphine is converted by hepatic metabolism to the morphine-3-glucuronide (M3G) and morphine-6-glucuronide (M6G) metabolites (about 50% and 15%, respectively). M6G has been shown to have analgesic activity but crosses the blood-brain barrier poorly, while M3G has no significant analgesic activity.

Excretion

Most of a dose of morphine sulfate is excreted in urine as M3G and M6G, with elimination of morphine occurring primarily as renal excretion of M3G. Approximately 10% of the dose is excreted unchanged in urine. A small amount of glucuronide conjugates are excreted in bile, with minor enterohepatic recycling. Seven to 10% of administered morphine sulfate is excreted in the feces.

The mean adult plasma clearance is approximately 20 to 30 mL/min/kg. The effective terminal half-life of morphine after intravenous administration is reported to be approximately 2 hours. In some studies involving longer periods of plasma sampling, a longer terminal half-life of morphine of about 15 hours was reported.

Race

There may be some pharmacokinetic differences associated with race. In one published study, Chinese subjects given intravenous morphine sulfate had a higher clearance when compared to Caucasian subjects (1852 +/- 116 mL/min compared to 1495 +/- 80 mL/min).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies in animals to evaluate the carcinogenic potential of morphine sulfate have not been conducted. No formal studies to assess the mutagenic potential of morphine sulfate have been conducted. In the published literature, the results of *in vitro* studies showed that morphine sulfate is non-mutagenic in the *Drosophila melanogaster* lethal mutation assay and produced no evidence of chromosomal aberrations when incubated with murine splenocytes. Contrary to these results, morphine sulfate was found to increase DNA fragmentation when incubated *in vitro* with a human lymphoma line. *In vivo*, morphine sulfate has been reported to produce an increase in the frequency of micronuclei in bone marrow cells and immature red blood cells in the mouse micronucleus test and to induce chromosomal aberrations in murine lymphocytes and spermatids. Some of the *in vivo* clastogenic effects reported with morphine sulfate in mice may be directly related to increases in glucocorticoid levels produced by morphine sulfate in this species.

A literature report indicated that morphine sulfate impairs fertility in rats. In a fertility study in which male rats were administered morphine sulfate subcutaneous prior to mating (up to 30 mg/kg twice daily) and during mating (20 mg/kg twice daily) with untreated females, a number of adverse reproductive effects were observed. These included reduction in total pregnancies, higher incidence of pseudopregnancies, and reduction in implantation sites.

16 HOW SUPPLIED/STORAGE AND HANDLING

Morphine Sulfate Oral Solution

Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is a clear/colorless solution available as follows:

100 mg per 5 mL (20 mg per mL) Oral Solution

NDC 0527-1425-35: Bottles of 15 mL with an oral syringe

NDC 0527-1425-36: Bottles of 30 mL with an oral syringe

NDC 0527-1425-62: Bottles of 120 mL with an oral syringe

NDC 0527-1425-63: Bottles of 240 mL with an oral syringe

Storage

Store at Controlled Room Temperature, 15° to 30°C (59° to 86°F).

PROTECT FROM MOISTURE.

All opioids are liable to diversion and misuse both by the general public and healthcare workers and should be handled accordingly.

17 PATIENT COUNSELING INFORMATION

See Medication Guide

Provide the following information to patients receiving morphine sulfate or their caregivers:

- Advise patients that morphine sulfate is a narcotic pain medication, and should be taken only as directed.

- Advise patients that Morphine Sulfate Oral Solution is marketed in multiple concentrations including concentrations lower than 100 mg per 5 mL (20 mg per mL). Carefully instruct patients about which concentration they have been prescribed and how to measure and take the correct dose of Morphine Sulfate Oral Solution. Advise patients whenever the prescribed concentration is changed to avoid dosing errors, which could result in accidental overdose and death.

- Instruct patients to always use the enclosed calibrated oral syringe when administering Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) to ensure the dose is measured and administered accurately.

- Instruct patients that Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL) is only for patients who are already receiving opioid-therapy and have demonstrated opioid-tolerance. Use of this formulation may cause fatal respiratory depression when administered to patients who have not had previous exposure to opioids.

- Instruct patients NEVER to share their Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL), even if they have the same symptoms, because doing so could result in death.

- Instruct patients not to adjust the dose of morphine sulfate without consulting with a physician or other healthcare professional.

- Instruct patients that if they miss one dose of Morphine Sulfate Oral Solution 100 mg per 5 mL (20 mg per mL), they can take the dose when they remember it if they have pain, or they can wait for the next dose.

- Advise patients that morphine sulfate may cause drowsiness, dizziness, or lightheadedness and may impair mental and/or physical ability required for the performance of potentially hazardous tasks (e.g., driving, operating machinery). Advise patients started on morphine sulfate or patients whose dose has been adjusted to refrain from any potentially dangerous activity until it is established that they are not adversely affected.

- Advise patients that morphine sulfate will add to the effect of alcohol and other CNS depressants (such as antihistamines, sedatives, hypnotics, tranquilizers, general anesthetics, phenothiazines, other opioids, and monoamine oxidase [MAO] inhibitors).

- Instruct patients not to combine morphine sulfate with central nervous system depressants (sleep aids, tranquilizers) except by the orders of the prescribing physician, and not to combine with alcohol because dangerous additive effects may occur, resulting in serious injury or death.
- Instruct women of childbearing potential who become or are planning to become pregnant to consult a physician prior to initiating or continuing therapy with morphine sulfate.

- Advise patients that safe use in pregnancy has not been established and that prolonged use of opioid analgesics during pregnancy may cause fetal-neonatal physical dependence, and neonatal withdrawal may occur.

- If patients have been receiving treatment with morphine sulfate for more than a few weeks and cessation of therapy is indicated, counsel them on the importance of safely tapering the dose and that abruptly discontinuing the medication could precipitate withdrawal symptoms. Provide a dose schedule to accomplish a gradual discontinuation of the medication.

- Advise patients that morphine sulfate is a potential drug of abuse. They must protect it from theft. It should never be given to anyone other than the individual for whom it was prescribed.

- Instruct patients to keep morphine sulfate in a secure place out of the reach of children. When morphine sulfate is no longer needed, the unused oral solution should be destroyed by flushing down the toilet.

- Advise patients taking morphine sulfate of the potential for severe constipation; appropriate laxatives and/or stool softeners as well as other appropriate treatments should be initiated from the onset of opioid therapy.

- Advise patients of the most common adverse events that may occur while taking morphine sulfate: constipation, nausea, somnolence, lightheadedness, dizziness, sedation, vomiting, and sweating.

Manufactured by:
Cody Laboratories, Inc.
Cody, WY 82414

Manufactured for:
Lannett Company, Inc.
Philadelphia, PA 19136

Made in the USA

Revised 10/2012; Revision 3

11-138

Solution, ask how to slowly reduce this medicine. Do not stop taking Morphine Sulfate Oral Solution without talking to your healthcare provider.

- If you take too much Morphine Sulfate Oral Solution, call your healthcare provider or go to the nearest hospital emergency room right away.**

- If you miss one dose of Morphine Sulfate Oral Solution** you can take the dose when you remember it if you have pain, or you can wait for the next dose.

Talk with your healthcare provider regularly about your pain to see if you still need to take Morphine Sulfate Oral Solution.

What should I avoid while taking Morphine Sulfate Oral Solution?

- You should not drink alcohol while using Morphine Sulfate Oral Solution. Drinking alcohol with Morphine Sulfate Oral Solution may increase your risk of having dangerous side effects or death.**

- Do not drive, operate heavy machinery, or do other dangerous activities,** especially when you start taking Morphine Sulfate Oral Solution and when your dose is changed, until you know how Morphine Sulfate Oral Solution affects you. Morphine Sulfate Oral Solution can make you sleepy. Ask your healthcare provider to tell you when it is okay to do these activities.

What are the possible side effects of Morphine Sulfate Oral Solution?

See "What is the most important information I should know about Morphine Sulfate Oral Solution?"

Morphine Sulfate Oral Solution can cause serious breathing problems that can become life-threatening, especially if Morphine Sulfate Oral Solution is used the wrong way. Call your healthcare provider or get help right away if:

- your breathing slows down
- you have shallow breathing (little chest movement with breathing)
- you feel faint, dizzy, confused, or
- you have any other unusual symptoms