

12 of 5  
PP#



**Drug Facts (continued)**

**Warnings:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 4 doses in 24 hours, which is the maximum daily amount for ages 6 to 11 months. More than 5 doses in 24 hours, which is the maximum daily amount for ages 12 to 36 months. Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, hives, rash. If a skin reaction occurs, stop use and seek medical help. For rectal use only. Do not use if you are allergic to acetaminophen. With any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Ask a doctor before use if the child has liver disease. The child is taking the blood thinning drug warfarin.

**Drug Facts (continued)**

**Stop use and ask a doctor if:**

- fever lasts more than 3 days (72 hours), or recurs.
- pain lasts more than 3 days or gets worse.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition. **Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children. Even if you do not notice any signs or symptoms.

**Directions** - This product does not contain directions or warnings for adult use. Do not use more than directed. Remove wrapper carefully insert suppository well up into the rectum.

**Dosing Chart**

Age	Dose
under 6 months	Do not use unless directed by a doctor
6 to 11 months	Use 1 suppository every 6 hours (maximum of 4 doses in 24 hours)
12 to 36 months	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Drug Facts**

**Active ingredient (in each rectal suppository)** Acetaminophen, USP 80 mg

**Purposes** Pain reliever/fever reducer

**Uses** temporarily

- reduces fever
- relieves minor aches, pains, and headache

NDC 0472-1200-08

Infants'

Ages 6-36 months

80 mg



# FeverAll

Acetaminophen Suppositories  
Pain reliever/fever reducer

6 Rectal Suppositories

80 mg each



3

047211200061

Actavis  
Distributed by Actavis Mid Atlantic, LLC, Lincoln, NC 28022 USA. Made in USA.  
Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.

**Drug Facts (continued)**

**Other information** - store at 2°-27°C (36°-80°F) - do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polyorbate 80

**Questions?** 1-800-432-8334 between 9 am and 4 pm EST, Monday-Friday.

1200813M1 VC101916

**FeverAll**  
Acetaminophen  
Suppositories  
Pain reliever/fever reducer

(b) (4)



**FeverAll.**  
Acetaminophen Suppositories  
Pain reliever/fever reducer  
**80 mg each**



**Infants'** **FeverAll**  
Acetaminophen Suppositories  
Pain reliever/fever reducer



**50 Rectal Suppositories**

**80 mg each**

**NOT FOR RETAIL SALE**  
NDC 0472-1200-50



Dist. by Actavis Mid Atlantic LLC  
Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped.  
Do not use if imprinted wrapper is opened or damaged.**

12000813M1 VC101917



**Drug Facts** (continued)

**Do not use**

- if you are allergic to acetaminophen,
- with any other drug containing acetaminophen (prescription or nonprescription),
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- the child has liver disease,
- the child is taking the blood thinning drug warfarin,
- fever lasts more than 3 days (72 hours), or recurs,
- pain lasts more than 3 days or gets worse,
- new symptoms occur,
- redness or swelling is present in the painful area.

**Stop use and ask a doctor if**

- these symptoms occur,
- pain lasts more than 3 days or gets worse,
- new symptoms occur,
- redness or swelling is present in the painful area.

**Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Drug Facts** (continued)

**Directions**

- this product does not contain directions or warnings for adult use
- do not use more than directed
- remove wrapper
- carefully insert suppository well up into the rectum

**Dosing Chart**

Age	Dose
under 6 months	Do not use unless directed by a doctor
6 to 11 months	Use 1 suppository every 6 hours (maximum of 4 doses in 24 hours)
12 to 36 months	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Other Information**

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.



(b) (4)

12 of 5  
PP#



3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours).
under 3 years	Do not use unless directed by a doctor.
<b>Dose</b>	

**Dosing Chart**

**Drug Facts (continued)**

**Warnings:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** if you are allergic to acetaminophen, or if you are taking other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** the child has liver disease, or if the child is taking the blood thinning drug warfarin.

**Drug Facts (continued)**


**Warnings:** This product contains acetaminophen. Severe liver damage may occur if your child takes more than 5 doses in 24 hours, which is the maximum daily amount.

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** if you are allergic to acetaminophen, or if you are taking other drugs containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if** the child has liver disease, or if the child is taking the blood thinning drug warfarin.



**Drug Facts**

**Active ingredient (in each rectal suppository)** Acetaminophen, USP 120 mg

**Purposes** Pain reliever/fever reducer

**Uses** temporarily • reduces fever • relieves minor aches, pains, and headaches

NDC 0472-1201-08

**Children's**

**Ages 3-6 years**

**120 mg**



**FeverAll**

**Acetaminophen Suppositories**  
Pain reliever/fever reducer

**6 Rectal Suppositories**

**120 mg each**



Actavis  
Distributed by Actavis Mid Atlantic, LLC, Lincoln, NC 28092, USA. Made in USA.  
Temperature sensitive: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.

**Drug Facts (continued)**

**Other information** • store at 2°-27°C (35°-80°F) • do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

12010813M1 VC101919

**FeverAll**  
Acetaminophen  
Suppositories  
Pain reliever/fever reducer

(b) (4)



**FeverAll.**  
Acetaminophen Suppositories  
Pain reliever/fever reducer  
120 mg each



**Children's**

**Ages 3-6 years**

**FeverAll**

**Acetaminophen Suppositories**

Pain reliever/fever reducer

**50 Rectal Suppositories**

**120 mg each**

**NOT FOR RETAIL SALE**  
NDC 0472-1201-50



Dist. by Actavis Mid Atlantic LLC  
Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped.  
Do not use if imprinted wrapper is opened or damaged.**

12010813M1 VC101920



3 04721 20150 1

**Drug Facts**  
**Active ingredient (in each rectal suppository)** Acetaminophen, USP 120 mg  
**Purposes** Pain reliever/fever reducer

**Uses** temporarily • reduces fever • relieves minor aches, pains, and headache

**Warnings** Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 5 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.  
For rectal use only.

**Drug Facts (continued)**

**Do not use**

- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription).
- if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- the child has liver disease.
- the child is taking the blood thinning drug warfarin.

**Ask a doctor before use if**

- fever lasts more than 3 days (72 hours), or recurs.
- pain lasts more than 5 days or gets worse.
- new symptoms occur.
- redness or swelling is present in the painful area.

**Stop use and ask a doctor if**

- these may be signs of a serious condition.

**Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- this product does not contain directions or warnings for adult use
- do not use more than directed
- remove wrapper
- carefully insert suppository well up into the rectum

**Dosing Chart**


Age	Dose
under 3 years	Do not use unless directed by a doctor
3 to 6 years	Use 1 suppository every 4 to 6 hours (maximum of 5 doses in 24 hours)

**Other information**

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.



(b) (4)

12 of 5  
PP#



**Drug Facts (continued)**

**Warnings:** This product contains acetaminophen. Severe liver damage may occur if:  
 • a child 6 to 12 years takes more than 5 doses in 24 hours  
 • an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount  
 • taken with other drugs containing acetaminophen  
 • an adult takes 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 • skin reddening • blisters • rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
 • in children under 6 years  
 • if you are allergic to acetaminophen.  
 • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**  
 • you have liver disease.  
 • you are taking the blood thinning drug warfarin.

**Drug Facts**

**Active ingredient (in each rectal suppository)**  
 Acetaminophen, USP 325 mg

**Purposes**  
 Pain reliever/fever reducer

**Uses**  
 temporarily • reduces fever • relieves minor aches, pains, and headache

**Drug Facts (continued)**

**Other information**  
 • store at 2°-27°C (35°-80°F)  
 • do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients**  
 glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-534 between 9 am and 4 pm EST, Monday-Friday.

1202081301 VC101921

**325 mg**

**Jr. Strength**

**Ages 6-12 years**

**Doctor Recommended**

**FeverAll®**

**Acetaminophen Suppositories**

**Pain reliever/fever reducer**

**6 Rectal Suppositories**

**325 mg each**



M 04721120206 5

Actavis  
 Div. by Actavis Midco, LLC, Lincoln, NC 28020 USA. Made in USA.  
 Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.

**Drug Facts (continued)**

**Other information**  
 • store at 2°-27°C (35°-80°F)  
 • do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients**  
 glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-534 between 9 am and 4 pm EST, Monday-Friday.

1202081301 VC101921

**FeverAll®**

**Acetaminophen Suppositories**

**Pain reliever/fever reducer**

(b) (4)



**FeverAll.**  
Acetaminophen Suppositories  
Pain reliever/fever reducer  
325 mg each



**Jr. Strength**

**Ages 6-12 years**

**FeverAll**

**Acetaminophen Suppositories**

Pain reliever/fever reducer



**50 Rectal Suppositories**

**325 mg each**

**NOT FOR RETAIL SALE**  
NDC 0472-1202-50



Dist. by Actavis Mid Atlantic LLC  
Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped.  
Do not use if imprinted wrapper is opened or damaged.**

12020813Q1 VC101922



**Drug Facts** (continued)

**Do not use**

- in children under 6 years
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease.
- you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.
- These may be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- carefully insert suppository well up into the rectum
- remove wrapper

**Drug Facts** (continued)

**Other information**

- store at 2°-27°C (35°-80°F)
- do not use if imprinted suppository wrapper is opened or damaged

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday-Friday.

**Drug Facts** (continued)

**Active ingredient (in each rectal suppository)** Acetaminophen, USP 325 mg

**Purposes** Pain reliever/fever reducer

**Uses** temporarily • reduces fever • relieves minor aches, pains, and headache

**Warnings** Liver warning: This product contains acetaminophen. Severe liver damage may occur if a child 6 to 12 years takes more than 5 doses in 24 hours • an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • an adult takes 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**For rectal use only.**

(b) (4)



**Drug Facts (continued)**

**Do not use**

- in children under 12 years.
- if you are allergic to acetaminophen.
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease.
- you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- fever lasts more than 3 days (72 hours), or recurs.
- pain gets worse or lasts more than 10 days.
- new symptoms occur.
- redness or swelling is present in the painful area.

These may be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** If swallowed or in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical in case of overdose for adults and for children even if you do not notice any signs or symptoms.

**Questions?** 1-800-432-8534 between 9 am and 4 pm EST, Monday–Friday.

**Inactive ingredients** glycerol monostearate, hydrogenated vegetable oil, polyoxyethylene stearate, polysorbate 80

**Drug Facts**

**Active ingredient (in each rectal suppository)** Acetaminophen, USP 650 mg

**Purposes** Pain reliever/fever reducer

**Uses** temporarily • reduces fever • relieves minor aches, pains, and headache

**Warnings** This product contains acetaminophen. Severe liver damage may occur if • an adult or child 12 years and older takes more than 6 doses in 24 hours, which is the maximum daily amount • taken with other drugs containing acetaminophen • an adult takes 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**For rectal use only.**

**Adults'** **Ages 12 years and older**

**NOT FOR RETAIL SALE**  
NDC 0472-1203-50

**FeverAll®**  
Acetaminophen Suppositories  
Pain reliever/fever reducer

**650 mg each**

**50 Rectal Suppositories**

**Doctor Recommended**



**FeverAll®**  
Acetaminophen Suppositories  
Pain reliever/fever reducer  
**650 mg each**

**actavis**  
Dist. by Actavis Mid Atlantic LLC  
Lincolnton, NC 28092 USA Made in USA

**Tamper-evident: Suppositories are individually wrapped. Do not use if imprinted wrapper is opened or damaged.**

12030813J1 VC101923



3 04721 20350 5

