

(b) (4)

FRONT LABEL

(b) (4)



(b) (4)



TAMPER EVIDENT Do not use if the neckband printed on the right is broken or missing.

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema
 ■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

Inactive ingredients citric acid anhydrous, D&C Red #33, edetate disodium, ethylcellulose, FD&C Blue #1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-888-963-3382
 You may also report side effects to this phone number.

Reckitt Benckiser

C I A 71978J
 PT# C I A 60333
 8137782

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224 © 2014 RB 022014

Code 128
71978J

12 Hour Cough Relief
Children's Delsym
 Dextromethorphan Polistirex
 Extended-Release Suspension (Cough Suppressant)

COUGH

*Also Available
 In Orange Flavor*



Please visit our web site:
www.delsym.com

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

12 Hour Cough Relief
Children's Delsym
COUGH
 Grape Flavored Liquid

12 Hour Cough Relief
Children's Delsym
 Dextromethorphan Polistirex
 Extended-Release Suspension (Cough Suppressant)

COUGH

DELSYM® DOSING
 SHAKE WELL BEFORE USE.
 Measure only with dosing cup provided. Do not use dosing cup with other products. mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.



Dosing Cup Included

Contains No Fever Reducer or Pain Reliever

Alcohol-free
 148 mL (5 fl oz) Day or Night
 Grape Flavored Liquid

3-63824-27265-8
 (UPC @ 100% truncated)

Lot No.:
 Exp. Date:

UPC area is 1/16" below score and is 1-7/32"w x 11/16"h

(b) (4)

FRONT LABEL

(b) (4)

(b) (4)



(b) (4)



TAMPER EVIDENT Do not use if the neckband printed on the cap is broken or missing.

Drug Facts

Active ingredient (in each 5 mL) **Purpose**
 Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant

Uses temporarily relieves
 ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 ■ the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
 ■ chronic cough that lasts as occurs with smoking, asthma or emphysema
 ■ cough that occurs with too much phlegm (mucus)
Stop use and ask a doctor if cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use
 ■ measure only with dosing cup provided
 ■ do not use dosing cup with other products
 ■ dose as follows or as directed by a doctor
 ■ mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information ■ each 5 mL contains: sodium 7 mg
 ■ store at 20-25°C (68-77°F) ■ dosing cup provided

Inactive ingredients citric acid, edetate disodium, ethylcellulose, FD&C Yellow No. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-888-963-3382
 You may also report side effects to this phone number.

Reckitt Benckiser

C I A 70236M
 PT# C I A 60333
 8137779

Distributed by: Reckitt Benckiser, Parsippany, NJ 07054-0224 © 2014 RB 022114

Code 128
70236M

12 Hour Cough Relief

Children's Delsym

Dextromethorphan Polistirex
Extended-Release Suspension (Cough Suppressant)

12 Hour Cough Relief

Children's Delsym

COUGH

Orange Flavored Liquid

NDC 63824-176-65

12 Hour Cough Relief

Children's Delsym

Dextromethorphan Polistirex
Extended-Release Suspension (Cough Suppressant)

Also Available in Grape Flavor

COUGH

Dextromethorphan Polistirex
Extended-Release Suspension (Cough Suppressant)

COUGH

Please visit our web site:
www.delsym.com

Age 4+
For Children & Adults

12 HOUR COUGH RELIEF

Alcohol-free
148 mL (5 fl oz) Day or Night

Orange Flavored Liquid

DELSYM® DOSING
SHAKE WELL BEFORE USE.
Measure only with dosing cup provided. Do not use dosing cup with other products. mL = milliliter

Age (yr)	Dose
12 years to adult	10 mL EVERY 12 HOURS
6 to under 12	5 mL EVERY 12 HOURS
4 to under 6	2.5 mL EVERY 12 HOURS
Under 4	Do not use

See back panel for full dosing directions.

Dosing Cup Included

Contains No Fever Reducer or Pain Reliever

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

3-63824-27665-6
(UPC @ 100% truncated)

Lot No.:
Exp. Date:

UPC area is 1/16" below score and is 1-7/32" w x 11/16" h

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
06/19/2014