Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	70%
Headers:	8.00 pt	Leading:	6.50 pt	Barlines:	2.50 pt		
Subheaders:	6.00 pt	Drug Facts (continued):	8.00 pt	Hairlines:	0.50 pt	Average Kerning:	0

McNeil Submission Artwork

(b) (4)

Active ingredient (in each caplet) — **Purpose**

Acetaminophen 650 mg ..... Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:
■ muscle aches ■ backache ■ minor pain of arthritis ■ toothache
■ premenstrual and menstrual cramps ■ headache ■ the common cold
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Do not use** ■ with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if you have** the following disease. **Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin. **Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not take more than directed (see overdose warning).**
Adults and children 12 years and over: ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor.
Children under 12 years: ■ do not use

Other Information ■ **store between 20-25°C (68-77°F)** ■ **do not use if the inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients carnauba wax, hydroxypropylcellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? call **1-877-895-3665** (toll-free) or **215-273-6755** (collect)

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2014

TO OPEN: PUSH AND TURN CAP
READ THE LABEL

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

TYLENOL® 8HR
MUSCLE ACHES & PAIN

Acetaminophen Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets

24 Caplets 650 mg each

NDC 50580-309-01

LOT

EXP.

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.70 pt	Barlines: NA pt	
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: -25

McNeil **Submission Artwork**

(b) (4)

Labeling Format Information:	Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 80%	McNeil Submission Artwork
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Kerning: 0	

(b) (4)

Active ingredient (in each caplet) **Purpose**

Acetaminophen 650 mg..... .... Pain reliever/fever reducer

Uses ■ temporarily relieves minor aches and pains due to:

■ muscular aches ■ backache ■ minor pain of arthritis ■ toothache
■ premenstrual and menstrual cramps ■ headache ■ the common cold

■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if you have liver disease. Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

TO OPEN: PUSH AND TURN CAP
READ THE LABEL

NDC 50580-309-02

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

TYLENOL® 8HR

MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets



Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ **do not take more than directed (see overdose warning)**

Adults and children 12 years and over: ■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor

Children under 12 years: ■ do not use

Other information ■ store between 20-25°C (68-77°F) ■ **do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? **call 1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA

©McNEIL-PPC, INC. 2014

00000000

LOT:

EXP.

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 75%
Headers: 6.00 pt	Leading: 6.50 pt	Barlines: NA pt	
Subheaders: 6.00 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: 0

McNeil Submission Artwork

(b) (4)

Labeling Format Information:	Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: 9.00 pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%	McNeil
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: 2.50 pt		
Subheaders: 6.00 pt	Drug Facts (continued): 8.00 pt	Hairlines: 0.50 pt	Average Kerning: 0	

(b) (4)

Active ingredient (in each caplet) *Purpose*

Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses ■ temporar ly relieves minor aches and pains due to:

- ■ muscular aches
- ■ backache
- ■ minor pain of arthritis
- ■ toothache
- ■ premenstrual and menstrual cramps
- ■ headache
- ■ the common cold

■ temporar ly reduces fever

Warnings

Liver warning: This p oduct contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ w th other drugs containing acetaminophen ■ 3 or more alcoholic d inks every day while using this p oduct. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. **Do not use** ■ w th any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pha macist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this p oduct. **Ask a doctor before use if you have** liver disease. **Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin. **Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professiona before use.

00000000

EXP

LOT

TO OPEN: PUSH AND TURN CAP

READ THE LABEL

NDC 50580-309-03

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

TYLENOL® 8HR

MUSCLE ACHES & PAIN

Acetaminophen

Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets

225 Caplets*

650 mg each

Keep out of reach of children.

Overdose warning: In case of ove dose, get medica help or contact a Poison Cont ol Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ **do not take more than directed (see overdose warning)**

Adults and children 12 years and over:

- ■ take 2 caplets every 8 hours w th water
- ■ swallow whole; do not crush, chew, split or dissolve
- ■ do not take more than 6 caplets in 24 hours
- ■ do not use fo more than 10 days unless directed by a doctor

Children under 12 years:

■ do not use

Other information

- ■ store between 20-25°C (68-77°F)
- ■ **do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients

carnauba wax, hyd oxyethyl cellulose, hypromellose, magnesium stearate, mic ocrystalline cellulose, modified starch povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:
McNeil Consumer Healthcare Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2014

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%
Headers: 8.00 pt	Leading: 6.50 pt	Barlines: NA pt	
Subheaders: 6.00 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: 0

McNeil Submission Artwork

(b) (4)

To Open: While Folded on Line, Tear At Slit

See New Warning

TYLENOL® 8HR

MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*2 Caplets**
650 mg each

*Capsule-Shaped Bi-Layer Tablets

Active ingredient (in each caplet)	*Purpose*
Acetaminophen 650 mg	**Pain Reliever/Fever Reducer**

Uses temporarily relieves minor aches and pains due to: muscular aches backache minor pain of arthritis toothache sprains menstrual and menstrual cramps headache the common cold temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product. **Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: skin reddening blisters rash. If a skin reaction occurs, stop use and seek medical help right away. Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are allergic to acetaminophen or any of the inactive ingredients in this product. Ask a doctor before use if you have liver disease Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. Stop use and ask a doctor if pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions do not take more than directed (see overdose warning). Adults and children 12 years and over: take 2 caplets every 8 hours with water swallow whole; do not crush, chew, split or dissolve do not take more than 6 caplets in 24 hours do not use for more than 10 days unless directed by a doctor. Children under 12 years: do not use

Other information store between 20-25 C (68-77 F) do not use if pouch is opened

Inactive ingredients carnauba wax, hydroxypropylcellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch povidone powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? call 1-877-895-3665 (toll-free) or 215-273-6755 (collect)

Distributed by: McNeil Consumer Healthcare Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA ©McNEIL-PPC, INC. 2014

00000000

EXP: LOT:

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%
Headers: 4.50 pt	Leading: 4.70 pt	Barlines: NA pt	
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: -25

McNeil Submission Artwork

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
12/23/2014