Panel 7

Reference ID: 3593466

TOP PLY

121/300/033

OVERALL LABEL SIZE .875" HIGH X 3.625" WIDE

BACK OF TOP PLY

BASE

121/300/033



Reference ID: 3593466



BACK OF FRONT PLY



FRONT OF BASE PLY

Lot/EXP Area

3 00067 2039 91 0

UPC FPO

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622 ©2014 2008779
Visit us at www.excedrin.com

02008779

CODE 128 FPO

02008779

CODE 128 FPO

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 CAPLETS®

*CAPLETS-SHAPED TABLETS

NDC 0067-2039-91

Glue

TAMPER EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Drug Facts (continued)

Ask a doctor before use if
• you have never had migraines diagnosed by a health professional
• you have a headache that is different from your usual migraines
• you have the worst headache of your life
• you have fever and stiff neck
• you have headaches beginning after or caused by head injury, exertion, coughing or bending
• you experienced your first headache after the age of 50
• you have daily headaches
• you have a migraine so severe as to require bed rest
• you have liver disease • stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic • you have asthma
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition
• taking any other drug
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur
• ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed**
• adults: take 2 caplets with a glass of water

Drug Facts (continued)

• if symptoms persist or worsen, ask your doctor
• do not take more than 2 caplets in 24 hours, unless directed by a doctor
• under 18 years of age: ask a doctor

Other information
• store at controlled room temperature 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, silicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

CODE 128 FPO

02008779

CODE 128 FPO

02008779

Reference ID: 3593466

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

NDC 0067 2038 91

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 CAPLETS®

*CAPSULE-SHAPED TABLETS

READ ALL PRODUCT INFORMATION BEFORE USING. KEEP BOX FOR IMPORTANT INFORMATION.

Active ingredients (in each caplet)	*Purpose*
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID)*	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use

- treats migraine

Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be a nearly sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock. Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Bleeding warning: The risk of serious stomach bleeding increases if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed • adults: take 2 caplets with a glass of water • for symptoms persist or worsen, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor. Store at controlled room temperature 20°-25°C (68°-77°F). Close cap tightly after use. **Questions or comments?** 1-800-468-7846

Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0672. ©2014. Visit us at www.excedrin.com

Lot/EXP Area

2038780

NDC 0067-2039-92

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

200 CAPLETS*

*CAPSULE-SHAPED TABLETS

Lot/EXP Area

3 0067 2039 92 7

UPC FPO

Distributed by Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0632 ©2014 20087811
Visit us at www.excedrin.com

02008781

CODE 128 FPO

02008781

CODE 128 FPO

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH
"SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 caplets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older
• have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if
• you have never had migraines diagnosed by a health professional
• you have a headache that is different from your usual migraines
• you have the worst headache of your life • you have fever and stiff neck

Drug Facts (continued)

• you have headaches beginning after or caused by head injury, exertion, coughing or bending
• you experienced your first headache after the age of 50
• you have daily headaches • you have a migraine so severe as to require bed rest
• you have liver disease • stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic • you have asthma
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition
• taking any other drug
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur
• ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed**
• adults: take 2 caplets with a glass of water
• if symptoms persist or worsen, ask your doctor
• do not take more than 2 caplets in 24 hours, unless directed by a doctor
• under 18 years of age: ask a doctor

Other information • store at controlled room temperature 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

Glue

EXCEDRIN®
MIGRAINE
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever / Pain Reliever Aid
200 CAPLETS*
*CAPSULE-SHAPED TABLETS

Reference ID: 3593466

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

NDC 0067 2038 92

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

200 CAPLETS®

*CAPSULE-SHAPED TABLETS

READ ALL PRODUCT INFORMATION BEFORE USING. KEEP BOX FOR IMPORTANT INFORMATION.

Active ingredients (in each caplet) — *Purpose*

Acetaminophen 250 mg ..... Pain reliever

Aspirin 250 mg (NSAID)* ..... Pain reliever

Caffeine 65 mg ..... Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be a nearly sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock. Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Risk of bleeding: The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed • adults: take 2 caplets with a glass of water • for symptoms past 5 days or women, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor. Store at controlled room temperature 20°-25°C (68°-77°F). Close cap tightly after use. *Questions or comments?* 1-800-468-7846

Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 Visit us at www.excedrin.com

Lot/EXP Area

2038782

Lot/EXP Area

UPC FPO

3 0067 2039 33 0

Distributed by Novartis Consumer Health, Inc.
Basking Ridge, NJ 07954-6822 ©2014 2008783
Visit us at www.excedrin.com

02008783

CODE 128 FPO

02008783

CODE 128 FPO

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

300 CAPLETS®

*CAPLETS-SHAPED TABLETS

NDC 0067-2039-33

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

300 CAPLETS®

*CAPLETS-SHAPED TABLETS

NDC 0067-2039-33

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

300 CAPLETS®

*CAPLETS-SHAPED TABLETS

TAMPER-EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use
- treats migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use
- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if
- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck

Drug Facts (continued)
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
- taking a prescription drug for: • diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  • feel faint • vomit blood • have bloody or black stools
  • have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
- **do not use more than directed**
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information
- store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

Glue

CODE 128 FPO

02008783

Reference ID: 3593466

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

NDC 0067 2038 33

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

300 CAPLETS®

*CAPSULE-SHAPED TABLETS

READ ALL PRODUCT INFORMATION BEFORE USING. KEEP BOX FOR IMPORTANT INFORMATION.

Active ingredients (in each caplet)	*Purpose*
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be a nearly sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock. Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 caplets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Bleeding warning: The risk of serious stomach bleeding increases if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed • adults: take 2 caplets with a glass of water • for symptoms past or worsen, ask your doctor • do not take more than 2 caplets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor. Store at controlled room temperature 20°-25°C (68°-77°F). Close cap tightly after use. *Questions or comments?* 1-800-468-7846

Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 Visit us at www.excedrin.com

Lot/EXP Area

2038784

127645SC

FPO

TAMPER-EVIDENT PACKETS. DO NOT USE IF SEAL ON INDIVIDUAL PACKET IS BROKEN.

Drug Facts

Active ingredient (in each tablet)	Purpose
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use
- treats migraine

Warnings

Allergy alert: Children and teenagers who have or are recovering from chickenpox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. If you take more than 4 packets in 24 hours, vomiting may occur. If pain gets worse or lasts more than 10 days, stop use and ask a doctor.

If you take too much, you could get liver damage. Ask a doctor before use if you have
- liver disease
- heart disease
- asthma
- high blood pressure
- kidney disease
- stomach problems (such as heartburn)
- a bleeding problem
- diabetes
- ulcers

Ask a doctor before use if you are taking a diuretic.

Do not use
- with any other drug containing acetaminophen
- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- if you have 3 or more alcoholic drinks every day while using this product
- if you are allergic to NSAIDs (nonsteroidal anti-inflammatory drugs) such as ibuprofen or naproxen sodium
- if you have had asthma, nasal problems, or an allergic reaction after taking aspirin or other NSAIDs
- if you have had stomach problems that are made worse by NSAIDs
- if you have ulcers or bleeding problems
- if you have had recent bleeding or clotting disorders
- if you have kidney disease and your doctor has told you not to take NSAIDs
- if you have severe liver disease
- if you are in the last 3 months of pregnancy unless directed by a doctor
- if you are taking a diuretic
- if you are taking a steroid
- if you have phenylketonuria
- if you have an upcoming surgery
- if you are taking other products containing caffeine

Caffeine warning: This nonprescription dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medicines, foods, or beverages while using this product because too much caffeine (usually more than 400 mg) may cause nervousness, shakiness, or sleeplessness.

Stop use and ask a doctor if
- you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- you have ever had an allergic reaction to other NSAIDs (nonsteroidal anti-inflammatory drugs) such as ibuprofen or naproxen sodium
- you have a fever that lasts more than 3 days or recurs
- you have redness or swelling
- you have pain that gets worse or lasts more than 10 days
- you have a headache that is different than your usual migraines
- you have a migraine attack that is different than those you've had before
- you have a migraine or stomach pain that doesn't get better
- you experience ringing in the ears or loss of hearing
- you experience any unexplained rash or skin reaction
- you experience your first headache after the age of 50
- you have a migraine or stomach pain that is severe or lasts longer than usual
- you have high blood pressure, heart disease, liver problems, or kidney disease
- you have a migraine that doesn't get better
- you are taking a diuretic
- you have asthma
- you have a migraine attack that is different than those you've had before
- you have a migraine or stomach pain that doesn't get better
- you have a migraine that is severe or lasts longer than usual
- you have a migraine or stomach pain that is severe or lasts longer than usual
- you have a migraine or stomach pain that is severe or lasts longer than usual
- you have a migraine or stomach pain



Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Do not use** • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain re lever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist **Stop use and ask a** doctor if an a lergic reaction occurs. Seek medical help right away **If pregnant or** breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn ch ld or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. *Directions* • do not use more than directed • adults: take 2 tablets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 tablets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor *Inactive ingredients:* benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, t tanium dioxide Store at 20°-25°C (68°-77°F) *Questions or comments?* 1-800-468-7746

Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 12477 02PC

Lot/EXP Area

TO OPEN, FOLD ALONG DOTTED LINE AND TEAR AT SLIT. TAMPER-EVIDENT PACKET. DO NOT USE IF SEAL IS BROKEN.

NDC 0067-4307-00

READ ALL PRODUCT INFORMATION BEFORE USING.

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

2 TABLETS

Active ingredients (in each tablet)	*Purposes*
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings **Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. **Allergy alert:** Aspirin may cause a severe allergic reaction which may include: • hives • facial swe ling • asthma (wheezing) • shock **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day wh le using this product


49.3
42.9
62.0
70.0
57.3
50.9
3.2
3.18
57.3
70.0
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain re lever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist Stop use and ask a doctor if an a lergic reaction occurs. Seek medical help right away If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn ch ld or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions • do not use more than directed • adults: take 2 tablets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 tablets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor Inactive ingredients: benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, t tanium dioxide Store at 20°-25°C (68°-77°F) Questions or comments? 1-800-468-7746
Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 12477 02PP
Lot/EXP Area
NDC 0067-3037-02
PROFESSIONAL SAMPLE. NOT FOR RESALE.
TO OPEN, FOLD ALONG DOTTED LINE AND TEAR AT NOTCH. TAMPER-EVIDENT PACKET. DO NOT USE IF SEAL IS BROKEN.
EXCEDRIN®
MIGRAINE
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid
2 TABLETS
Active ingredients (in each tablet) Purposes
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID*) Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug
Use • treats migraine
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swe ling • asthma (wheezing) • shock Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day wh le using this product


TAMPER EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING
Drug Facts
Active ingredients (in each tablet) Purposes
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID*) Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug
Use • treats migraine
Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product
Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.
Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
Ask a doctor before use if • you have never had migraines diagnosed by a health professional • you have a headache that is different from your usual migraines • you have the worst headache of your life • you have fever and stiff neck • you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease,
Drug Facts (continued)
liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache
Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis • under a doctor's care for any serious condition • taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer
Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur
• ringing in the ears or loss of hearing occurs
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions • do not use more than directed • adults: take 2 tablets with a glass of water • if symptoms persist
Drug Facts (continued)
or worsen, ask your doctor • do not take more than 2 tablets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor
Other information • store at controlled room temperature 20°-25°C (68°-77°F) • close cap tightly after use • read all product information before using • keep this box for important information.
Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide
Questions or comments? 1-800-468-7746
Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622 ©2014 12477 24C
Visit us at www.excedrin.com
Lot/EXP Area
UPC FPO
3 0067 2037 24 4
12477 24C
CODE 128 FPO
EXCEDRIN®
MIGRAINE
Acetaminophen, Aspirin (NSAID) and Caffeine
Pain Reliever/Pain Reliever Aid
24 TABLETS
Glue
NDC 0067-2037-24
CODE 128 FPO
12477 24C
CODE 128 FPO
12477 24C

BACK OF FRONT PLY

FRONT OF BASE PLY

Lot/EXP Area

3 00067 2037 91 6

UPC FPO

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622 ©2014 12477 100C
Visit us at www.excedrin.com

12477 100C

CODE 128 FPO

12477 100C

CODE 128 FPO

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 TABLETS

MDC 0067-2037-91

Glue

TAMPER EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each tablet)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
• more than 2 tablets in 24 hours, which is the maximum daily amount
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
• are age 60 or older • have had stomach ulcers or bleeding problems
• take a blood thinning (anticoagulant) or steroid drug
• take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
• have 3 or more alcoholic drinks every day while using this product
• take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Drug Facts (continued)

Ask a doctor before use if
• you have never had migraines diagnosed by a health professional
• you have a headache that is different from your usual migraines
• you have the worst headache of your life
• you have fever and stiff neck
• you have headaches beginning after or caused by head injury, exertion, coughing or bending
• you experienced your first headache after the age of 50
• you have daily headaches
• you have a migraine so severe as to require bed rest
• you have liver disease • stomach bleeding warning applies to you
• you have a history of stomach problems, such as heartburn
• you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
• you are taking a diuretic • you have asthma
• you have problems or serious side effects from taking pain relievers or fever reducers
• you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition
• taking any other drug
• taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding:
• feel faint • vomit blood • have bloody or black stools
• have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur
• ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed**
• adults: take 2 tablets with a glass of water

Drug Facts (continued)

• if symptoms persist or worsen, ask your doctor
• do not take more than 2 tablets in 24 hours, unless directed by a doctor
• under 18 years of age: ask a doctor

Other information
• store at controlled room temperature 20°-25°C (68°-77°F)
• close cap tightly after use
• read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, carnauba wax, FD&C blue #1, hydroxypropylcellulose, hypromellose, light mineral oil, microcrystalline cellulose, polysorbate 20, povidone, propylene glycol, silicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

Reference ID: 3593466

DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

NDC 0067 2037 91

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

100 TABLETS

READ ALL PRODUCT INFORMATION BEFORE USING. KEEP BOX FOR IMPORTANT INFORMATION.

Active ingredients (in each tablet)	*Purpose*
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg [NSAID*]	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be a nearly sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock. Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product. Bleeding warning: The risk of serious stomach bleeding increases if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed. Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not use more than directed • adults: take 2 tablets with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 tablets in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor. Store at controlled room temperature 20°-25°C (68°-77°F). Close cap tightly after use. *Questions or comments?* 1-800-468-7846

Dist. by: Novartis Consumer Health, Inc., Parsippany, NJ 07054-0622 ©2014 Visit us at www.excedrin.com

Lot/EXP Area

12477 1001

TAMPER EVIDENT BOTTLE
DO NOT USE IF INNER FOIL SEAL IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each geltab) Purposes

Acetaminophen 250 mg ..... Pain reliever
Aspirin 250 mg (NSAID*) ..... Pain reliever
Caffeine 65 mg ..... Pain reliever aid
*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• skin reddening • blisters • rash
If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
• hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 geltabs in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if • you have never had migraines diagnosed by a health professional • you have a headache that is different from your usual migraines • you have the worst headache of your life • you have fever and stiff neck • you have headaches beginning after or caused by head injury, exertion, coughing or bending • you experienced your first headache after the age of 50 • you have daily headaches • you have a migraine so severe as to require bed rest • you have liver disease • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic

Drug Facts (continued)

• you have asthma • you have problems or serious side effects from taking pain relievers or fever reducers • you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for: • diabetes • gout • arthritis
• under a doctor's care for any serious condition • taking any other drug • taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
• an allergic reaction occurs. Seek medical help right away.
• you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
• your migraine is not relieved or worsens after first dose
• new or unexpected symptoms occur
• ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed**
• adults: take 2 geltabs with a glass of water
• if symptoms persist or worsen, ask your doctor

Drug Facts (continued)

• do not take more than 2 geltabs in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor

Other information • store at controlled room temperature 20°-25°C (68°-77°F) • close cap tightly after use • read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, D&C yellow #10 laka, disodium EDTA, FD&C blue #1 lake, FD&C red #40 lake, ferric oxide, gelatin, glycerin, hydroxypropylcellulose, hypromellose, light mineral oil, maltitol solution, polydextrose, polyethylene glycol, polysorbate 20, povidone, propylene glycol, purified water, silicon emulsion, sorbitan monostearate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622 ©2014 1350630C24
Visit us at www.excedrin.com

UPC FPO

3 00067 2035 24 0

Lot/EXP Area

CODE 128 FPO

1350630C24

CODE 128 FPO

1350630C24

CODE 128 FPO

1350630C24

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

24 GELTABS*

*GELATIN COATED TABLETS

NDC 0067-2035-24

BACK OF FRONT PLY

FRONT OF BASE PLY

Reference ID: 3593466

DO NOT USE IF INNER FOIL SEAL IMPROPERLY VENTED. THIS PRODUCT CARTON IS BROKEN OR MISSING

NDC 0067 2035 50

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever/Pain Reliever Aid

50 GELTABS®

*GELATIN COATED TABLETS

READ ALL PRODUCT INFORMATION BEFORE USING.
KEEP BOX FOR IMPORTANT INFORMATION.

Drug Facts

Active ingredients (in each gel tab) — Purpose

Acetaminophen 250 mg ..... Pain reliever
Aspirin 250 mg (NSAID) ..... Pain reliever
Caffeine 65 mg ..... Pain reliever aid

Uses
• temporarily relieves minor aches and pains due to migraine

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • rash • blisters If a skin reaction occurs, stop use and seek medical help right away. Allergy alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 2 geltabs in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Bleeding warning: This product contains aspirin (NSAID), which may cause a bleeding problem. The chance is higher if you • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or other pain relievers/fever reducers (nonsteroidal anti-inflammatory drug), caffeine, acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions • do not take more than directed • adults: take 2 geltabs with a glass of water • if symptoms persist or worsen, ask your doctor • do not take more than 2 geltabs in 24 hours, unless directed by a doctor • under 18 years of age: ask a doctor Store at controlled room temperature 20°-25°C (68°-77°F). Dispose of properly. Protect from excessive moisture.

Distributed by Novartis Consumer Health, Inc., East Hanover, NJ 07936-1080 ©2014

2035715

Lot/EXP Area

Glue

NDC 0067-2035-91

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

100 GELTABS*

*GELATIN COATED TABLETS

100 GELTABS IN TWO 50 COUNT BOTTLES

135063C100

FPO

135063C100

FPO

135063C100

FPO

135063C100

FPO

Lot/EXP Area

3 0067 2035 91 2

TAMPER-EVIDENT BOTTLES
DO NOT USE IF PRINTED SEAL IS MISSING OR IMPRINTED WITH "SEALED for YOUR PROTECTION" IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each geltab)	Purposes
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID*)	Pain reliever
Caffeine 65 mg	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Use • treats migraine

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
- hives • facial swelling • asthma (wheezing) • shock

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 2 geltabs in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older • have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use • if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Drug Facts (continued)

Ask a doctor before use if • you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic • you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

Ask a doctor or pharmacist before use if you are
- taking a prescription drug for:
  - diabetes • gout • arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

Stop use and ask a doctor if
- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding: • feel faint
- vomit blood • have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • **do not use more than directed**
- adults: take 2 geltabs with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 geltabs in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

Other information • store at controlled room temperature 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

Inactive ingredients benzoic acid, D&C yellow #10 lake, disodium EDTA, FD&C blue #1 lake, FD&C red #40 lake, ferric oxide, gelatin, glycerin, hydroxypropylcellulose, hypromellose, light mineral oil, maltitol solution, microcrystalline cellulose, pepsin, polysorbate 20, povidone, propylene glycol, propyl gallate, simethicone emulsion, sorbitan monolaurate, stearic acid, titanium dioxide

Questions or comments? 1-800-468-7746

Distributed by: Novartis Consumer Health, Inc.
Parsippany, NJ 07054-0622 USA ©2014 135063C00
Visit us at www.excedrin.com

NDC 0067-2035-91

EXCEDRIN®

MIGRAINE

Acetaminophen, Aspirin (NSAID) and Caffeine

Pain Reliever / Pain Reliever Aid

100 GELTABS*

*GELATIN COATED TABLETS

100 GELTABS IN TWO 50 COUNT BOTTLES

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
07/15/2014