Reference ID: 3598813

5HB10800_05_20_14Draft_Clean

Reference ID: 3598813

NET CONTENTS: 10 BOXES OF 100 INDIVIDUAL SWABS
REORDER NO. B10800
NDC 10819-1080-1

Prevantics® Swab

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Antiseptic
Non-sterile Solution
*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to an injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call 1-800-999-6423, M-F (9:00 am-5:00 pm EST)

(01)30310819000018

8GB10800_05_20_14Draft

Each swab contains 0.034 fl.oz. (1 mL)

PDI® **Professional Disposables International, Inc.**, Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8GB10800_05_20_14Draft_Clean

Reference ID: 3598813

REORDER NO. B11400
NDC 10819-1080-2

Prevantics® Swab

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Antiseptic
Non-sterile Solution
Packaging is sterile when in sterile kit *equivalent to 167 mg of Chlorhexidine Gluconate per pouch
Contains: 3000 Individual Swabs, 0.034 fl. oz. (1 mL) Each

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to an injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

(01)30310819000865

8KB11400_05_20_14Draft

Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8KB11400_05_20_14Draft_Clean

Reference ID: 3598813

Prevantics® Swab

REORDER NO. B11400
NDC 10819-1080-2

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab] Antiseptic Non-sterile Solution
Packaging is sterile when in sterile kit
*equivalent to 167 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to an injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition. ▶

Drug Facts (continued)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swab is approximately 2.5 by 2.5 inches (6 by 6 cm)
- tear open packet and remove swab. Do not unfold swab.
- prior to injection, apply swab to the procedure site by holding swab between thumb and index finger. Apply swab to skin using repeated back-and-forth strokes for 15 seconds.
- allow the prepped area to air dry for 30 seconds
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

Each Swab contains 0.034 fl. oz. (1 mL)

PDI
Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 • USA • 1-800-999-6423
www.pdihc.com

5KB114IS_05_20_14Draft

5KB114IS_05_20_14Draft_Clean

Reference ID: 3598813

NDC 10819 4075 1 REORDER NO. S40750

Prevantics® Swabstick

TEAR ALONG PERFORATION FOR DISPENSING

Drug Facts

Active Ingredients — **Purpose**
Chlorhexidine gluconate 3.15% w/v — Antiseptic
Isopropyl alcohol 70% v/v — Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings
- For external use only
- Flammable, keep away from fire or flame
- Do not use with electrocautery procedures

Do not use
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- For lumbar puncture or contact with the meninges
- On open skin wounds or as a general skin cleanser
- Under occlusive patch

When using this product keep out of eyes, ears, mouth and mucous membranes. May cause severe or permanent injury if placed on broad surfaces of the body or used with occlusive dressings. Discontinue use if irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

Stop use and ask a doctor If irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
- Use when hands, nares, or nares under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick: approximately 6 x 6 inches (30 by 30 centimeters or 3 by 5 inches [7.5 by 12.5 cm])
- peel apart the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip
- prior to surgery or injection: Apply the skin to the procedure site
- press one of the dark foam tips against the prepped skin and vigorously back and forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by applying to the same site
- allow the prepped area to dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information
- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive Ingredient
USP purified water

Questions? Call 1-800-999-6423 M-F (9:00 am-5:00 pm EST)

Contains: 50 Individual Swabsticks, 0.054 fl.oz. (1.6 mL) Each

PDI

5HS40750_05_20_14Draft

NDC 10819 4075 1 REORDER NO. S40750

Prevantics® Swabstick

Do not use hypochlorite bleach on fabrics exposed to chlorhexidine gluconate.

PDI

Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 USA
1-800-999-6423
www.pdihc.com

(01)20310819000028

NDC 10819 4075 1 REORDER NO. S40750

Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

- **For external use only**
- Single step, broad spectrum, persistent antiseptic that significantly reduces the number of microorganisms on intact skin
- Use for the preparation of the patient's skin prior to surgery or injection
- Helps reduce bacteria that potentially can cause skin infection
- Latex free
- For hospital and professional use only

Contains:
50 Individual Swabsticks,
0.054 fl.oz. (1.6 mL) Each

PDI

NDC 10819 4075 1 REORDER NO. S40750

Prevantics® Swabstick

CONTENTS: 50 packets of swabsticks premoistened with 1.6 mL of a solution of 3.15% (w/v) Chlorhexidine gluconate, 70% (v/v) Isopropyl alcohol, and USP purified water

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)

PDI

LOT UNVARNISHED AREA

EXP. UNVARNISHED AREA

5HS40750_05_20_14Draft_Clean

Reference ID: 3598813

NET CONTENTS: 10 BOXES OF 50 INDIVIDUAL SWABSTICKS

Prevantics® Swabstick

REORDER NO. S40750
NDC 10819-4075-1

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use
- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. ▶
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)
- peel apart the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

Each swabstick contains 0.054 fl.oz. (1.6 mL)

PDI Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

(01)30310819000025

8GS40750_05_20_14Draft

8GS40750_05_20_14Draft_Clean

Reference ID: 3598813

Prevantics® Swabstick

REORDER NO. S32450
NDC 10819-4075-2

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit *equivalent to 268 mg of Chlorhexidine Gluconate per pouch
Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

Drug Facts

Active ingredients	*Purpose*
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site:3 by 5 inches (7.5 by 12.5 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

(01)30310819000872
8KS32450_05_20_14Draft

PDI

Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8KS32450_05_20_14Draft_Clean

REORDER NO. S32450
NDC 10819-4075-2

Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit

*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures ▶

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute ▶

Drug Facts (continued)

- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient
USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

Each swabstick contains 0.054 fl. oz. (1.6 mL)

PDI Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdihc.com

5KS324 S_05_20_14Dra t

5KS324IS_05_20_14Draft_Clean

Reference ID: 3598813



5HS41950_05_20_14Draft_Clean

Reference ID: 3598813

NET CONTENTS: 10 BOXES OF 30 INDIVIDUAL SWABSTICKS

Prevantics® Maxi Swabstick

REORDER NO. S41950
NDC 10819-4075-4

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution

*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)
- peel apart the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

(01)30310819000032

8GS41950_05_20_14Draft

Each swabstick contains 0.172 fl.oz. (5.1 mL)

PDI Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8GS41950_05_20_14Draft_Clean

Reference ID: 3598813

Prevantics® Maxi Swabstick

REORDER NO. S27350
NDC 10819-4075-3

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit
*equivalent to 855 mg of Chlorhexidine Gluconate per pouch
Contains: 300 Individual Swabsticks, 0.172 fl. oz. (5.1 mL) Each

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient USP purified water

Questions?

Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

(01)30310819000889
8L27350_05_20_14Draft

PDI Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8L27350_05_20_14Draft_Clean

Reference ID: 3598813

REORDER NO. S27350
NDC 10819-4075-3

Prevantics® Maxi Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit
*equivalent to 855 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures ▶

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor. ▶

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 7 by 7 inches (18 by 18 cm) or moist site: 3 by 7.5 inches (7.5 by 19 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site ▶

Drug Facts (continued)

- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

Each Maxi Swabstick contains 0.172 fl. oz. (5.1 mL)

PDI Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdihc.com

5KS273IS_05_20_14Draft

5KS273IS_05_20_14Draft_Clean

Reference ID: 3598813

Prevantics® Swabstick

REORDER NO. S42850
NDC 10819-4075-5

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch
Contains: 500 Individual Swabsticks, 0.054 fl. oz. (1.6 mL) Each

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm)
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute
- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store at 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

(01)30310819000995
8KS428CK_05_20_14Draft

PDI Professional Disposables International, Inc., Orangeburg, NY 10962-1376
USA • 1-800-999-6423 • www.pdihc.com

8KS428CK_05_20_14Draft_Clean

Reference ID: 3598813

REORDER NO. S42850
NDC 10819-4075-5

Prevantics® Swabstick

[Chlorhexidine Gluconate (3.15%)* and Isopropyl Alcohol (70%) Swab]
Patient preoperative skin preparation
Non-sterile Solution
Packaging is sterile when in sterile kit
*equivalent to 268 mg of Chlorhexidine Gluconate per pouch

Drug Facts

Active ingredients	Purpose
Chlorhexidine gluconate 3.15% w/v	Antiseptic
Isopropyl alcohol 70% v/v	Antiseptic

Use For the preparation of the patient's skin prior to surgery or injection

Warnings

- **For external use only**
- **Flammable, keep away from fire or flame**
- Do not use with electrocautery procedures

Drug Facts (continued)

Do not use

- on patients with known allergies to chlorhexidine gluconate or isopropyl alcohol
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser
- under occlusive patch

When using this product keep out of eyes, ears, mouth, and mucous membranes. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Drug Facts (continued)

Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.
- maximum treatment area for one swabstick is approximately: dry site: 4 by 4 inches (10 by 10 cm) or moist site: 3 by 5 inches (7.5 by 12.5 cm).
- tear open the pouch to expose the swabstick
- remove the swabstick from package. Do not touch the foam tip.
- prior to surgery or injection, apply swabstick to the procedure site
- place one flat side of foam tip to the proposed skin site and prep the skin in **vigorous** back-and-forth repeated strokes for one (1) minute

Drug Facts (continued)

- turn the swabstick over (unused side of the foam tip) and repeat the procedure by prepping for one (1) minute
- allow the prepped area to air dry for one and one half minutes (1.5 minutes)
- do not blot or wipe dry
- discard after a single use

Other information

- store between 68-77°F (20-25°C)
- avoid freezing and excessive heat above 104°F (40°C)
- for hospital and professional use only

Inactive ingredient

USP purified water

Questions? Call **1-800-999-6423**, M-F (9:00 am-5:00 pm EST)

Each swabstick contains 0.054 fl. oz. (1.6 mL)

PDI® Professional Disposables International, Inc.
Orangeburg, NY 10962-1376 • USA • 1-800-999-6423 • www.pdihc.com

5KS428IS_05_20_14Draft