



(b) (4)

3.625

(b) (4)

3.875"

Drug Facts		Drug Facts (continued)	
Active ingredient (in each dose) Purpose Polyethylene Glycol 3350, 17 g (cap filled to line)..... Osmotic Laxative		If pregnant or breast-feeding, ask a health professional before use.	
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days		Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor Ask a doctor before use if you have ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug When using this product you may have loose, watery, more frequent stools Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week		Directions ■ do not take more than directed unless advised by your doctor ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) ■ adults and children 17 years of age and older: ■ fill to top of white section in cap which is marked to indicate the correct dose (17g) ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor	
		Other information ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken	
		Inactive ingredients none	
		Questions or comments? 1-800-MiraLAX (1-800-647-2529) or www.MiraLAX.com	
38889-02		• Sugar Free 33069928	



(b) (4)

S-08-91r1 MiraLAX 30 Dose Fron Label

3.625

3.875"

<p>Drug Facts</p> <p>Active ingredient (in each dose) Purpose Polyethylene Glycol 3350, 17 g (cap filled to line)..... Osmotic Laxative</p> <p>Use</p> <ul style="list-style-type: none"> ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to polyethylene glycol</p> <p>Do not use if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ nausea, vomiting or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome <p>Ask a doctor or pharmacist before use if you are taking a prescription drug</p> <p>When using this product you may have loose, watery, more frequent stools</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you get diarrhea ■ you need to use a laxative for longer than 1 week 	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ do not take more than directed unless advised by your doctor ■ the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap) ■ adults and children 17 years of age and older: <ul style="list-style-type: none"> ■ fill to top of white section in cap which is marked to indicate the correct dose (17g) ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ use once a day ■ use no more than 7 days ■ children 16 years of age or under: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ store at 20° - 25°C (68° - 77°F) ■ tamper-evident: do not use if foil seal under cap, printed with "SEALED for YOUR PROTECTION" is missing, open or broken <p>Inactive ingredients none</p> <p>Questions or comments? 1-800-MiraLAX (1-800-647-2529) or www.MiraLAX.com</p>
---	---

• Sugar Free 38500-00

(b) (4)



(b) (4)

