

No Varnish

MUST BE REFRIGERATED

No Varnish

No Varnish

No Varnish

No Varnish

Must be refrigerated

Picato[®]
(ingenol mebutate) gel
0.015%

3 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Dosage: Apply one tube per day to the treatment area on the face and scalp.
See insert for complete information.
Store in a refrigerator at 36°F - 46°F (2°C - 8°C); excursions permitted between 32°F - 59°F (0°C - 15°C) (see USP for controlled cold temperature). Protect from freezing.
Discard after single use. Keep out of reach of children.
Composition: Each gram of Picato[®] (ingenol mebutate) gel, 0.015% contains 150 mcg of ingenol mebutate in a gel base of isopropyl alcohol, hydroxyethyl cellulose, citric acid monohydrate, sodium citrate, benzyl alcohol and purified water.
To report SUSPECTED ADVERSE REACTIONS, contact LEO Pharma Inc. at Phone 1-877-494-4536 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

026281-XX

LOT:
EXP:

Must be refrigerated

Must be refrigerated

No Varnish



Manufactured by:
LEO Laboratories Ltd., (LEO Pharma), Dublin 12, Ireland
or
DPT Laboratories, Ltd., San Antonio, Texas 78215 USA
Net Wt. 0.47 g in each tube
3 Unit Dose Tubes

Picato[®]
(ingenol mebutate) gel
0.015%
Rx only

No Varnish

NDC 50222-502-91

Rx only

Picato[®]
(ingenol mebutate) gel
0.015%

For Topical Use on Face and Scalp Only

Not for Oral, Ophthalmic, or Intravaginal Use


3 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Physician's Sample - Not For Sale

Picato[®]
(ingenol mebutate) gel
0.015%

3 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Must be refrigerated



INSIDE

No Varnish



MUST BE REFRIGERATED



No Varnish

No Varnish

No Varnish

No Varnish

Must be refrigerated

Picato[®]
(ingenol mebutate) gel
0.015%
3 Unit Dose Tubes
Net Wt. 0.47 g in each tube



Dosage: Apply one tube per day to the treatment area on the face and scalp.
See insert for complete information.
Store in a refrigerator at 36°F–46°F (2°C–8°C); excursions permitted between 32°F–59°F (0°C–15°C) (see USP for controlled cold temperature). Protect from freezing.
Discard after single use. Keep out of reach of children.
Composition: Each gram of Picato[®] (ingenol mebutate) gel, 0.015% contains 150 mcg of ingenol mebutate in a gel base of isopropyl alcohol, hydroxyethyl cellulose, citric acid monohydrate, sodium citrate, benzyl alcohol and purified water.
To report SUSPECTED ADVERSE REACTIONS, contact LEO Pharma Inc. at Phone 1-877-494-4536 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

025667-XX

LOT:
EXP:

Must be refrigerated

Must be refrigerated

No Varnish



Manufactured by:
LEO Laboratories Ltd, (LEO Pharma), Dublin 12, Ireland
or
DPT Laboratories, Ltd, San Antonio, Texas 78215 USA
Distributed by:
LEO Pharma Inc., 1 Sylvan Way,
Parlpspany, NJ 07054, USA, 1-877-494-4536
U.S. Patent Nos. US6432452, US6787161

Picato[®]
(ingenol mebutate) gel
0.015%
3 Unit Dose Tubes
Net Wt. 0.47 g in each tube
LEO[®]
Rx only

No Varnish

NDC 50222-502-47

Rx only

Picato[®]
(ingenol mebutate) gel
0.015%

For Topical Use on Face and Scalp Only
Not for Oral, Ophthalmic, or Intravaginal Use


3 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Picato[®]
(ingenol mebutate) gel
0.015%

3 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Must be refrigerated

OUTSIDE



No Varnish

MUST BE REFRIGERATED

No Varnish

No Varnish

No Varnish

No Varnish

Must be refrigerated

Must be refrigerated

LOT:
EXP:
025668-XX

Must be refrigerated

Picato[®]
(ingenol mebutate) gel
0.05%

2 Unit Dose Tubes
Net Wt. 0.47 g in each tube



Dosage: Apply one tube per day to the treatment area on the trunk and extremities. See insert for complete information.
Store in a refrigerator at 36°F - 46°F (2°C - 8°C); excursions permitted between 32°F - 59°F (0°C - 15°C) (see USP for controlled cold temperature). Protect from freezing. Discard after single use. Keep out of reach of children.
Composition: Each gram of Picato[®] (ingenol mebutate) gel, 0.05% contains 500 mcg of ingenol mebutate in a gel base of isopropyl alcohol, hydroxyethyl cellulose, citric acid monohydrate, sodium citrate, benzyl alcohol and purified water.
To report SUSPECTED ADVERSE REACTIONS, contact LEO Pharma Inc. at Phone 1-877-494-4536 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

No Varnish

Picato[®]
(ingenol mebutate) gel
0.05%
2 Unit Dose Tubes
Net Wt. 0.47 g in each tube
Rx only
LEO[®]
Manufactured by:
LEO Laboratories Ltd., (LEO Pharma), Dublin 12, Ireland
or
DPT Laboratories, Ltd., San Antonio, Texas 78215 USA
Distributed by:
LEO Pharma Inc., 1 Sylvan Way,
Parlappan, NJ 07054, USA, 1-877-494-4536
U.S. Patent Nos. US6432452, US6787161
LEO[®]

No Varnish

NDC 50222-503-47

Rx only


Picato[®]
(ingenol mebutate) gel
0.05%

For Topical Use on Trunk and Extremities Only
Not for Oral, Ophthalmic, or Intravaginal Use

2 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Picato[®]
(ingenol mebutate) gel
0.05%
2 Unit Dose Tubes
Net Wt. 0.47 g in each tube

Must be refrigerated



025668-XX

No Varnish

No Varnish

No Varnish


No Varnish

No Varnish

No Varnish

No Varnish

No Varnish



D



C



B



A