Gloss and Matt Varnish -
Full size file on Varnish layer





BASE



RELEASED BACK

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	1 pt, .5 pt

UNCOATED
NO COPY
NO PRINT

128 Code: 048361

128 CODE AREA

128 Code: 048361

128 CODE AREA

128 Code: 048361

128 CODE AREA

MICRO PRINT

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) — **Purpose**

Esomeprazole 20 mg ..... Acid reducer

(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free 1-866-226-1600

128 CODE AREA

128 Code: 048361

Nexium® 24HR | Clinically Proven to **Treat Frequent Heartburn**

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium® (esomeprazole magnesium) Delayed Release Capsules 22.3 mg/Acid Reducer 24HR

Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

NEW

Treats Frequent Heartburn

Nexium® (esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

2 CAPSULES
First two doses of a 14-day course of treatment.

SAMPLE - NOT FOR SALE

May take 1 to 4 days for full effect

128 CODE AREA

128 Code: 048361

1 BOTTLE INSIDE

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.

128 CODE AREA

128 Code: 048361

PAA048361

FPO

3 0573 2450 01 8

128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON
128 Code: 048361

MICRO PRINT

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.

3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
Data Matrix: 048361

Gloss and Matt Varnish -
Full size file on Varnish layer



PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Gloss and Matt Varnish -
Full size file on Varnish layer







PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	3.5 pt
• SPACE BEFORE BULLET	2 ems
• WARNING BOX LINE, HAIRLINES	1 pt, .5 pt

UNCOATED
NO COPY
NO PRINT

128 Code: 048376

128 CODE AREA

128 Code: 048376

128 CODE AREA

128 Code: 048376

128 CODE AREA

MICRO PRINT

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) — **Purpose**

Esomeprazole 20 mg ..... Acid reducer

(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free 1-866-226-1600

128 CODE AREA

128 Code: 048376

128 CODE AREA - SUBSET C 10 MIL DENSITY
WITH 4 OR 6 DIGITS (CODE TO EXTEND
TO EDGE OF FLAP - UNVARNISHED)
PRINTED SIDE OF CARTON
128 Code: 048376

MICRO PRINT

Nexium® 24HR | Clinically Proven to **Treat Frequent Heartburn**

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium® (esomeprazole magnesium) Delayed Release Capsules 22.3 mg/Acid Reducer 24HR

Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

NEW

Treats Frequent Heartburn

Nexium® (esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

14 CAPSULES
One 14-day course of treatment

May take 1 to 4 days for full effect

128 CODE AREA

128 Code: 048376

LOT & EXP AREA.
DARK COLOR FOR
LASER PRINTING.

PAA048376

3/16" DATA MATRIX CODE
WITH 1/8" QUIET ZONE
ON ALL SIDES
Data Matrix: 048376

1 BOTTLE INSIDE

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.

128 CODE AREA

128 Code: 048376

Gloss and Matt Varnish -
Full size file on Varnish layer



PRODUCTION (NOT COLOR ACCURATE)

(b) (4)

Gloss and Matt Varnish -
Full size file on Varnish layers





PRODUCTION (NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Gloss and Matt Varnish -
Full size file on Varnish layer



UNCOATED
NO COPY
NO PRINT

128 Code = 048374

128 CODE AREA

128 Code = 048374

128 CODE AREA

128 Code = 048374

128 CODE AREA

MICRO PRINT

KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) ***Purpose***

Esomeprazole 20 mg ..... Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking
- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment
- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (If needed)
- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?

call toll-free 1-866-226-1600

Nexium® 24HR | Clinically Proven to **Treat Frequent Heartburn**

Tips for Managing Heartburn
- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium® (esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.

Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

NEW **Treats Frequent Heartburn**

Nexium®
(esomeprazole magnesium)
Delayed-Release Capsules
22.3 mg/Acid Reducer
24HR

28 CAPSULES
Two 14-day courses of treatment

May take 1 to 4 days for full effect

PAA048374

FPO
3 0573 2450 28 5

2 BOTTLES INSIDE

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.

128 CODE AREA

128 CODE AREA

128 CODE AREA - SUBSET C 10 MIL DENSITY
WITH 4 OR 6 DIGITS (CODE TO EXTEND
TO EDGE OF FLAP - UNVARNISHED)
PRINTED SIDE OF CARTON
128 Code = 048374

MICRO PRINT

128 Code = 048374

LOT & EXP AREA.
DARK COLOR FOR
LASER PRINTING.

3/16" DATA MATRIX CODE
WITH 1/8" QUIET ZONE
ON ALL SIDES
DATA MATRIX CODE = 048374

128 Code = 048374

PRODUCTION (NOT COLOR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.

Nexium®
(esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer
24HR

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) ***Purpose***
Esomeprazole 20 mg ..... Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking
- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment
- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
- keep the card. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?

call toll-free 1-866-226-1600

FPO
3 0573 2450 29 2

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.
Made in France
For most recent product information, visit www.Nexium24HR.com
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

LOT/EXP
WHITE HOLDOUT
REMOVE VARNISH

PAA048365

Inside bleed

Outside bleed

2 Pack

28 CAPSULES
Two 14-day courses of treatment

May take 1 to 4 days for full effect

NEW

Treats Frequent Heartburn

Nexium® 24HR
(esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer

PRINT SIDE

Gloss and Matt Varnish - Full size file on Varnish layers



PRODUCTION (NOT COL OR ACCURATE)

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Heat Seal Area
No critical text in

UNCOATED
NO COPY
NO PRINT

128 CODES = 048372

128 CODE AREA

128 CODE AREA

128 CODE AREA

128 CODE AREA

128 CODE AREA

MICRO PRINT

MICRO PRINT

Gloss and Matt Varnish -
Full size file on Varnish layer



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) — **Purpose**
Esomeprazole 20 mg ..... Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses
- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings
Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:
- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking
- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

Drug Facts (continued)
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment
- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)
- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information
- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?
call toll-free 1-866-226-1600

Tips for Managing Heartburn
- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

Nexium® (esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

Marketed by:
Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.

Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

Nexium® 24HR

Clinically Proven to **Treat Frequent Heartburn**

NEW Treats Frequent Heartburn

Nexium®
(esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

42 CAPSULES
Three 14-day courses of treatment

May take 1 to 4 days for full effect

PAA048372

FPO
3 0573 2450 42 1

3 BOTTLES INSIDE

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.



128 CODE AREA - SUBSET C 10 MIL DENSITY WITH 4 OR 6 DIGITS (CODE TO EXTEND TO EDGE OF FLAP - UNVARNISHED) PRINTED SIDE OF CARTON

LOT & EXP AREA. DARK COLOR FOR LASER PRINTING.

3/16" DATA MATRIX CODE WITH 1/8" QUIET ZONE ON ALL SIDES
DATA MATRIX CODE = 048372

PRODUCTION

(b) (4)

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	10 pt
• DRUG FACTS CONTINUED	9 pt
• HEADINGS	9 pt
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	6 pt
• SPACE BEFORE BULLET	N/A
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

PRODUCTION

(NOT COLOR ACCURATE)

(b) (4)

Do Not Use if seal under bottle cap imprinted with *"SEALED for YOUR PROTECTION"* or yellow band around the center of each capsule is broken or missing.

Nexium®

(esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

Drug Facts

Active ingredient (in each capsule) ***Purpose***

Esomeprazole 20 mg ..... Acid reducer

(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**

These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription antifungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- **do not take for more than 14 days or more often than every 4 months unless directed by a doctor**

- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

Other Information

- read the directions and warnings before use
- keep the card. It contains important information.
- store at 20-25°C (68-77°F)

Inactive ingredients

corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

Questions or comments?

call toll-free 1-866-226-1600

FPO

3 0573 2450 43 8

PAA048363

Marketed by: Pfizer, Madison, NJ 07940 USA
© 2014 Pfizer Inc.

Made in France

For most recent product information, visit www.Nexium24HR.com

Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

LOT/EXP
WHITE HOLDOUT
REMOVE VARNISH

DATA MATRIX CODE

WHITE HOLDOUT QUIET AREA

REMOVE VARNISH IN THIS AREA ONLY

DATA MATRIX CODE = 048363

42 CAPSULES

Three 14-day courses of treatment

3 Pack

May take 1 to 4 days for full effect

NEW

Treats Frequent Heartburn

Nexium® (esomeprazole magnesium) Delayed-Release Capsules 22.3 mg/Acid Reducer 24HR

Gloss and Matt Varnish -
Full size file on Varnish layers