Nexium 24HR
Clinically Proven to Treat Frequent Heartburn
NEW
Treats Frequent Heartburn
Nexium®
(esomeprazole magnesium)
Delayed-Release Capsules
22.3 mg/Acid Reducer
24HR
2 CAPSULES
First 2 doses of a 14-day course of treatment
May take 1 to 4 days for full effect
SAMPLE - NOT FOR SALE
Marketed by: Pfizer, Madison, NJ 07940 USA Made in France
For most recent product information, visit www.Nexium24HR.com
© 2014 Pfizer Inc.
Do Not Use if foil blister barrier or yellow band on capsule is broken.
Lot:
Exp:
Tips for Managing Heartburn • Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables. • Eat slowly and do not eat big meals. • Do not eat late at night or just before bedtime. • Do not lie flat or bend over soon after eating. • Raise the head of your bed. • Wear loose-fitting clothing around your stomach. • If you are overweight, lose weight. • If you smoke, quit smoking.
PAA000000
KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.
Drug Facts
Active ingredient (in each capsule) Purpose
Esomeprazole 20 mg ..... Acid reducer
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)
Uses
■ treats frequent heartburn (occurs 2 times per week or more) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
Warnings
Allergy alert: Do not use if you are allergic to esomeprazole
Do not use if you have: ■ trouble or pain in swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor.
Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain
Ask a doctor or pharmacist before use if you are taking
■ warfarin or other oral blood-thinning medicines ■ phenytoin or antifungal or antiyeast medicines ■ digoxin or heart medicine) ■ diazepam (anxiety medicine) ■ tacrolimus (immune system medicine) ■ prescription antiretrovirals (medicines for HIV infection) ■ methotrexate (arthritis medicine)
Stop use and ask a doctor if
■ your heartburn continues or worsens ■ you need to take this product for more than 14 days ■ you need to take more than 1 course of treatment every 4 months ■ you get diarrhea
Drug Facts (continued)
If pregnant or breastfeeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ adults 18 years and older ■ this product: take once a day every 24 hours, everyday for 14 days ■ may take 1 to 4 days for full effect
14-Day Course of Treatment
■ swallow 1 capsule with a glass of water before eating in the morning
■ take every day for 14 days
■ do not take more than 1 capsule a day
■ swallow whole. Do not crush or chew capsules.
■ do not use for more than 14 days unless directed by your doctor
Drug Facts (continued)
Repeated 14-Day Courses (if needed)
■ you may repeat a 14-day course every 4 months
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor
■ children under 18 years of age ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.
Other Information
■ read the warnings before use ■ store carefully. It contains important information ■ store at 20-25°C (68-77°F)
Inactive ingredients
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glycerylmonostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethylcitrate
Questions or comments? call toll-free 1-866-226-1600
FPO
FPO
FPO

(b) (4)


Keep directions and warnings. Do not detach individual blisters until ready to use.
Swallow whole. Do not crush or chew capsules.
First two doses of a 14-day course of treatment.
Do not take for more than 14 days or more often than every 4 months unless directed by a doctor.
PAA048370
Nexium®
24HR
1 Esomeprazole Magnesium
Delayed-Release
Capsule 22.3 mg/Acid Reducer
Marketed by: Pfizer
Madison, NJ 07940 USA
LOT/EXP
PEEL
SEPARATE
PEEL
SEPARATE
Nexium®
24HR
1 Esomeprazole Magnesium
Delayed-Release
Capsule 22.3 mg/Acid Reducer
Marketed by: Pfizer
Madison, NJ 07940 USA
LOT/EXP

(b) (4)