LOT & EXP

LOT & EXP

without pain. Ask doctor before use if • stomach bleeding warning applies to you • you have problems or serious side effects from taking pain relievers or fever reducers • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma • you are taking a diuretic • you have a breathing problem such as emphysema or chronic bronchitis • you have gallbladder problems • you have trouble urinating due to an enlarged prostate gland. Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers, or any other sleep-aid • under a doctor's care for any serious condition • taking any other antihistamines • taking any other drug. When using this product • drowsiness will occur • avoid alcoholic drinks • do not drive a motor vehicle or operate machinery • take with food or milk if stomach upset occurs • the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask doctor if • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. • redness or swelling is present in the painful area • any new symptoms appear • you have difficulty swallowing • it feels like the pill is stuck in your throat. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin-containing products during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. *Directions* • do not take more than directed • drink a full glass of water with each dose • adults and children 12 years and over: take 2 caplets at bedtime • do not take more than 2 caplets in 24 hours • taken with food, this product may take longer to work. *Other information* • read all warnings and directions before use • each caplet contains sodium 20 mg • store at 20-25°C (68-77°F) • avoid high humidity and excessive heat above 40°C (104°F) *Inactive ingredients* carnauba wax, FD&C blue #2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, purified water, talc, titanium dioxide *Questions?* 1-800-385-0589 (Mon-Fri 9-5 EST)

Bayer HealthCare LLC, Morristown, NJ 07960 MADE IN GERMANY XXXXXXXX

Do not use if pouch is torn or open.

ALEVE pm

naproxen sodium, 220 mg (NSAID)/
diphenhydramine HCl, 25 mg
Pain reliever/nighttime sleep-aid

BAYER

2 CAPLETS
*CAPSULE-SHAPED TAB

Active ingredients (in each caplet)	*Purposes*
Diphenhydramine hydrochloride 25 mg	Nighttime sleep-aid
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever

*nonsteroidal anti-inflammatory drug

Uses • for relief of occasional sleeplessness when associated with minor aches and pains • helps you fall asleep and stay asleep *Warnings* **Allergy alert:** Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed **Do not use** • if you have ever had an allergic reaction to any other pain reliever/fever reducer • unless you have time for a full night's sleep • in children under 12 years of age • right before or after heart surgery • with any other product containing diphenhydramine, even one used on skin • if you have sleeplessness

Actual Size 100%

(b) (4)

Reference ID: 3438368



Actual Size 100%

MODIFIED DFL

(b) (4)

Reference ID: 3438368

Actual Size 100%

Reference ID: 3438368

Actual Size 100%

(b) (4)

Reference ID: 3438368

Actual Size 100%

Reference ID: 3438368

Actual Size 100%

(b) (4)

Reference ID: 3438368

Actual Size 100%