


Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Outside Layout

11.05.14

NAS_Childrens_Allergy_24HR_60sprays



Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Drug Facts Only

11.05.14

NAS_Childrens_Allergy_24HR_60sprays

(b) (4)

(b) (4)

(b) (4)


<p>Drug Facts</p> <p>Active ingredient (in each spray) Purpose</p> <p>Triamcinolone acetonide (glucocorticoid) 55 mcg Nasal allergy symptom reliever</p> <p>Uses</p> <p>temporarily relieves these symptoms of hay fever or other upper respiratory allergies:</p> <ul style="list-style-type: none"> ■ nasal congestion ■ runny nose ■ sneezing ■ itchy nose <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ in children under 2 years of age ■ if you have ever had an allergic reaction to any of the ingredients <p>Ask a doctor before use if you</p> <ul style="list-style-type: none"> ■ have had recent nose ulcers or nose surgery ■ have had a nose injury that has not healed ■ are using a steroid medicine for asthma, allergies or skin rash ■ have an eye infection ■ have or had glaucoma or cataracts <p>When using this product</p> <ul style="list-style-type: none"> ■ the growth rate of some children may be slower ■ some symptoms may get better on the first day of treatment. It may take up to one week of daily use to feel the most symptom relief. ■ do not share this bottle with anyone else as this may spread germs ■ remember to tell your doctor about all the medicines you take, including this one <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have, or come into contact with someone who has, chickenpox, measles or tuberculosis ■ you have or develop symptoms of an infection such as a persistent fever ■ you have any change in vision ■ you have severe or frequent nosebleeds <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	<p>Drug Facts (continued)</p> <p>Directions</p> <p>Read insert (inside package) on how to:</p> <ul style="list-style-type: none"> ■ get a new bottle ready (primed) before first use ■ prime bottle again if not used for more than 2 weeks ■ use the spray ■ clean the spray nozzle <p>ADULTS AND CHILDREN 12 YEARS OF AGE AND OLDER</p> <p>adults and children 12 years of age and older</p> <ul style="list-style-type: none"> ■ once daily, spray 2 times into each nostril while sniffing gently ■ once your allergy symptoms improve, reduce to 1 spray in each nostril per day <p>CHILDREN 2 TO UNDER 12 YEARS OF AGE</p> <ul style="list-style-type: none"> ■ the growth rate of some children may be slower while using this product. Talk to your child's doctor if your child needs to use the spray for longer than two months a year. ■ an adult should supervise use <p>children 6 to under 12 years of age</p> <ul style="list-style-type: none"> ■ once daily, spray 1 time into each nostril while sniffing gently ■ if allergy symptoms do not improve, increase to 2 sprays in each nostril per day. Once allergy symptoms improve, reduce to 1 spray in each nostril per day. <p>children 2 to under 6 years of age</p> <ul style="list-style-type: none"> ■ an adult should supervise use ■ once daily, spray 1 time into each nostril while sniffing gently <p>children under 2 years of age</p> <ul style="list-style-type: none"> ■ do not use <p>do not use more than directed</p> <ul style="list-style-type: none"> ■ if you forget a dose, do not double the next dose ■ do not spray into eyes or mouth ■ if allergy symptoms do not improve after one week, stop using and talk to a doctor ■ do not use for the common cold ■ shake well before each use <p>Other information</p> <ul style="list-style-type: none"> ■ do not use if sealed package is torn or opened ■ keep package and insert. They contain important information. ■ store between 20°-25°C (68°-77°F)
---	---

(b) (4)

Children's Nasacort Allergy 24HR 60 Sprays Partial Cover Clamshell - Inside Layout

11.05.14

NAS_Childrens_Allergy_24HR_60sprays




Children's Nasacort Allergy 24HR 60 LBL


11.05.14

NAS_Childrens_Allergy_24HR_60_LBL


PAGE 1 (Front Page)




PAGE 2 (Inside Front Page)




PAGE 3




PAGE 4




PAGE 5




PAGE 6




PAGE 7



PAGE 8



BASE LABEL



(b) (4)

Children's **Nasacort[®] Allergy 24HR** Multi-Symptom Nasal Allergy Spray
Product Information xxxxxxxx Title Panel triamcinolone acetonide (glucocorticoid)
55 mcg per spray/Nasal Allergy Symptom Reliever
FPO

IMPORTANT INFORMATION


- Read both sides of this insert for complete instructions on how to get a bottle ready (primed), how to use the spray bottle and how to clean the spray nozzle.
- Keep this insert as it contains important information.

STEPS TO GET A NEW BOTTLE READY FOR USE (PRIMED)

1. Before first use, a new bottle must be primed.

- Remove cap (Figure A).
- Shake bottle.
- Press and release spray nozzle until a fine mist is produced (as shown in Figure B). This may take several times. Take care not to spray in face.

Figure A



Cap

Spray Nozzle

Bottle

PRIME BOTTLE AGAIN IF NOT USED FOR MORE THAN 2 WEEKS

- Repeat steps for priming a bottle (see above).

USE INSTRUCTIONS

- Blow nose gently to clear nostrils.
- Remove cap, then shake bottle.
- Hold bottle with thumb under bottle and spray nozzle between fingers (as shown in Figure B).
- Press against the outside of your nose with your finger to close off one nostril (as shown in Figure C).
- Place the tip of the spray nozzle into the other nostril. The spray nozzle should not reach far into the nose. Aim nozzle toward back of nose (as shown in Figure C).



Figure C

Correct Position



Figure D

Wrong Position

- While sniffing gently, spray into nostril

IMPORTANT: For complete dosing instructions, See "Directions for use" on next side.

- Repeat steps "4" through "6" for the other nostril.
- After using the nasal spray, wipe nozzle with a tissue and replace cap by pressing it down over the spray nozzle.

NOTE: Avoid blowing nose for 15 minutes after use.

If nozzle does not spray properly, see cleaning instructions on next side.

(continued on side 2)

SIDE 1

