



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart—if possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C red no. 33, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)
Grape Flavor

NDC 50580-603-04

For Ages 2 to 11 Years

Children's Motrin
contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL
Pain Reliever / Fever Reducer

Lasts up to 8 hours

Alcohol Free

Grape Flavor

4 fl oz (120 ml)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purpose (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

146 mm

63,5 mm

NDC 50580-603-04

Children's Motrin®

contains **Ibuprofen**
Oral Suspension (NSAID)
100 mg per 5 mL 4 fl oz (120 mL)

Pain Reliever/Fever Reducer

Grape Flavor

Stomach Bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems while taking aspirin (acetylsalicylic acid) or other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) or takes more or for a longer time than directed. **Severe or persistent stomach or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Call your doctor promptly.** Do not use more than 3 days or administer to children under 3 years of age unless directed by doctor. Do not use until the child has been given any other pain reliever/fever reducer. **Weight before or after heart surgery. Ask a doctor before use if stomach bleeding warning applies by ur child who has a history of stomach problems, such as heartburn or indigestion, or if the child has problems or serious side effects from taking pain relievers or fever reducers or if the child has not been drinking, has lost a lot of fluid due to vomiting or diarrhea, or if the child has high blood pressure, heart disease, liver or kidney disease, or if the child has asthma. Do not use if the child is taking a diuretic. Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition or if not taking any other drug. When using this product, do not take with food or milk. Stomach upset occurs as the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if such experiences or if the following signs of stomach bleeding occur: vomit blood or bloody or black stools; has stomach pain that does not get better; or the child does not get any relief within 1 day (24 hours) of treatment. If fever or pain gets worse or lasts more than 3 days, or redness or swelling is present in the painful area, or any new symptoms appear. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). **Directions:** This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter; tsp = teaspoon; light dose = half dose. If possible, use weight to dose; otherwise use age. Use only enclosed dosing cup. Do not use any other dosing device. If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.**

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-5 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information: Each 5 mL (1 tsp) contains sodium 2 mg, active between 20-25°C (68-77°F). Do not use if bottle cap is imprinted "SAFETY SEAL" is broken or missing. **Active ingredients:** aceclofenac potassium, arbutyrus citric acid, D,L-CALCITRIOL, FD&C blue no. 1, FD&C red no. 40, iron oxide black, polyethylene glycol 400, purified water, sodium benzoate, sucrose, xanthan gum.

Questions or comments? Call 1-877-899-5665 (toll-free) or 215-273-8785 (collect).

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children.

Active ingredient: (in each 5 mL = 7 teaspoonsful) Purpose
 Ibuprofen 100 mg (NSAID) Pain reliever/fever reducer
 nonsteroidal anti-inflammatory drug

Use: Temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache. Reduces fever.

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), rash, skin redness, or stomach bleeding. If an allergic reaction occurs, stop use and seek medical help right away.

Lot: 01000000
Exp: 01000000

(b) (4)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart—if possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)
Original Berry Flavor

NDC 50580-601-20

For Ages 2 to 11 Years

Children's Motrin
contains **Ibuprofen Oral Suspension (NSAID)**
100 mg per 5 mL
Pain Reliever / Fever Reducer

Lasts up to 8 hours

Alcohol Free

Original Berry Flavor

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purpose (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

3 0045- 0192- 04 2

LOT:

EXP:

USA 00000 000



(b) (4)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25° C (68-77° F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?

Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)



Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com

00000000

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purpose (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*.....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:


- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

USA
000000
000



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL®" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil
Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com 00000000

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)
Bubble Gum Flavor

NDC 51980-0404
For Ages 2 to 11 Years
Children's Motrin
contains **Ibuprofen Oral Suspension**
100 mg per 5 mL
Lasts up to 8 hours
Bubble Gum Flavor
4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts


Active ingredient Ibuprofen 100 mg (NSAID)*
Purpose Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily:
relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
reduces fever


Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives, asthma (wheezing), skin reddening, rash, blisters, facial swelling, shock
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use
if the child has ever had an allergic reaction to any other pain reliever/fever reducer
right before or after heart surgery

3 0045-0604-04 0
LOT:
EXP:

USA
000000
000



(b) (4)



Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

* or as directed by a doctor

Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive ingredients acesulfame potassium, anhydrous citric acid, D&C yellow no. 10, FD&C red no. 40, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

McNeil

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
© McNEIL-PPC, INC. 2015
www.motrin.com 00000000

FOR HOSPITAL/GOVERNMENT USE ONLY
For Ages 2 to 11 Years

Children's Motrin
contains Ibuprofen Oral Suspension (NSAID)
Pain Reliever / Fever Reducer
Original Berry Flavor

NDC 50580-601-21

FOR HOSPITAL/GOVERNMENT USE ONLY
For Ages 2 to 11 Years

Children's Motrin
contains Ibuprofen Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer

Original Berry Flavor

Lasts up to 8 hours

Alcohol Free

Not for Retail Sale

4 fl oz (120 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient Purpose (in each 5 mL = 1 teaspoonful)
Ibuprofen 100 mg (NSAID).....Pain reliever/fever reducer

*nonsteroidal anti-inflammatory drug

Uses temporarily:
relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
hives facial swelling
asthma (wheezing) shock
skin reddening rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use
if the child has ever had an allergic reaction to any other pain reliever/fever reducer
right before or after heart surgery

LOT:

EXP:

w 3.9375"

Convenient 2-PACK!

NDC 50580-601-25

NDC 50580-601-25

For Ages 2 to 11 Years

For Ages 2 to 11 Years

Children's Motrin®

Children's Motrin®

contains *Ibuprofen*
Oral Suspension (NSAID)
100 mg per 5 mL

contains *Ibuprofen*
Oral Suspension (NSAID)
100 mg per 5 mL

Pain Reliever / Fever Reducer

Pain Reliever / Fever Reducer

Lasts up to
8 hours

Lasts up to
8 hours

Alcohol Free

Alcohol Free

Original
Berry Flavor

Original
Berry Flavor

h 5.6562"

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNeil-PPC, INC. www.motrin.com




McN-XX-0000

3 0045-0192-08 0

h 0.7"

(b) (4)



Drug Facts (continued)

Ask a doctor before use if:

- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is:

- under a doctor's care for any serious condition
- taking any other drug

When using this product:

- take with food or milk if stomach upset occurs
- do not use more than directed or for longer than directed

Stop use and ask a doctor if:

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
 - child does not get any relief within last day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use—do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose, otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

Drug Facts (continued)

If needed, repeat dose every 6-8 hours. Do not use more than 4 times a day. Replace original bottle cap to maintain child resistance.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)*
under 24	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

*or as directed by a doctor

Other Information

- each 5 mL (1 tsp) contains:
 - 50 mg ibuprofen
 - store between 20-25°C (68-77°F)
- do not use if carton is opened or if carton tape or bottle wrap imprinted "SAFETY SEAL" is broken or missing

Inactive Ingredients: acesulfame potassium, anhydrous citric acid, D-C-phenylalanine, FD&C red no. 40, flavors, glycerin, polyethylene glycol, propyl hydroxybenzoate, purified water, sodium benzoate, sucrose, xanthan gum.

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

3 0 0 4 5 - 0 1 9 2 - 0 1 1

Children's Motrin
contains
Ibuprofen Oral Suspension (NSAID)

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19004 USA
©McNeil-PPC, INC. 2015 www.motrin.com

Original Berry Flavor

For Ages 2 to 11 Years

Motrin
contains
Ibuprofen Oral Suspension
100 mg per 5 mL

Original Berry Flavor
1 fl oz (30 mL)

Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active Ingredient (in each 5 mL = 1 teaspoonful)
Ibuprofen 100mg (NSAID)

Purpose
Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses: temporarily relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters.
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child has had stomach ulcers or bleeding problems.
Do not take with aspirin (anticoagulant) or steroid drug.
Do not take other drugs containing aspirin or other NSAIDs (aspirin, ibuprofen, naproxen, or others) unless directed by a doctor.
Do not take for a longer time than directed.
Severe throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer.
Do not use right before or after heart surgery.

USA
000000
000

LOT:

EXP:

[Barcode]

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

THERESA M MICHELE
04/16/2015