




COMPOSITE VIEW – FOR REFERENCE ONLY




COMPOSITE OUTSIDE VIEW AFTER ASSEMBLY













COMPOSITE VIEW – FOR REFERENCE ONLY

















Drug Facts:	pt Body Text:	pt Bullets:	pt Horizontal Scale:	<i>McNeil</i>)	Submission
Headers: Subheaders:	pt Leading: pt Drug Facts (continued):	pt Barlines: pt Hairlines:	pt Average Tracking:		Artwork
	REDACTED				/ Spray 3x120
				File No.: FDA3010823	Version: D-4
					(b) (4)

REDACTED

2. STEPS TO GET A BOTTLE **READY FOR USE (PRIMED)**

- 1. Shake bottle gently.
- 2. Remove protective cap as shown in Figure A.
- 3. Hold bottle firmly, as shown in Figure B.



firmly pressing down on the white

collar while holding the base of the

bottle with your thumb for 8 sprays,

days, reprime the pump until a

b. If not used for more than 14 days,

rinse the spray tip of the pump

until a fine spray appears,

using the cleaning steps listed in

Section 4. After cleaning, reprime

The pump is now ready to use.


5. a. If not used for two consecutive

fine mist appears.

- RHINOCORT® Allergy Spray for complete instructions on how to get a bottle ready for use (primed), how to use the spray bottle, and 4. Prime the pump by quickly and how to clean the spray nozzle.
- 2. Keep this insert as it contains important information.

1. IMPORTANT INFORMATION

Read entire insert before using



Follow these instructions for daily use of RHINO CORT® Allergy Spray:

Gently blow nose to clear nostrils.

3. INSTRUCTIONS

FOR USE

- 2. Shake bottle gently and remove cap.
- 3. Hold bottle firmly, (See Figure B) Insert spray tip into nostril

(tip should not reach far into nose). Close other nostril with a finger and

lean head slightly forward so the spray will aim toward the back of nose. (See Figure C)

DO NOT spray towards nasal septum (wall between two nostrils). (See Figure D)





- 4. For each spray, press down quickly and firmly on white collar while holding the base of the bottle with your thumb. Breathe gently inward.
- 5. After spraying into your nostril, lean your head slightly backward for a few seconds.
- 6. If a second spray is needed in the same nostril, repeat steps 3 through 5. For complete dosing instructions see Table 1 "Directions for Use".
- 7. Repeat steps 3 through 6, for the other nostril
- 8. Avoid blowing your nose for 15 MINUTES after using RHINOCORT_® Allergy Spray.
- 9. Wipe spray tip with a clean tissue (See Figure E), and replace outer cap. Store the bottle in an upright position.



TABLE 1: **DIRECTIONS FOR USE**

- Before first use, you must get a new bottle ready (primed). See Section 2.
- If bottle is not used for two consecutive days,
- reprime the pump until a fine spray appears. If bottle is not used for more than two weeks, rinse the spray tip of the pump usina After spray a

using the cleaning steps listed in Section 4. After cleaning, reprime the pump until a fine spray appears.				
ADULTS AND CHIL	DREN 12 YEARS OF AGE AND OLDER			
adults and children 12 years of age and older	■ once daily, spray 2 times in o each nostril while sniffing gently ■ once your allergy symp oms improve, reduce o 1 spray in each nostril per day			
CHILDREN 6 TO UNDER 12 YEARS OF AGE				
he g ow h rate of some children may be slower while using his p oduct, Talk o your child's doctor if your child needs to use he				


■ hegow spray for longer han wo mon hs a year,

children 6	■ an adult should supervise use				
to under 12	■ once daily, spray 1 time in o each				
years of age	nostril while sniffing gently				
	■ if allergy symp oms do not imp ove,				
	increase to 2 sprays in each nostril				
	per day. Once allergy symp oms				
	improve, reduce o 1 pray in each				
	nostril per day.				
children under	■ do not use				
6 years of age					
Other important information for use					
= do not use more than directed					

- do not use more than directed
- if you forget a dose, do not double he next dose
- do not spray in o eyes or mou h ■ if allergy symptoms do not imp ove after two weeks, stop using and talk to a doc or
- do not use for he common cold ■ shake well before each use

4. HOW TO CLEAN YOUR RHINOCORT® ALLERGY SPRAY

ise cap and spray tip regularly.



- Remove cap and gently pull spray tip away from the bottle. (See Figure G)
- 2. Wash only cap and spray tip in warm water and rinse in cold tap water, Shake to remove excess water,
- Allow cap and spray tip to air dry completely before reassembling the nasal spray.
- 4. Press and release spray nozzle until a fine spray is produced, taking care not to spray in the face,
- 5. If the spray tip becomes blocked, it can be cleared by repeating steps 1 through 3. Do not unblock the nasal applicator with a pin or other sharp object.

REDACTED

REDACTED

Submission McNeil Artwork

REDACTED

REDACTED

Description: IFU Rhinocort Allergy Spray File No.: FDA3010823

C-3

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.
/s/
THERESA M MICHELE 03/23/2015