Lot
Exp
$5.00
Coupon Inside
NDC 0135-0510-06
gsk
Nicorette®
nicotine polacrilex lozenge, 2 mg
stop smoking aid
Lozenge
Poppac®
Mint
Includes User's Guide
2 mg
FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES AFTER WAKING UP.
If you smoke your first cigarette WITHIN 30 MINUTES of waking up, use Nicorette 4 mg Lozenge
24 LOZENGES, 2 mg Each
Nicorette
nicotine polacrilex lozenge, 2 mg · stop smoking aid
Lozenge
Drug Facts (continued)
use 2 mg nicotine lozenge according to the following 12 week schedule:
Weeks 1 to 6: 1 lozenge every 1 to 2 hours
Weeks 7 to 9: 1 lozenge every 2 to 4 hours
Weeks 10 to 12: 1 lozenge every 4 to 8 hours
• nicotine lozenge is a medicine and must be used a certain way to get the best results
• place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow lozenge.
• you may feel a warm or tingling sensation
• occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 - 30 minutes)
• do not eat or drink 15 minutes before using or while the lozenge is in your mouth
• to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
• do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
• do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
• it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.
Other information
• each lozenge contains: sodium, 18 mg
• Phenylketonurics: Contains Phenylalanine 3.4 mg per lozenge
• store at 20 - 25°C (68 - 77°F)
• keep POPPAC tightly closed and protect from light
Inactive ingredients acacia, aspartame, calcium polycarbophil, corn syrup solids, flavors, lactose, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, soy protein, triethyl citrate, xanthan gum
Questions or comments? call toll-free 1-888-569-1743 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)
PS
NICORETTE, POPPAC and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.
Distributed By:
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108
Made in Switzerland
©2015 GlaxoSmithKline
not for sale to those under 18 years of age
proof of age required
not for sale in vending machines or from any source where proof of age cannot be verified
TAMPER EVIDENT FEATURE: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken.
Retain outer carton for full product uses, directions and warnings.
TO INCREASE YOUR SUCCESS IN QUITTING:
1. You must be motivated to quit.
2. Use Enough - Use at least 9 Nicorette® Lozenges per day during the first six weeks.
3. Use Long Enough - Use Nicorette® Lozenges for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.
Nicorette® POPPAC®
To open vial, push in child resistant band on the POPPAC® container with thumb.
Flip up the top of the POPPAC® and remove lozenge. A small amount of powder on opening of the POPPAC® is normal.
For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.
3 0766-7845-10 5
Drug Facts
Active ingredient (in each lozenge) Purpose
Nicotine polacrilex, 2 mg ..... Stop smoking aid
Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking
Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.
Do not use
• if you are allergic to soya
Ask a doctor before use if you have
• a sodium-restricted diet
• heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
• high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
• stomach ulcer or diabetes
Ask a doctor or pharmacist before use if you are
• using a non-nicotine stop smoking drug
• taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
Stop use and ask a doctor if
• mouth problems occur
• persistent indigestion or severe sore throat occurs
• irregular heartbeat or palpitations occur
• you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
• you have symptoms of an allergic reaction (such as difficulty breathing or rash)
Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
• if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
• before using this product, read the enclosed User's Guide for complete directions and other important information
• begin using the lozenge on your quit day
• if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge
• if you smoke your first cigarette more than 30 minutes after waking up, ▶

(b) (4)


No copy areas.
SAVE
$5.00
on any
Nicorette®
Lozenge
144ct
MANUFACTURER'S COUPON
EXPIRES 12/31/2018
Redeemable at food, drug and mass merchandise stores.
030766-074621

(b) (4)

Lot
Exp
0000000000
Drug Facts (continued)
4 mg nicotine lozenge according to the following 12 week schedule:
Weeks 1 to 6 | Weeks 7 to 9 | Weeks 10 to 12
1 lozenge every 1 to 2 hours | 1 lozenge every 2 to 4 hours | 1 lozenge every 4 to 8 hours
• nicotine lozenge is a medicine and must be used a certain way to get the best results
• place the lozenge in your mouth and allow the lozenge to slowly dissolve (about 20 - 30 minutes). Minimize swallowing. Do not chew or swallow lozenge.
• you may feel a warm or tingling sensation
• occasionally move the lozenge from one side of your mouth to the other until completely dissolved (about 20 - 30 minutes)
• do not eat or drink 15 minutes before using or while the lozenge is in your mouth
• to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
• do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
• do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
• it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.
Other information
• each lozenge contains: sodium, 18 mg
• Phenylketonurics: Contains Phenylalanine 3.4 mg per lozenge
• store at 20 - 25°C (68 - 77°F)
• keep POPPAC tightly closed and protect from light
Inactive ingredients acacia, aspartame, calcium polycarbophil, corn syrup solids, flavors, lactose, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, sodium alginate, sodium carbonate, soy protein, triethyl citrate, xanthan gum
Questions or comments? call toll-free 1-888-569-1743 (English/Spanish) weekdays (9:00 am - 4:30 pm ET)
$5.00 Coupon Inside
NDC 0135-0511-06
gsk
Nicorette®
nicotine polacrilex lozenge, 4 mg
stop smoking aid
Lozenge
Poppac®
Mint
Includes User's Guide
4 mg
FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicorette 2 mg Lozenge
24 LOZENGES, 4 mg Each
Nicorette
nicotine polacrilex lozenge, 4 mg • stop smoking aid
Lozenge
NICORETTE, POPPAC and the NICORETTE sunburst design are registered trademarks of the GSK group of companies.
Distributed By:
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Township, PA 15108
Made in Switzerland
©2015 GlaxoSmithKline
■ not for sale to those under 18 years of age
■ proof of age required
■ not for sale in vending machines or from any source where proof of age cannot be verified
TAMPER EVIDENT FEATURE: Do not use if clear neckband printed "SEALED FOR SAFETY" is missing or broken.
Retain outer carton for full product uses, directions and warnings.
TO INCREASE YOUR SUCCESS IN QUITTING:
1. You must be motivated to quit.
2. Use Enough - Use at least 9 Nicorette® Lozenges per day during the first six weeks.
3. Use Long Enough - Use Nicorette® Lozenges for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.
Nicorette® POPPAC®
To open vial, push in child resistant band on the POPPAC® container with thumb.
Flip up the top of the POPPAC® and remove lozenge. A small amount of powder on opening of the POPPAC® is normal.
For more information and for a FREE individualized stop smoking program, please visit www.Nicorette.com or see inside for more details.
3 0766-7847-10 9
Drug Facts
Active ingredient (in each lozenge) Purpose
Nicotine polacrilex, 4 mg ..... Stop smoking aid
Use • reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking
Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.
Do not use
• if you are allergic to soya
Ask a doctor before use if you have
• a sodium-restricted diet
• heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
• high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
• stomach ulcer or diabetes
Ask a doctor or pharmacist before use if you are
• using a non-nicotine stop smoking drug
• taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.
Stop use and ask a doctor if
• mouth problems occur
• persistent indigestion or severe sore throat occurs
• irregular heartbeat or palpitations occur
• you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat
• you have symptoms of an allergic reaction (such as difficulty breathing or rash)
Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
• if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
• before using this product, read the enclosed User's Guide for complete directions and other important information
• begin using the lozenge on your quit day
• if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge
• if you smoke your first cigarette within 30 minutes of waking up, use ▶
PS
0000000000

(b) (4)


No copy areas.
SAVE
$5.00
on any
Nicorette®
Lozenge
144ct
MANUFACTURER'S COUPON
EXPIRES 12/31/2018
Redeemable at food, drug and mass merchandise stores.
030766-074621

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

SHERRY A STEWART
08/10/2015

THERESA M MICHELE
08/10/2015