

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use REYATAZ safely and effectively. See full prescribing information for REYATAZ.

**REYATAZ<sup>®</sup> (atazanavir) capsules, for oral use**

**REYATAZ<sup>®</sup> (atazanavir) oral powder**

**Initial U.S. Approval: 2003**

### RECENT MAJOR CHANGES

Indications and Usage (1)	6/2014
Dosage and Administration,	
Overview (2.1)	6/2014
Dosage and Administration of REYATAZ Oral Powder in Pediatric Patients (2.4)	6/2014
Contraindications (4)	6/2014
Warnings and Precautions,	
Patients with Phenylketonuria (5.4)	6/2014
Risk of Serious Adverse Reactions Due to Drug Interactions (5.7)	3/2015

### INDICATIONS AND USAGE

REYATAZ is a protease inhibitor indicated for use in combination with other antiretroviral agents for the treatment of HIV-1 infection for patients 3 months and older weighing at least 10 kg. (1)

### DOSAGE AND ADMINISTRATION

- *Treatment-naïve adults:* REYATAZ 300 mg with ritonavir 100 mg once daily with food or REYATAZ 400 mg once daily with food. (2.2)
- *Treatment-experienced adults:* REYATAZ 300 mg with ritonavir 100 mg once daily with food. (2.2)
- *Pediatric patients:* REYATAZ capsule dosage is based on body weight not to exceed the adult dose and must be taken with food. (2.3)
- *REYATAZ oral powder:* Must be taken with ritonavir and food and should not be used in children who weigh less than 10 kg or who weigh 25 kg or more. (2.4)
- *Pregnancy:* REYATAZ 300 mg with ritonavir 100 mg once daily with food, with dosing modifications for some concomitant medications. (2.5)
- *Concomitant therapy:* Dosing modifications may be required. (2.2, 2.3, 2.4, 7)
- *Renal impairment:* Dosing modifications may be required. (2.6)
- *Hepatic impairment:* Dosing modifications may be required. (2.7)

### DOSAGE FORMS AND STRENGTHS

- Capsules: 150 mg, 200 mg, 300 mg. (3, 16)
- Oral powder: 50 mg packet. (3, 16)

### CONTRAINDICATIONS

- REYATAZ is contraindicated in patients with previously demonstrated hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of this product. (4)
- Coadministration with alfuzosin, triazolam, orally administered midazolam, ergot derivatives, rifampin, irinotecan, lovastatin, simvastatin, indinavir, cisapride, pimozide, St. John's wort, nevirapine, and sildenafil when dosed as REVATIO<sup>®</sup>. (4)

### WARNINGS AND PRECAUTIONS

- *Cardiac conduction abnormalities:* PR interval prolongation may occur in some patients. ECG monitoring should be considered in patients with preexisting conduction system disease or when administered with other drugs that may prolong the PR interval. (5.1, 7.3, 12.2, 17)
- *Rash:* Discontinue if severe rash develops. (5.2, 17)
- *Hyperbilirubinemia:* Most patients experience asymptomatic increases in indirect bilirubin, which is reversible upon discontinuation. Do not dose reduce. If a concomitant transaminase increase occurs, evaluate for alternative etiologies. (5.3)
- *Hepatotoxicity:* Patients with hepatitis B or C infection are at risk of increased transaminases or hepatic decompensation. Monitor hepatic laboratory tests prior to therapy and during treatment. (2.5, 5.5, 8.8)
- *Nephrolithiasis and cholelithiasis* have been reported. Consider temporary interruption or discontinuation. (5.6)
- The concomitant use of REYATAZ/ritonavir and certain other medications may result in known or potentially significant drug interactions. Consult the full prescribing information prior to and during treatment for potential drug interactions. (5.7, 7.3)
- Patients receiving REYATAZ may develop new onset or exacerbations of diabetes mellitus/hyperglycemia (5.8), immune reconstitution syndrome (5.9), and redistribution/accumulation of body fat (5.10).
- *Hemophilia:* Spontaneous bleeding may occur and additional factor VIII may be required. (5.11)
- *Phenylketonuria:* REYATAZ oral powder contains phenylalanine which can be harmful to patients with phenylketonuria. (5.4)

### ADVERSE REACTIONS

Most common adverse reactions (≥2%) are nausea, jaundice/scleral icterus, rash, headache, abdominal pain, vomiting, insomnia, peripheral neurologic symptoms, dizziness, myalgia, diarrhea, depression, and fever. (6.1, 6.2)

**To report SUSPECTED ADVERSE REACTIONS, contact Bristol-Myers Squibb at 1-800-721-5072 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### DRUG INTERACTIONS

Coadministration of REYATAZ can alter the concentration of other drugs and other drugs may alter the concentration of atazanavir. The potential drug-drug interactions must be considered prior to and during therapy. (4, 7, 12.3)

### USE IN SPECIFIC POPULATIONS

- *Pregnancy:* Use only if the potential benefit justifies the potential risk. (8.1)
- *Nursing mothers* should be instructed not to breastfeed due to the potential for postnatal HIV transmission. (8.3)
- *Hepatitis B or C co-infection:* Monitor liver enzymes. (5.5, 6.3)
- *Renal impairment:* REYATAZ is not recommended for use in treatment-experienced patients with end stage renal disease managed with hemodialysis. (2.6, 8.7)
- *Hepatic impairment:* REYATAZ is not recommended in patients with severe hepatic impairment. REYATAZ/ritonavir is not recommended in patients with any degree of hepatic impairment. (2.7, 8.8)

**See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.**

**Revised: 03/2015**

**FULL PRESCRIBING INFORMATION: CONTENTS\***

<b>1</b>	<b>INDICATIONS AND USAGE</b>	<b>7</b>	<b>DRUG INTERACTIONS</b>
<b>2</b>	<b>DOSAGE AND ADMINISTRATION</b>	7.1	Potential for REYATAZ to Affect Other Drugs
2.1	Overview	7.2	Potential for Other Drugs to Affect REYATAZ
2.2	Dosage in Adult Patients	7.3	Established and Other Potentially Significant Drug Interactions
2.3	Dosage of REYATAZ Capsules in Pediatric Patients	7.4	Drugs with No Observed or Predicted Interactions with REYATAZ
2.4	Dosage and Administration of REYATAZ Oral Powder in Pediatric Patients	<b>8</b>	<b>USE IN SPECIFIC POPULATIONS</b>
2.5	Dosage Adjustments in Pregnant Patients	8.1	Pregnancy
2.6	Renal Impairment	8.3	Nursing Mothers
2.7	Dosage Adjustments in Patients with Hepatic Impairment	8.4	Pediatric Use
<b>3</b>	<b>DOSAGE FORMS AND STRENGTHS</b>	8.5	Geriatric Use
<b>4</b>	<b>CONTRAINDICATIONS</b>	8.6	Age/Gender
<b>5</b>	<b>WARNINGS AND PRECAUTIONS</b>	8.7	Impaired Renal Function
5.1	Cardiac Conduction Abnormalities	8.8	Impaired Hepatic Function
5.2	Rash	<b>10</b>	<b>OVERDOSAGE</b>
5.3	Hyperbilirubinemia	<b>11</b>	<b>DESCRIPTION</b>
5.4	Patients with Phenylketonuria	<b>12</b>	<b>CLINICAL PHARMACOLOGY</b>
5.5	Hepatotoxicity	12.1	Mechanism of Action
5.6	Nephrolithiasis and Cholelithiasis	12.2	Pharmacodynamics
5.7	Risk of Serious Adverse Reactions Due to Drug Interactions	12.3	Pharmacokinetics
5.8	Diabetes Mellitus/Hyperglycemia	12.4	Microbiology
5.9	Immune Reconstitution Syndrome	<b>13</b>	<b>NONCLINICAL TOXICOLOGY</b>
5.10	Fat Redistribution	13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
5.11	Hemophilia	<b>14</b>	<b>CLINICAL STUDIES</b>
5.12	Resistance/Cross-Resistance	14.1	Adult Patients without Prior Antiretroviral Therapy
<b>6</b>	<b>ADVERSE REACTIONS</b>	14.2	Adult Patients with Prior Antiretroviral Therapy
6.1	Clinical Trial Experience in Adults	14.3	Pediatric Patients
6.2	Clinical Trial Experience in Pediatric Patients	<b>16</b>	<b>HOW SUPPLIED/STORAGE AND HANDLING</b>
6.3	Patients Co-Infected with Hepatitis B and/or Hepatitis C Virus	<b>17</b>	<b>PATIENT COUNSELING INFORMATION</b>
6.4	Postmarketing Experience	* Sections or subsections omitted from the full prescribing information are not listed.	

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

REYATAZ<sup>®</sup> (atazanavir) is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection for patients 3 months and older weighing at least 10 kg.

Limitations of Use:

- REYATAZ is not recommended for use in pediatric patients below the age of 3 months due to the risk of kernicterus.
- Use of REYATAZ/ritonavir in treatment-experienced patients should be guided by the number of baseline primary protease inhibitor resistance substitutions [see *Microbiology (12.4)*].

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Overview

- REYATAZ capsules and oral powder must be taken with food.
- Do not open the capsules.
- The recommended oral dosage of REYATAZ depends on the treatment history of the patient and the use of other coadministered drugs. When coadministered with H<sub>2</sub>-receptor antagonists or proton-pump inhibitors, dose separation may be required [see *Dosage and Administration (2.2, 2.3, 2.4, 2.5) and Drug Interactions (7)*].
- REYATAZ capsules without ritonavir are not recommended for treatment-experienced adult or pediatric patients with prior virologic failure [see *Clinical Studies (14)*].
- REYATAZ oral powder must be taken with ritonavir and is not recommended for use in children who weigh less than 10 kg or who weigh 25 kg or more. [See *Dosage and Administration (2.4)*.]
- Efficacy and safety of REYATAZ with ritonavir when ritonavir is administered in doses greater than 100 mg once daily have not been established. The use of higher ritonavir doses may alter the safety profile of atazanavir (cardiac effects, hyperbilirubinemia) and, therefore, is not recommended. Prescribers should consult the complete prescribing information for ritonavir when using ritonavir.

#### 2.2 Dosage in Adult Patients

Table 1 displays the recommended dosage of REYATAZ capsules in treatment-naïve and treatment-experienced adults. Table 1 also displays recommended dosage of REYATAZ and ritonavir when given concomitantly with other antiretroviral drugs and H<sub>2</sub>-receptor antagonists (H2RA). Ritonavir is required with several REYATAZ dosage regimens (see the ritonavir

complete prescribing information about the safe and effective use of ritonavir). The use of REYATAZ in treatment-experienced adult patients without ritonavir is not recommended.

**Table 1: Recommended REYATAZ and Ritonavir Dosage in Adults<sup>a</sup>**

	REYATAZ Once Daily Dosage	Ritonavir Once Daily Dosage
<b>Treatment-Naive Adult Patients</b>		
recommended regimen	300 mg	100 mg
unable to tolerate ritonavir	400 mg	N/A
in combination with efavirenz	400 mg	100 mg
<b>Treatment-Experienced Adult Patients</b>		
recommended regimen	300 mg	100 mg
in combination with both H2RA and tenofovir	400 mg	100 mg

<sup>a</sup> See *Drug Interactions* (7) for instructions concerning coadministration of acid reducing medications (eg, H2RA or proton pump inhibitors [PPIs]), and other antiretroviral drugs (eg, efavirenz, tenofovir, and didanosine).

## 2.3 Dosage of REYATAZ Capsules in Pediatric Patients

The recommended daily dosage of REYATAZ capsules and ritonavir in pediatric patients (6 years of age to less than 18 years of age) is based on body weight (see Table 2).

**Table 2: Recommended Dosage of REYATAZ Capsules and Ritonavir in Pediatric Patients (6 to less than 18 years of age)<sup>a,b</sup>**

Body weight	REYATAZ Daily Dosage	Ritonavir Daily Dosage
<b>Treatment-Naive and Treatment-Experienced<sup>c</sup></b>		
Less than 15 kg	Capsules not recommended	N/A
15 kg to less than 20 kg	150 mg	100 mg
20 kg to less than 40 kg	200 mg	100 mg
At least 40 kg	300 mg	100 mg
<b>Treatment-Naive, at least 13 years old and cannot tolerate ritonavir<sup>c</sup></b>		
At least 40 kg	400 mg	N/A

<sup>a</sup> Administer REYATAZ capsules and ritonavir simultaneously with food.

<sup>b</sup> The same recommendations regarding the timing and maximum doses of concomitant PPIs and H2RAs in adults also apply to pediatric patients. See *Drug Interactions* (7) for instructions concerning coadministration of acid reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir, and didanosine).

<sup>c</sup> In treatment-experienced patients, REYATAZ capsules must be administered with ritonavir.

## 2.4 Dosage and Administration of REYATAZ Oral Powder in Pediatric Patients

REYATAZ oral powder is for use in treatment-naïve or treatment-experienced pediatric patients who are at least 3 months of age and weighing at least 10 kg and less than 25 kg. REYATAZ oral powder must be mixed with food or beverage for administration and ritonavir must be given immediately afterwards. Table 3 displays the recommended dosage of REYATAZ oral powder and ritonavir.

**Table 3: Recommended Dosage of REYATAZ Oral Powder and Ritonavir in Pediatric Patients (at least 3 months of age and weighing at least 10 kg and less than 25 kg)<sup>a</sup>**

Body Weight	Daily Dosage of REYATAZ Oral Powder	Daily Dosage of Ritonavir Oral Solution
10 kg to less than 15 kg	200 mg (4 packets) <sup>b</sup>	80 mg
15 kg to less than 25 kg	250 mg (5 packets) <sup>b</sup>	80 mg

<sup>a</sup> The same recommendations regarding the timing and maximum doses of concomitant PPIs and H2RAs in adults also apply to pediatric patients. See *Drug Interactions* (7) for instructions concerning coadministration of acid reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir, and didanosine).

<sup>b</sup> Each packet contains 50 mg of REYATAZ.

### Instructions for Mixing REYATAZ Oral Powder [see FDA-approved *Instructions for Use*]

- It is preferable to mix REYATAZ oral powder with food such as applesauce or yogurt. Mixing REYATAZ oral powder with a beverage (milk, infant formula, or water) may be used for infants who can drink from a cup. For young infants (less than 6 months) who cannot eat solid food or drink from a cup, REYATAZ oral powder should be mixed with infant formula and given using an oral dosing syringe. Administration of REYATAZ and infant formula using an infant bottle is not recommended because full dose may not be delivered.
- Determine the number of packets (4 or 5 packets) that are needed.
- Prior to mixing, tap the packet to settle the powder. Use a clean pair of scissors to cut each packet along the dotted line.

- **Mixing with food:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with a minimum of one tablespoon of food (such as applesauce or yogurt). Feed the mixture to the infant or young child. Add an additional one tablespoon of food to the small container, mix, and feed the child the residual mixture.
- **Mixing with a beverage such as milk or water in a small drinking cup:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with a minimum of 30 mL of the beverage. Have the child drink the mixture. Add an additional 15 mL more of beverage to the drinking cup, mix, and have the child drink the residual mixture. If water is used, food should also be taken at the same time.
- **Mixing with liquid infant formula using an oral dosing syringe and a small medicine cup:** Using a spoon, mix the recommended number of REYATAZ oral powder packets with 10 mL of prepared liquid infant formula. Draw up the full amount of the mixture into an oral syringe and administer into either right or left inner cheek of infant. Pour another 10 mL of formula into the medicine cup to rinse off remaining REYATAZ oral powder in cup. Draw up residual mixture into the syringe and administer into either right or left inner cheek of infant.
- Administer ritonavir immediately following REYATAZ powder administration.
- Administer the entire dosage of REYATAZ oral powder (mixed in the food or beverage) within one hour of preparation (may leave the mixture at room temperature during this one hour period). Ensure that the patient eats or drinks all the food or beverage that contains the powder. Additional food may be given after consumption of the entire mixture.

## 2.5 Dosage Adjustments in Pregnant Patients

Table 4 includes the recommended dosage of REYATAZ capsules and ritonavir in treatment-naïve and treatment-experienced pregnant patients. In these patients, REYATAZ must be administered with ritonavir. There are no dosage adjustments for postpartum patients (see Table 1 for the recommended REYATAZ dosage in adults). *[See Use in Specific Populations (8.1).]*

**Table 4: Recommended Dosage of REYATAZ and Ritonavir in Pregnant Patients<sup>a</sup>**

	<b>REYATAZ Once Daily Dosage</b>	<b>Ritonavir Once Daily Dosage</b>
<b>Treatment-Naive and Treatment-Experienced</b>		
Recommended Regimen	300 mg	100 mg
<b>Treatment-Experienced During the Second or Third Trimester When Coadministered with either H2RA or Tenofovir<sup>b</sup></b>		
In combination with either H2RA or tenofovir	400 mg	100 mg

<sup>a</sup> See *Drug Interactions* (7) for instructions concerning coadministration of acid reducing medications (eg, H2RA or PPIs), and other antiretroviral drugs (eg, efavirenz, tenofovir, and didanosine).

<sup>b</sup> REYATAZ is not recommended for treatment-experienced pregnant patients during the second and third trimester taking REYATAZ with both tenofovir and H2RA.

## 2.6 Renal Impairment

For patients with renal impairment, including those with severe renal impairment who are not managed with hemodialysis, no dose adjustment is required for REYATAZ. Treatment-naive patients with end stage renal disease managed with hemodialysis should receive REYATAZ 300 mg with ritonavir 100 mg. REYATAZ should not be administered to HIV-treatment-experienced patients with end stage renal disease managed with hemodialysis. [See *Use in Specific Populations* (8.7).]

## 2.7 Dosage Adjustments in Patients with Hepatic Impairment

Table 5 displays the recommended REYATAZ dosage in treatment-naive patients with hepatic impairment. The use of REYATAZ in patients with severe hepatic impairment (Child-Pugh Class C) is not recommended. The coadministration of REYATAZ with ritonavir in patients with any degree of hepatic impairment is not recommended.

**Table 5: Recommended Dosage of REYATAZ Capsules in Treatment-Naive Adults with Hepatic Impairment**

	REYATAZ Once Daily Dosage
Mild hepatic impairment (Child-Pugh Class A)	400 mg
Moderate hepatic impairment (Child-Pugh Class B)	300 mg
Severe hepatic impairment (Child-Pugh Class C)	REYATAZ with or without ritonavir is not recommended

### 3 DOSAGE FORMS AND STRENGTHS

#### *REYATAZ Capsules:*

- 150 mg capsule with blue cap and powder blue body, printed with white ink “BMS 150 mg” on the cap and with blue ink “3624” on the body.
- 200 mg capsule with blue cap and blue body, printed with white ink “BMS 200 mg” on the cap and with white ink “3631” on the body.
- 300 mg capsule with red cap and blue body, printed with white ink “BMS 300 mg” on the cap and with white ink “3622” on the body.

#### *REYATAZ Oral Powder:*

- 50 mg of atazanavir as an oral powder in a packet.

### 4 CONTRAINDICATIONS

REYATAZ is contraindicated:

- in patients with previously demonstrated clinically significant hypersensitivity (eg, Stevens-Johnson syndrome, erythema multiforme, or toxic skin eruptions) to any of the components of REYATAZ capsules or REYATAZ oral powder [see *Warnings and Precautions (5.2)*].
- when coadministered with drugs that are highly dependent on CYP3A or UGT1A1 for clearance, and for which elevated plasma concentrations of the interacting drugs are associated with serious and/or life-threatening events (see Table 6).
- when coadministered with drugs that strongly induce CYP3A and may lead to lower exposure and loss of efficacy of REYATAZ (see Table 6).

Table 6 displays drugs that are contraindicated with REYATAZ.



**Table 6: Drugs that are Contraindicated with REYATAZ (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

Drug Class	Drugs within class that are contraindicated with REYATAZ	Clinical Comment
Alpha 1-Adrenoreceptor Antagonist	Alfuzosin	Potential for increased alfuzosin concentrations, which can result in hypotension.
Antimycobacterials	Rifampin	Rifampin substantially decreases plasma concentrations of atazanavir, which may result in loss of therapeutic effect and development of resistance.
Antineoplastics	Irinotecan	Atazanavir inhibits UGT1A1 and may interfere with the metabolism of irinotecan, resulting in increased irinotecan toxicities.
Benzodiazepines	Triazolam, orally administered midazolam <sup>a</sup>	Triazolam and orally administered midazolam are extensively metabolized by CYP3A4. Coadministration of triazolam or orally administered midazolam with REYATAZ may cause large increases in the concentration of these benzodiazepines. Potential for serious and/or life-threatening events such as prolonged or increased sedation or respiratory depression.
Ergot Derivatives	Dihydroergotamine, ergotamine, ergonovine, methylergonovine	Potential for serious and/or life-threatening events such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
GI Motility Agent	Cisapride	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
Herbal Products	St. John's wort ( <i>Hypericum perforatum</i> )	Coadministration of St. John's wort and REYATAZ may result in loss of therapeutic effect and development of resistance.
HMG-CoA Reductase Inhibitors	Lovastatin, simvastatin	Potential for serious reactions such as myopathy, including rhabdomyolysis.
Neuroleptic	Pimozide	Potential for serious and/or life-threatening reactions such as cardiac arrhythmias.
PDE5 Inhibitor	Sildenafil <sup>b</sup> when dosed as REVATIO <sup>®</sup> for the treatment of pulmonary arterial hypertension	Potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, priapism, and syncope).
Protease Inhibitors	Indinavir	Both REYATAZ and indinavir are associated with indirect (unconjugated) hyperbilirubinemia.
Non-nucleoside Reverse Transcriptase Inhibitors	Nevirapine	Nevirapine substantially decreases atazanavir exposure which may result in loss of therapeutic effect and development of resistance. Potential risk for nevirapine-associated adverse reactions due to increased nevirapine exposures.

<sup>a</sup> See *Drug Interactions, Table 16 (7)* for parenterally administered midazolam.

<sup>b</sup> See *Drug Interactions, Table 16 (7)* for sildenafil when dosed as VIAGRA<sup>®</sup> for erectile dysfunction.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Cardiac Conduction Abnormalities

REYATAZ has been shown to prolong the PR interval of the electrocardiogram in some patients. In healthy volunteers and in patients, abnormalities in atrioventricular (AV) conduction were asymptomatic and generally limited to first-degree AV block. There have been reports of second-degree AV block and other conduction abnormalities [*see Adverse Reactions (6.4) and Overdosage (10)*]. In clinical trials that included electrocardiograms, asymptomatic first-degree AV block was observed in 5.9% of atazanavir-treated patients (n=920), 5.2% of lopinavir/ritonavir-treated patients (n=252), 10.4% of nelfinavir-treated patients (n=48), and 3.0% of efavirenz-treated patients (n=329). In Study AI424-045, asymptomatic first-degree AV block was observed in 5% (6/118) of atazanavir/ritonavir-treated patients and 5% (6/116) of lopinavir/ritonavir-treated patients who had on-study electrocardiogram measurements. Because of limited clinical experience in patients with preexisting conduction system disease (eg, marked first-degree AV block or second- or third-degree AV block). ECG monitoring should be considered in these patients. [*See Clinical Pharmacology (12.2).*]

### 5.2 Rash

In controlled clinical trials, rash (all grades, regardless of causality) occurred in approximately 20% of patients treated with REYATAZ. The median time to onset of rash in clinical studies was 7.3 weeks and the median duration of rash was 1.4 weeks. Rashes were generally mild-to-moderate maculopapular skin eruptions. Treatment-emergent adverse reactions of moderate or severe rash (occurring at a rate of  $\geq 2\%$ ) are presented for the individual clinical studies [*see Adverse Reactions (6.1)*]. Dosing with REYATAZ was often continued without interruption in patients who developed rash. The discontinuation rate for rash in clinical trials was  $<1\%$ . Cases of Stevens-Johnson syndrome, erythema multiforme, and toxic skin eruptions, including drug rash, eosinophilia, and systemic symptoms (DRESS) syndrome, have been reported in patients receiving REYATAZ. [*See Contraindications (4) and Adverse Reactions (6.1).*] REYATAZ should be discontinued if severe rash develops.

### **5.3 Hyperbilirubinemia**

Most patients taking REYATAZ experience asymptomatic elevations in indirect (unconjugated) bilirubin related to inhibition of UDP-glucuronosyl transferase (UGT). This hyperbilirubinemia is reversible upon discontinuation of REYATAZ. Hepatic transaminase elevations that occur with hyperbilirubinemia should be evaluated for alternative etiologies. No long-term safety data are available for patients experiencing persistent elevations in total bilirubin >5 times the upper limit of normal (ULN). Alternative antiretroviral therapy to REYATAZ may be considered if jaundice or scleral icterus associated with bilirubin elevations presents cosmetic concerns for patients. Dose reduction of atazanavir is not recommended since long-term efficacy of reduced doses has not been established. [See *Adverse Reactions* (6.1, 6.2).]

### **5.4 Patients with Phenylketonuria**

Phenylalanine can be harmful to patients with phenylketonuria (PKU). REYATAZ oral powder contains phenylalanine (a component of aspartame). Each packet of REYATAZ oral powder contains 35 mg of phenylalanine. REYATAZ capsules do not contain phenylalanine.

### **5.5 Hepatotoxicity**

Patients with underlying hepatitis B or C viral infections or marked elevations in transaminases before treatment may be at increased risk for developing further transaminase elevations or hepatic decompensation. In these patients, hepatic laboratory testing should be conducted prior to initiating therapy with REYATAZ and during treatment. [See *Adverse Reactions* (6.3) and *Use in Specific Populations* (8.8).]

### **5.6 Nephrolithiasis and Cholelithiasis**

Cases of nephrolithiasis and/or cholelithiasis have been reported during postmarketing surveillance in HIV-infected patients receiving REYATAZ therapy. Some patients required hospitalization for additional management and some had complications. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis and/or cholelithiasis occur, temporary interruption or discontinuation of therapy may be considered. [See *Adverse Reactions* (6.4).]

### **5.7 Risk of Serious Adverse Reactions Due to Drug Interactions**

Initiation of REYATAZ with ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving REYATAZ with ritonavir, may increase plasma concentrations of medications

metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of REYATAZ with ritonavir, respectively. These interactions may lead to:

- clinically significant adverse reactions, potentially leading to severe, life threatening, or fatal events from greater exposures of concomitant medications.
- clinically significant adverse reactions from greater exposures of REYATAZ with ritonavir.
- loss of therapeutic effect of REYATAZ with ritonavir and possible development of resistance.

See Table 16 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations [see *Drug Interactions (7)*]. Consider the potential for drug interactions prior to and during REYATAZ/ritonavir therapy; review concomitant medications during REYATAZ/ritonavir therapy; and monitor for the adverse reactions associated with the concomitant medications [see *Contraindications (4)* and *Drug Interactions (7)*].

## 5.8 Diabetes Mellitus/Hyperglycemia

New-onset diabetes mellitus, exacerbation of preexisting diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-infected patients receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and a causal relationship between protease inhibitor therapy and these events has not been established. [See *Adverse Reactions (6.4)*.]

## 5.9 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including REYATAZ. During the initial phase of combination antiretroviral treatment, patients whose immune system responds may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jiroveci* pneumonia, or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

## 5.10 Fat Redistribution

Redistribution/accumulation of body fat including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and “cushingoid appearance” have been observed in patients receiving antiretroviral therapy. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

## 5.11 Hemophilia

There have been reports of increased bleeding, including spontaneous skin hematomas and hemarthrosis, in patients with hemophilia type A and B treated with protease inhibitors. In some patients additional factor VIII was given. In more than half of the reported cases, treatment with protease inhibitors was continued or reintroduced. A causal relationship between protease inhibitor therapy and these events has not been established.

## 5.12 Resistance/Cross-Resistance

Various degrees of cross-resistance among protease inhibitors have been observed. Resistance to atazanavir may not preclude the subsequent use of other protease inhibitors. *[See Microbiology (12.4).]*

# 6 ADVERSE REACTIONS

The following adverse reactions are discussed in greater detail in other sections of the labeling:

- cardiac conduction abnormalities *[see Warnings and Precautions (5.1)]*
- rash *[see Warnings and Precautions (5.2)]*
- hyperbilirubinemia *[see Warnings and Precautions (5.3)]*
- nephrolithiasis and cholelithiasis *[see Warnings and Precautions (5.6)]*

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

## 6.1 Clinical Trial Experience in Adults

### Treatment-Emergent Adverse Reactions in Treatment-Naive Patients

The safety profile of REYATAZ in treatment-naive adults is based on 1625 HIV-1 infected patients in clinical trials. 536 patients received REYATAZ 300 mg with ritonavir 100 mg and 1089 patients received REYATAZ 400 mg or higher (without ritonavir).

The most common adverse reactions were nausea, jaundice/scleral icterus, and rash.

Selected clinical adverse reactions of moderate or severe intensity reported in  $\geq 2\%$  of treatment-naïve patients receiving combination therapy including REYATAZ 300 mg with ritonavir 100 mg and REYATAZ 400 mg (without ritonavir) are presented in Tables 7 and 8, respectively.

**Table 7: Selected Treatment-Emergent Adverse Reactions<sup>a</sup> of Moderate or Severe Intensity Reported in  $\geq 2\%$  of Adult Treatment-Naïve Patients,<sup>b</sup> Study AI424-138**

	96 weeks <sup>c</sup> REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir with emtricitabine <sup>d</sup> (n=441)	96 weeks <sup>c</sup> lopinavir 400 mg with ritonavir 100 mg (twice daily) and tenofovir with emtricitabine <sup>d</sup> (n=437)
<b>Digestive System</b>		
Nausea	4%	8%
Jaundice/scleral icterus	5%	*
Diarrhea	2%	12%
<b>Skin and Appendages</b>		
Rash	3%	2%

\* None reported in this treatment arm.

<sup>a</sup> Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

<sup>b</sup> Based on the regimen containing REYATAZ.

<sup>c</sup> Median time on therapy.

<sup>d</sup> As a fixed-dose combination: 300 mg tenofovir, 200 mg emtricitabine once daily.

**Table 8: Selected Treatment-Emergent Adverse Reactions<sup>a</sup> of Moderate or Severe Intensity Reported in  $\geq 2\%$  of Adult Treatment-Naïve Patients,<sup>b</sup> Studies AI424-034, AI424-007, and AI424-008**

Study AI424-034		Studies AI424-007, -008	
64 weeks <sup>c</sup> REYATAZ 400 mg once daily + lamivudine + zidovudine <sup>e</sup> (n=404)	64 weeks <sup>c</sup> efavirenz 600 mg once daily + lamivudine + zidovudine <sup>e</sup> (n=401)	120 weeks <sup>c,d</sup> REYATAZ 400 mg once daily + stavudine + lamivudine or didanosine (n=279)	73 weeks <sup>c,d</sup> nelfinavir 750 mg TID or 1250 mg BID + stavudine + lamivudine or didanosine (n=191)

**Table 8: Selected Treatment-Emergent Adverse Reactions<sup>a</sup> of Moderate or Severe Intensity Reported in  $\geq 2\%$  of Adult Treatment-Naive Patients,<sup>b</sup> Studies AI424-034, AI424-007, and AI424-008**

	Study AI424-034		Studies AI424-007, -008	
	64 weeks <sup>c</sup> REYATAZ 400 mg once daily + lamivudine + zidovudine <sup>e</sup>  (n=404)	64 weeks <sup>c</sup> efavirenz 600 mg once daily + lamivudine + zidovudine <sup>e</sup>  (n=401)	120 weeks <sup>c,d</sup> REYATAZ 400 mg once daily + stavudine + lamivudine or didanosine  (n=279)	73 weeks <sup>c,d</sup> nelfinavir 750 mg TID or 1250 mg BID + stavudine + lamivudine or didanosine  (n=191)
<b>Body as a Whole</b>				
Headache	6%	6%	1%	2%
<b>Digestive System</b>				
Nausea	14%	12%	6%	4%
Jaundice/scleral icterus	7%	*	7%	*
Vomiting	4%	7%	3%	3%
Abdominal pain	4%	4%	4%	2%
Diarrhea	1%	2%	3%	16%
<b>Nervous System</b>				
Insomnia	3%	3%	<1%	*
Dizziness	2%	7%	<1%	*
Peripheral neurologic symptoms	<1%	1%	4%	3%
<b>Skin and Appendages</b>				
Rash	7%	10%	5%	1%

\* None reported in this treatment arm.

<sup>a</sup> Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

<sup>b</sup> Based on regimens containing REYATAZ.

<sup>c</sup> Median time on therapy.

<sup>d</sup> Includes long-term follow-up.

<sup>e</sup> As a fixed-dose combination: 150 mg lamivudine, 300 mg zidovudine twice daily.

### Treatment-Emergent Adverse Reactions in Treatment-Experienced Patients

The safety profile of REYATAZ in treatment-experienced adults is based on 119 HIV-1 infected patients in clinical trials.

The most common adverse reactions are jaundice/scleral icterus and myalgia.

Selected clinical adverse reactions of moderate or severe intensity reported in  $\geq 2\%$  of treatment-experienced patients receiving REYATAZ/ritonavir are presented in Table 9.

**Table 9: Selected Treatment-Emergent Adverse Reactions<sup>a</sup> of Moderate or Severe Intensity Reported in  $\geq 2\%$  of Adult Treatment-Experienced Patients,<sup>b</sup> Study AI424-045**

	48 weeks <sup>c</sup> REYATAZ/ritonavir 300/100 mg once daily + tenofovir + NRTI (n=119)	48 weeks <sup>c</sup> lopinavir/ritonavir 400/100 mg twice daily <sup>d</sup> + tenofovir + NRTI (n=118)
<b>Body as a Whole</b>		
Fever	2%	*
<b>Digestive System</b>		
Jaundice/scleral icterus	9%	*
Diarrhea	3%	11%
Nausea	3%	2%
<b>Nervous System</b>		
Depression	2%	<1%
<b>Musculoskeletal System</b>		
Myalgia	4%	*

\* None reported in this treatment arm.

<sup>a</sup> Includes events of possible, probable, certain, or unknown relationship to treatment regimen.

<sup>b</sup> Based on the regimen containing REYATAZ.

<sup>c</sup> Median time on therapy.

<sup>d</sup> As a fixed-dose combination.

### Laboratory Abnormalities in Treatment-Naive Patients

The percentages of adult treatment-naive patients treated with combination therapy including REYATAZ 300 mg with ritonavir 100 mg and REYATAZ 400 mg (without ritonavir) with Grade 3–4 laboratory abnormalities are presented in Tables 10 and 11, respectively.



**Table 10: Grade 3–4 Laboratory Abnormalities Reported in ≥2% of Adult Treatment-Naive Patients,<sup>a</sup> Study AI424-138**

Variable	Limit <sup>c</sup>	96 weeks <sup>b</sup> REYATAZ 300 mg with ritonavir 100 mg (once daily) and tenofovir with emtricitabine <sup>d</sup> (n=441)	96 weeks <sup>b</sup> lopinavir 400 mg with ritonavir 100 mg (twice daily) and tenofovir with emtricitabine <sup>d</sup> (n=437)
Chemistry	<u>High</u>		
SGOT/AST	≥5.1 × ULN	3%	1%
SGPT/ALT	≥5.1 × ULN	3%	2%
Total Bilirubin	≥2.6 × ULN	44%	<1%
Lipase	≥2.1 × ULN	2%	2%
Creatine Kinase	≥5.1 × ULN	8%	7%
Total Cholesterol	≥240 mg/dL	11%	25%
Hematology	<u>Low</u>		
Neutrophils	<750 cells/mm <sup>3</sup>	5%	2%

<sup>a</sup> Based on the regimen containing REYATAZ.

<sup>b</sup> Median time on therapy.

<sup>c</sup> ULN = upper limit of normal.

<sup>d</sup> As a fixed-dose combination: 300 mg tenofovir, 200 mg emtricitabine once daily.

**Table 11: Grade 3–4 Laboratory Abnormalities Reported in ≥2% of Adult Treatment-Naive Patients,<sup>a</sup> Studies AI424-034, AI424-007, and AI424-008**

Variable	Limit <sup>d</sup>	Study AI424-034		Studies AI424-007, -008	
		64 weeks <sup>b</sup> REYATAZ 400 mg once daily + lamivudine + zidovudine <sup>e</sup> (n=404)	64 weeks <sup>b</sup> efavirenz 600 mg once daily + lamivudine + zidovudine <sup>e</sup> (n=401)	120 weeks <sup>b,c</sup> REYATAZ 400 mg once daily + stavudine + lamivudine or + stavudine + didanosine (n=279)	73 weeks <sup>b,c</sup> nelfinavir 750 mg TID or 1250 mg BID + stavudine + lamivudine or + stavudine + didanosine (n=191)
Chemistry	<u>High</u>				
SGOT/AST	≥5.1 × ULN	2%	2%	7%	5%
SGPT/ALT	≥5.1 × ULN	4%	3%	9%	7%
Total Bilirubin	≥2.6 × ULN	35%	<1%	47%	3%
Amylase	≥2.1 × ULN	*	*	14%	10%
Lipase	≥2.1 × ULN	<1%	1%	4%	5%
Creatine Kinase	≥5.1 × ULN	6%	6%	11%	9%
Total Cholesterol	≥240 mg/dL	6%	24%	19%	48%
Triglycerides	≥751 mg/dL	<1%	3%	4%	2%
Hematology	<u>Low</u>				
Hemoglobin	<8.0 g/dL	5%	3%	<1%	4%
Neutrophils	<750 cells/mm <sup>3</sup>	7%	9%	3%	7%

\* None reported in this treatment arm.

<sup>a</sup> Based on regimen(s) containing REYATAZ.

<sup>b</sup> Median time on therapy.

<sup>c</sup> Includes long-term follow-up.

<sup>d</sup> ULN = upper limit of normal.

<sup>e</sup> As a fixed-dose combination: 150 mg lamivudine, 300 mg zidovudine twice daily.

## Laboratory Abnormalities in Treatment-Experienced Patients

The percentages of adult treatment-experienced patients treated with combination therapy including REYATAZ/ritonavir with Grade 3–4 laboratory abnormalities are presented in Table 12.

**Table 12: Grade 3–4 Laboratory Abnormalities Reported in  $\geq 2\%$  of Adult Treatment-Experienced Patients, Study AI424-045<sup>a</sup>**

Variable	Limit <sup>c</sup>	48 weeks <sup>b</sup> REYATAZ/ritonavir 300/100 mg once daily + tenofovir + NRTI (n=119)	48 weeks <sup>b</sup> lopinavir/ritonavir 400/100 mg twice daily <sup>d</sup> + tenofovir + NRTI (n=118)
Chemistry	<u>High</u>		
SGOT/AST	$\geq 5.1 \times \text{ULN}$	3%	3%
SGPT/ALT	$\geq 5.1 \times \text{ULN}$	4%	3%
Total Bilirubin	$\geq 2.6 \times \text{ULN}$	49%	<1%
Lipase	$\geq 2.1 \times \text{ULN}$	5%	6%
Creatine Kinase	$\geq 5.1 \times \text{ULN}$	8%	8%
Total Cholesterol	$\geq 240 \text{ mg/dL}$	25%	26%
Triglycerides	$\geq 751 \text{ mg/dL}$	8%	12%
Glucose	$\geq 251 \text{ mg/dL}$	5%	<1%
Hematology	<u>Low</u>		
Platelets	$< 50,000 \text{ cells/mm}^3$	2%	3%
Neutrophils	$< 750 \text{ cells/mm}^3$	7%	8%

<sup>a</sup> Based on regimen(s) containing REYATAZ.

<sup>b</sup> Median time on therapy.

<sup>c</sup> ULN = upper limit of normal.

<sup>d</sup> As a fixed-dose combination.

## Lipids, Change from Baseline in Treatment-Naive Patients

For Study AI424-138 and Study AI424-034, changes from baseline in LDL-cholesterol, HDL-cholesterol, total cholesterol, and triglycerides are shown in Tables 13 and 14, respectively.

**Table 13: Lipid Values, Mean Change from Baseline, Study AI424-138**

	REYATAZ/ritonavir <sup>a,b</sup>					lopinavir/ritonavir <sup>b,c</sup>				
	Baseline mg/dL (n=428 <sup>e</sup> )	Week 48 mg/dL (n=372 <sup>e</sup> )	Change <sup>d</sup> (n=372 <sup>e</sup> )	Week 96 mg/dL (n=342 <sup>e</sup> )	Change <sup>d</sup> (n=342 <sup>e</sup> )	Baseline mg/dL (n=424 <sup>e</sup> )	Week 48 mg/dL (n=335 <sup>e</sup> )	Change <sup>d</sup> (n=335 <sup>e</sup> )	Week 96 mg/dL (n=291 <sup>e</sup> )	Change <sup>d</sup> (n=291 <sup>e</sup> )
LDL-Cholesterol <sup>f</sup>	92	105	+14%	105	+14%	93	111	+19%	110	+17%
HDL-Cholesterol <sup>f</sup>	37	46	+29%	44	+21%	36	48	+37%	46	+29%
Total Cholesterol <sup>f</sup>	149	169	+13%	169	+13%	150	187	+25%	186	+25%
Triglycerides <sup>f</sup>	126	145	+15%	140	+13%	129	194	+52%	184	+50%

<sup>a</sup> REYATAZ 300 mg with ritonavir 100 mg once daily with the fixed-dose combination: 300 mg tenofovir, 200 mg emtricitabine once daily.

<sup>b</sup> Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 1% in the lopinavir/ritonavir treatment arm and 1% in the REYATAZ/ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 8% in the lopinavir/ritonavir treatment arm and 2% in the REYATAZ/ritonavir arm. Through Week 96, serum lipid-reducing agents were used in 10% in the lopinavir/ritonavir treatment arm and 3% in the REYATAZ/ritonavir arm.

<sup>c</sup> Lopinavir 400 mg with ritonavir 100 mg twice daily with the fixed-dose combination 300 mg tenofovir, 200 mg emtricitabine once daily.

<sup>d</sup> The change from baseline is the mean of within-patient changes from baseline for patients with both baseline and Week 48 or Week 96 values and is not a simple difference of the baseline and Week 48 or Week 96 mean values, respectively.

<sup>e</sup> Number of patients with LDL-cholesterol measured.

<sup>f</sup> Fasting.

**Table 14: Lipid Values, Mean Change from Baseline, Study AI424-034**

	REYATAZ <sup>a,b</sup>			efavirenz <sup>b,c</sup>		
	Baseline mg/dL (n=383 <sup>e</sup> )	Week 48 mg/dL (n=283 <sup>e</sup> )	Week 48 Change <sup>d</sup> (n=272 <sup>e</sup> )	Baseline mg/dL (n=378 <sup>e</sup> )	Week 48 mg/dL (n=264 <sup>e</sup> )	Week 48 Change <sup>d</sup> (n=253 <sup>e</sup> )
LDL-Cholesterol <sup>f</sup>	98	98	+1%	98	114	+18%
HDL-Cholesterol	39	43	+13%	38	46	+24%
Total Cholesterol	164	168	+2%	162	195	+21%
Triglycerides <sup>f</sup>	138	124	-9%	129	168	+23%

<sup>a</sup> REYATAZ 400 mg once daily with the fixed-dose combination: 150 mg lamivudine, 300 mg zidovudine twice daily.

<sup>b</sup> Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 0% in the efavirenz treatment arm and <1% in the REYATAZ arm. Through Week 48, serum lipid-reducing agents were used in 3% in the efavirenz treatment arm and 1% in the REYATAZ arm.

<sup>c</sup> Efavirenz 600 mg once daily with the fixed-dose combination: 150 mg lamivudine, 300 mg zidovudine twice daily.

<sup>d</sup> The change from baseline is the mean of within-patient changes from baseline for patients with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.

- <sup>e</sup> Number of patients with LDL-cholesterol measured.  
<sup>f</sup> Fasting.

## Lipids, Change from Baseline in Treatment-Experienced Patients

For Study AI424-045, changes from baseline in LDL-cholesterol, HDL-cholesterol, total cholesterol, and triglycerides are shown in Table 15. The observed magnitude of dyslipidemia was less with REYATAZ/ritonavir than with lopinavir/ritonavir. However, the clinical impact of such findings has not been demonstrated.

**Table 15: Lipid Values, Mean Change from Baseline, Study AI424-045**

	REYATAZ/ritonavir <sup>a,b</sup>			lopinavir/ritonavir <sup>b,c</sup>		
	Baseline mg/dL (n=111 <sup>e</sup> )	Week 48 mg/dL (n=75 <sup>e</sup> )	Week 48 Change <sup>d</sup> (n=74 <sup>e</sup> )	Baseline mg/dL (n=108 <sup>e</sup> )	Week 48 mg/dL (n=76 <sup>e</sup> )	Week 48 Change <sup>d</sup> (n=73 <sup>e</sup> )
LDL-Cholesterol <sup>f</sup>	108	98	−10%	104	103	+1%
HDL-Cholesterol	40	39	−7%	39	41	+2%
Total Cholesterol	188	170	−8%	181	187	+6%
Triglycerides <sup>f</sup>	215	161	−4%	196	224	+30%

<sup>a</sup> REYATAZ 300 mg once daily + ritonavir + tenofovir + 1 NRTI.

<sup>b</sup> Values obtained after initiation of serum lipid-reducing agents were not included in these analyses. At baseline, serum lipid-reducing agents were used in 4% in the lopinavir/ritonavir treatment arm and 4% in the REYATAZ/ritonavir arm. Through Week 48, serum lipid-reducing agents were used in 19% in the lopinavir/ritonavir treatment arm and 8% in the REYATAZ/ritonavir arm.

<sup>c</sup> Lopinavir/ritonavir (400/100 mg) BID + tenofovir + 1 NRTI.

<sup>d</sup> The change from baseline is the mean of within-patient changes from baseline for patients with both baseline and Week 48 values and is not a simple difference of the baseline and Week 48 mean values.

<sup>e</sup> Number of patients with LDL-cholesterol measured.

<sup>f</sup> Fasting.

## 6.2 Clinical Trial Experience in Pediatric Patients

### Adverse Reactions in Pediatric Patients: REYATAZ Capsules

The safety and tolerability of REYATAZ Capsules with and without ritonavir have been established in pediatric patients at least 6 years of age from the open-label, multicenter clinical trial PACTG 1020A.

The safety profile of REYATAZ in pediatric patients (6 to less than 18 years of age) taking the capsule formulation was generally similar to that observed in clinical studies of REYATAZ in

adults. The most common Grade 2–4 adverse events ( $\geq 5\%$ , regardless of causality) reported in pediatric patients were cough (21%), fever (18%), jaundice/scleral icterus (15%), rash (14%), vomiting (12%), diarrhea (9%), headache (8%), peripheral edema (7%), extremity pain (6%), nasal congestion (6%), oropharyngeal pain (6%), wheezing (6%), and rhinorrhea (6%). Asymptomatic second-degree atrioventricular block was reported in  $<2\%$  of patients. The most common Grade 3–4 laboratory abnormalities occurring in pediatric patients taking the capsule formulation were elevation of total bilirubin ( $\geq 3.2$  mg/dL, 58%), neutropenia (9%), and hypoglycemia (4%). All other Grade 3–4 laboratory abnormalities occurred with a frequency of less than 3%.

### **Adverse Reactions in Pediatric Patients: REYATAZ Oral Powder**

The data described below reflect exposure to REYATAZ oral powder in 89 subjects weighing from 10 kg to less than 25 kg, including 65 patients exposed for 48 weeks. These data are from two pooled open-label, multi-center clinical trials in treatment-naïve and treatment-experienced pediatric patients (AI424-397 [PRINCE I] and AI424-451 [PRINCE II]). Age ranged from 15 months to less than 7.5 years of age. In these studies 53% were female and 47% were male. All patients received ritonavir and 2 nucleoside reverse transcriptase inhibitors (NRTIs).

The safety profile of REYATAZ in pediatric patients taking REYATAZ oral powder was generally similar to that observed in clinical studies of REYATAZ in pediatric patients taking REYATAZ capsules. The most common Grade 3–4 laboratory abnormalities occurring in pediatric patients weighing 10 kg to less than 25 kg taking REYATAZ oral powder were increased amylase (19%), neutropenia (12%), increased SGPT/ALT (5%), elevation of total bilirubin ( $\geq 2.6$  times ULN, 12%), increased lipase (5%), and decreased hemoglobin (3%). All other Grade 3–4 laboratory abnormalities occurred with a frequency of less than 3%.

### **6.3 Patients Co-Infected with Hepatitis B and/or Hepatitis C Virus**

In study AI424-138, 60 patients treated with REYATAZ/ritonavir 300 mg/100 mg once daily, and 51 patients treated with lopinavir/ritonavir 400 mg/100 mg twice daily, each with fixed dose tenofovir-emtricitabine, were seropositive for hepatitis B and/or C at study entry. ALT levels  $>5$  times ULN developed in 10% (6/60) of the REYATAZ/ritonavir-treated patients and 8% (4/50) of the lopinavir/ritonavir-treated patients. AST levels  $>5$  times ULN developed in 10% (6/60) of the REYATAZ/ritonavir-treated patients and none (0/50) of the lopinavir/ritonavir-treated patients.

In study AI424-045, 20 patients treated with REYATAZ/ritonavir 300 mg/100 mg once daily, and 18 patients treated with lopinavir/ritonavir 400 mg/100 mg twice daily, were seropositive for

hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 25% (5/20) of the REYATAZ/ritonavir-treated patients and 6% (1/18) of the lopinavir/ritonavir-treated patients. AST levels >5 times ULN developed in 10% (2/20) of the REYATAZ/ritonavir-treated patients and 6% (1/18) of the lopinavir/ritonavir-treated patients.

In studies AI424-008 and AI424-034, 74 patients treated with 400 mg of REYATAZ once daily, 58 who received efavirenz, and 12 who received nelfinavir were seropositive for hepatitis B and/or C at study entry. ALT levels >5 times ULN developed in 15% of the REYATAZ-treated patients, 14% of the efavirenz-treated patients, and 17% of the nelfinavir-treated patients. AST levels >5 times ULN developed in 9% of the REYATAZ-treated patients, 5% of the efavirenz-treated patients, and 17% of the nelfinavir-treated patients. Within REYATAZ and control regimens, no difference in frequency of bilirubin elevations was noted between seropositive and seronegative patients. *[See Warnings and Precautions (5.3).]*

## 6.4 Postmarketing Experience

The following events have been identified during postmarketing use of REYATAZ. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

*Body as a Whole:* edema

*Cardiovascular System:* second-degree AV block, third-degree AV block, left bundle branch block, QTc prolongation *[see Warnings and Precautions (5.1)]*

*Gastrointestinal System:* pancreatitis

*Hepatic System:* hepatic function abnormalities

*Hepatobiliary Disorders:* cholelithiasis *[see Warnings and Precautions (5.5)]*, cholecystitis, cholestasis

*Metabolic System and Nutrition Disorders:* diabetes mellitus, hyperglycemia *[see Warnings and Precautions (5.8)]*

*Musculoskeletal System:* arthralgia

*Renal System:* nephrolithiasis *[see Warnings and Precautions (5.6)]*, interstitial nephritis

*Skin and Appendages:* alopecia, maculopapular rash *[see Contraindications (4) and Warnings and Precautions (5.2)]*, pruritus, angioedema

## 7 DRUG INTERACTIONS

See also *Contraindications* (4) and *Clinical Pharmacology* (12.3).

### 7.1 Potential for REYATAZ to Affect Other Drugs

Atazanavir is an inhibitor of CYP3A and UGT1A1. Coadministration of REYATAZ and drugs primarily metabolized by CYP3A or UGT1A1 may result in increased plasma concentrations of the other drug that could increase or prolong its therapeutic and adverse effects.

Atazanavir is a weak inhibitor of CYP2C8. Use of REYATAZ without ritonavir is not recommended when coadministered with drugs highly dependent on CYP2C8 with narrow therapeutic indices (eg, paclitaxel, repaglinide). When REYATAZ with ritonavir is coadministered with substrates of CYP2C8, clinically significant interactions are not expected. [See *Clinical Pharmacology*, Table 22 (12.3).]

The magnitude of CYP3A-mediated drug interactions on coadministered drug may change when REYATAZ is coadministered with ritonavir. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.

### 7.2 Potential for Other Drugs to Affect REYATAZ

Atazanavir is a CYP3A4 substrate; therefore, drugs that induce CYP3A4 may decrease atazanavir plasma concentrations and reduce REYATAZ's therapeutic effect.

Atazanavir solubility decreases as pH increases. Reduced plasma concentrations of atazanavir are expected if proton-pump inhibitors, antacids, buffered medications, or H<sub>2</sub>-receptor antagonists are administered with REYATAZ [see *Dosage and Administration* (2.2, 2.3, 2.4, and 2.5)].

### 7.3 Established and Other Potentially Significant Drug Interactions

Table 16 provides dosing recommendations as a result of drug interactions with REYATAZ. These recommendations are based on either drug interaction studies or predicted interactions due to the expected magnitude of interaction and potential for serious events or loss of efficacy.



**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<b>Concomitant Drug Class: Specific Drugs</b>	<b>Effect on Concentration of Atazanavir or Concomitant Drug</b>	<b>Clinical Comment</b>
<b><i>HIV Antiviral Agents</i></b>		
<i>Nucleoside Reverse Transcriptase Inhibitors (NRTIs):</i> didanosine buffered formulations enteric-coated (EC) capsules	↓ atazanavir ↓ didanosine	Coadministration of REYATAZ with didanosine buffered tablets resulted in a marked decrease in atazanavir exposure. It is recommended that REYATAZ be given (with food) 2 h before or 1 h after didanosine buffered formulations. Simultaneous administration of didanosine EC and REYATAZ with food results in a decrease in didanosine exposure. Thus, REYATAZ and didanosine EC should be administered at different times.
<i>Nucleotide Reverse Transcriptase Inhibitors:</i> tenofovir disoproxil fumarate	↓ atazanavir ↑ tenofovir	Tenofovir may decrease the AUC and C <sub>min</sub> of atazanavir. When coadministered with tenofovir, it is recommended that REYATAZ 300 mg be given with ritonavir 100 mg and tenofovir 300 mg (all as a single daily dose with food). REYATAZ increases tenofovir concentrations. The mechanism of this interaction is unknown. Higher tenofovir concentrations could potentiate tenofovir-associated adverse reactions, including renal disorders. Patients receiving REYATAZ and tenofovir should be monitored for tenofovir-associated adverse reactions. For pregnant women taking REYATAZ with ritonavir and tenofovir, see <i>Dosage and Administration</i> (2.5).
<i>Non-nucleoside Reverse Transcriptase Inhibitors (NNRTIs):</i> efavirenz	↓ atazanavir	Efavirenz decreases atazanavir exposure.  <b><i>In treatment-naïve patients:</i></b> If REYATAZ is combined with efavirenz, REYATAZ 400 mg (two 200-mg capsules) should be administered with ritonavir 100 mg simultaneously once daily with food, and efavirenz 600 mg should be administered once daily on an empty stomach, preferably at bedtime.  <b><i>In treatment-experienced patients:</i></b> Coadministration of REYATAZ with efavirenz in treatment-experienced patients is not recommended due to decreased atazanavir exposure.
<i>Protease Inhibitors:</i> saquinavir (soft gelatin capsules)	↑ saquinavir	Appropriate dosing recommendations for this combination, with or without ritonavir, with respect to efficacy and safety have not been established. In a clinical study, saquinavir 1200 mg coadministered with REYATAZ 400 mg and tenofovir 300 mg (all given once daily) plus nucleoside analogue reverse transcriptase inhibitors did not provide adequate efficacy [see <i>Clinical Studies</i> (14.2)].
ritonavir	↑ atazanavir	If REYATAZ is coadministered with ritonavir, it is recommended that REYATAZ 300 mg once daily be given with ritonavir 100 mg once daily with food. See the complete prescribing information for ritonavir for information on drug interactions with ritonavir.
others	↑ other protease inhibitor	Although not studied, the coadministration of REYATAZ/ritonavir and an additional protease inhibitor would be expected to increase exposure to the other protease inhibitor. Such coadministration is not recommended.
<b><i>HCV Antiviral Agents</i></b>		
<i>Protease Inhibitors:</i> boceprevir	↓ atazanavir ↓ ritonavir	Concomitant administration of boceprevir and atazanavir/ritonavir resulted in reduced steady-state exposures to atazanavir and ritonavir. Coadministration of REYATAZ/ritonavir and boceprevir is not recommended.
telaprevir	↓ telaprevir ↑ atazanavir	Concomitant administration of telaprevir and atazanavir/ritonavir resulted in reduced steady-state telaprevir exposure, while steady-state atazanavir exposure was increased.

**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<i>Concomitant Drug Class: Specific Drugs</i>	<i>Effect on Concentration of Atazanavir or Concomitant Drug</i>	<i>Clinical Comment</i>
<b>Other Agents</b>		
<i>Antacids and buffered medications</i>	↓ atazanavir	Reduced plasma concentrations of atazanavir are expected if antacids, including buffered medications, are administered with REYATAZ. REYATAZ should be administered 2 hours before or 1 hour after these medications.
<i>Antiarrhythmics:</i> amiodarone, bepridil, lidocaine (systemic), quinidine	↑ amiodarone, bepridil, lidocaine (systemic), quinidine	Coadministration with REYATAZ has the potential to produce serious and/or life-threatening adverse events and has not been studied. Caution is warranted and therapeutic concentration monitoring of these drugs is recommended if they are used concomitantly with REYATAZ.
<i>Anticoagulants:</i> warfarin	↑ warfarin	Coadministration with REYATAZ has the potential to produce serious and/or life-threatening bleeding and has not been studied. It is recommended that International Normalized Ratio (INR) be monitored.
<i>Antidepressants:</i> tricyclic antidepressants	↑ tricyclic antidepressants	Coadministration with REYATAZ has the potential to produce serious and/or life-threatening adverse events and has not been studied. Concentration monitoring of these drugs is recommended if they are used concomitantly with REYATAZ.
trazodone	↑ trazodone	Concomitant use of trazodone and REYATAZ with or without ritonavir may increase plasma concentrations of trazodone. Nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone and ritonavir. If trazodone is used with a CYP3A4 inhibitor such as REYATAZ, the combination should be used with caution and a lower dose of trazodone should be considered.
<i>Antiepileptics:</i> carbamazepine	↓ atazanavir ↑ carbamazepine	Plasma concentrations of atazanavir may be decreased when carbamazepine is administered with REYATAZ without ritonavir. Coadministration of carbamazepine and REYATAZ without ritonavir is not recommended. Ritonavir may increase plasma levels of carbamazepine. If patients beginning treatment with REYATAZ/ritonavir have been titrated to a stable dose of carbamazepine, a dose reduction for carbamazepine may be necessary.
phenytoin, phenobarbital	↓ atazanavir ↓ phenytoin ↓ phenobarbital	Plasma concentrations of atazanavir may be decreased when phenytoin or phenobarbital is administered with REYATAZ without ritonavir. Coadministration of phenytoin or phenobarbital and REYATAZ without ritonavir is not recommended. Ritonavir may decrease plasma levels of phenytoin and phenobarbital. When REYATAZ with ritonavir is coadministered with either phenytoin or phenobarbital, a dose adjustment of phenytoin or phenobarbital may be required.
lamotrigine	↓ lamotrigine	Coadministration of lamotrigine and REYATAZ <b>with</b> ritonavir may decrease lamotrigine plasma concentrations. Dose adjustment of lamotrigine may be required when coadministered with REYATAZ and ritonavir. Coadministration of lamotrigine and REYATAZ <b>without</b> ritonavir is not expected to decrease lamotrigine plasma concentrations. No dose adjustment of lamotrigine is required when coadministered with REYATAZ without ritonavir.
<i>Antifungals:</i> ketoconazole, itraconazole	<b>REYATAZ/ritonavir:</b> ↑ ketoconazole ↑ itraconazole	Coadministration of ketoconazole has only been studied with REYATAZ without ritonavir (negligible increase in atazanavir AUC and C <sub>max</sub> ). Due to the effect of ritonavir on ketoconazole, high doses of ketoconazole and itraconazole (>200 mg/day) should be used cautiously with REYATAZ/ritonavir.

**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<b>Concomitant Drug Class: Specific Drugs</b>	<b>Effect on Concentration of Atazanavir or Concomitant Drug</b>	<b>Clinical Comment</b>
voriconazole	<p><b><i>REYATAZ/ritonavir in subjects with a functional CYP2C19 allele:</i></b></p> <p>↓ voriconazole ↓ atazanavir</p> <p><b><i>REYATAZ/ritonavir in subjects without a functional CYP2C19 allele:</i></b></p> <p>↑ voriconazole ↓ atazanavir</p>	<p>The use of voriconazole in patients receiving REYATAZ/ritonavir is not recommended unless an assessment of the benefit/risk to the patient justifies the use of voriconazole. Patients should be carefully monitored for voriconazole-associated adverse reactions and loss of either voriconazole or atazanavir efficacy during the coadministration of voriconazole and REYATAZ/ritonavir. Coadministration of voriconazole with REYATAZ (without ritonavir) may affect atazanavir concentrations; however, no data are available.</p>
Antigout: colchicine	↑ colchicine	<p>The coadministration of REYATAZ with colchicine in patients with renal or hepatic impairment is not recommended.</p> <p><b><i>Recommended dosage of colchicine when administered with REYATAZ:</i></b></p> <p><b><i>Treatment of gout flares:</i></b></p> <p>0.6 mg (1 tablet) for 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Not to be repeated before 3 days.</p> <p><b><i>Prophylaxis of gout flares:</i></b></p> <p>If the original regimen was 0.6 mg <i>twice</i> a day, the regimen should be adjusted to 0.3 mg <i>once</i> a day.</p> <p>If the original regimen was 0.6 mg <i>once</i> a day, the regimen should be adjusted to 0.3 mg <i>once every other day</i>.</p> <p><b><i>Treatment of familial Mediterranean fever (FMF):</i></b></p> <p>Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day).</p>
Antimycobacterials: rifabutin	↑ rifabutin	<p>A rifabutin dose reduction of up to 75% (eg, 150 mg every other day or 3 times per week) is recommended. Increased monitoring for rifabutin-associated adverse reactions including neutropenia is warranted.</p>
Antipsychotics: quetiapine	↑ quetiapine	<p><b><i>Initiation of REYATAZ with ritonavir in patients taking quetiapine:</i></b></p> <p>Consider alternative antiretroviral therapy to avoid increases in quetiapine exposures. If coadministration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring.</p> <p><b><i>Initiation of quetiapine in patients taking REYATAZ with ritonavir:</i></b></p> <p>Refer to the quetiapine prescribing information for initial dosing and titration of quetiapine.</p>

**Table 16:**                      **Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<i>Concomitant Drug Class: Specific Drugs</i>	<i>Effect on Concentration of Atazanavir or Concomitant Drug</i>	<i>Clinical Comment</i>
<i>Benzodiazepines:</i> parenterally administered midazolam <sup>b</sup>	↑ midazolam	Concomitant use of parenteral midazolam with REYATAZ may increase plasma concentrations of midazolam. Coadministration should be done in a setting which ensures close clinical monitoring and appropriate medical management in case of respiratory depression and/or prolonged sedation. Dosage reduction for midazolam should be considered, especially if more than a single dose of midazolam is administered. Coadministration of oral midazolam with REYATAZ is CONTRAINDICATED.
<i>Calcium channel blockers:</i> diltiazem	↑ diltiazem and desacetyl-diltiazem	Caution is warranted. A dose reduction of diltiazem by 50% should be considered. ECG monitoring is recommended. Coadministration of REYATAZ/ritonavir with diltiazem has not been studied.
felodipine, nifedipine, nicardipine, and verapamil	↑ calcium channel blocker	Caution is warranted. Dose titration of the calcium channel blocker should be considered. ECG monitoring is recommended.
<i>Endothelin receptor antagonists:</i> bosentan	↓ atazanavir ↑ bosentan	Plasma concentrations of atazanavir may be decreased when bosentan is administered with REYATAZ without ritonavir. Coadministration of bosentan and REYATAZ without ritonavir is not recommended.  <b>Coadministration of bosentan in patients on REYATAZ/ritonavir:</b> For patients who have been receiving REYATAZ/ritonavir for at least 10 days, start bosentan at 62.5 mg once daily or every other day based on individual tolerability.  <b>Coadministration of REYATAZ/ritonavir in patients on bosentan:</b> Discontinue bosentan at least 36 hours before starting REYATAZ/ritonavir. At least 10 days after starting REYATAZ/ritonavir, resume bosentan at 62.5 mg once daily or every other day based on individual tolerability.
<i>HMG-CoA reductase inhibitors:</i> atorvastatin, rosuvastatin	↑ atorvastatin ↑ rosuvastatin	Titrate atorvastatin dose carefully and use the lowest necessary dose. Rosuvastatin dose should not exceed 10 mg/day. The risk of myopathy, including rhabdomyolysis, may be increased when HIV protease inhibitors, including REYATAZ, are used in combination with these drugs.

**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<i>Concomitant Drug Class: Specific Drugs</i>	<i>Effect on Concentration of Atazanavir or Concomitant Drug</i>	<i>Clinical Comment</i>
<i>H<sub>2</sub>-Receptor antagonists</i>	↓ atazanavir	<p>Plasma concentrations of atazanavir were substantially decreased when REYATAZ 400 mg once daily was administered simultaneously with famotidine 40 mg twice daily, which may result in loss of therapeutic effect and development of resistance.</p> <p><b><i>In treatment-naïve patients:</i></b></p> <p>REYATAZ 300 mg with ritonavir 100 mg once daily with food should be administered simultaneously with, and/or at least 10 hours after, a dose of the H<sub>2</sub>-receptor antagonist (H2RA). An H2RA dose comparable to famotidine 20 mg once daily up to a dose comparable to famotidine 40 mg twice daily can be used with REYATAZ 300 mg with ritonavir 100 mg in treatment-naïve patients.</p> <p style="text-align: center;">OR</p> <p>For patients unable to tolerate ritonavir, REYATAZ 400 mg once daily with food should be administered at least 2 hours before and at least 10 hours after a dose of the H2RA. No single dose of the H2RA should exceed a dose comparable to famotidine 20 mg, and the total daily dose should not exceed a dose comparable to famotidine 40 mg. The use of REYATAZ without ritonavir in pregnant women is not recommended.</p> <p><b><i>In treatment-experienced patients:</i></b></p> <p>Whenever an H2RA is given to a patient receiving REYATAZ with ritonavir, the H2RA dose should not exceed a dose comparable to famotidine 20 mg twice daily, and the REYATAZ and ritonavir doses should be administered simultaneously with, and/or at least 10 hours after, the dose of the H2RA.</p> <ul style="list-style-type: none"> <li>• REYATAZ 300 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with an H2RA.</li> <li>• REYATAZ 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with both tenofovir and an H2RA.</li> <li>• REYATAZ 400 mg with ritonavir 100 mg once daily (all as a single dose with food) if taken with either tenofovir or an H2RA for pregnant women during the second and third trimester. REYATAZ is not recommended for pregnant women during the second and third trimester taking REYATAZ with both tenofovir and an H2RA.</li> </ul>
<i>Hormonal contraceptives: ethinyl estradiol and norgestimate or norethindrone</i>	↓ ethinyl estradiol ↑ norgestimate <sup>c</sup>	<p>Use with caution if coadministration of REYATAZ or REYATAZ/ritonavir with oral contraceptives is considered. If an oral contraceptive is administered with REYATAZ plus ritonavir, it is recommended that the oral contraceptive contain at least 35 mcg of ethinyl estradiol. If REYATAZ is administered without ritonavir, the oral contraceptive should contain no more than 30 mcg of ethinyl estradiol.</p>
	↑ ethinyl estradiol ↑ norethindrone <sup>d</sup>	<p>Potential safety risks include substantial increases in progesterone exposure. The long-term effects of increases in concentration of the progestational agent are unknown and could increase the risk of insulin resistance, dyslipidemia, and acne.</p> <p>Coadministration of REYATAZ or REYATAZ/ritonavir with other hormonal contraceptives (eg, contraceptive patch, contraceptive vaginal ring, or injectable contraceptives) or oral contraceptives containing progestogens other than norethindrone or norgestimate, or less than 25 mcg of ethinyl estradiol, has not been studied; therefore, alternative methods of contraception are recommended.</p>

**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<b>Concomitant Drug Class: Specific Drugs</b>	<b>Effect on Concentration of Atazanavir or Concomitant Drug</b>	<b>Clinical Comment</b>
<i>Immunosuppressants:</i> cyclosporine, sirolimus, tacrolimus	↑ immunosuppressants	Therapeutic concentration monitoring is recommended for these immunosuppressants when coadministered with REYATAZ.
<i>Inhaled beta agonist:</i> salmeterol	↑ salmeterol	Coadministration of salmeterol with REYATAZ is not recommended. Concomitant use of salmeterol and REYATAZ may result in increased risk of cardiovascular adverse reactions associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.
<i>Inhaled/nasal steroid:</i> fluticasone	<b>REYATAZ</b> ↑ fluticasone	Concomitant use of fluticasone propionate and REYATAZ (without ritonavir) may increase plasma concentrations of fluticasone propionate. Use with caution. Consider alternatives to fluticasone propionate, particularly for long-term use.
	<b>REYATAZ/ritonavir</b> ↑ fluticasone	Concomitant use of fluticasone propionate and REYATAZ/ritonavir may increase plasma concentrations of fluticasone propionate, resulting in significantly reduced serum cortisol concentrations. Systemic corticosteroid effects, including Cushing's syndrome and adrenal suppression, have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone propionate. Coadministration of fluticasone propionate and REYATAZ/ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects [see <i>Warnings and Precautions</i> (5.1)].
<i>Macrolide antibiotics:</i> clarithromycin	↑ clarithromycin ↓ 14-OH clarithromycin ↑ atazanavir	Increased concentrations of clarithromycin may cause QTc prolongations; therefore, a dose reduction of clarithromycin by 50% should be considered when it is coadministered with REYATAZ. In addition, concentrations of the active metabolite 14-OH clarithromycin are significantly reduced; consider alternative therapy for indications other than infections due to <i>Mycobacterium avium</i> complex. Coadministration of REYATAZ/ritonavir with clarithromycin has not been studied.
<i>Opioids:</i> Buprenorphine	↑ buprenorphine ↑ norbuprenorphine	Coadministration of buprenorphine and REYATAZ with or without ritonavir increases the plasma concentration of buprenorphine and norbuprenorphine. Coadministration of REYATAZ plus ritonavir with buprenorphine warrants clinical monitoring for sedation and cognitive effects. A dose reduction of buprenorphine may be considered. Coadministration of buprenorphine and REYATAZ with ritonavir is not expected to decrease atazanavir plasma concentrations. Coadministration of buprenorphine and REYATAZ without ritonavir may decrease atazanavir plasma concentrations. The coadministration of REYATAZ and buprenorphine without ritonavir is not recommended.

**Table 16: Established and Other Potentially Significant Drug Interactions: Alteration in Dose or Regimen May Be Recommended Based on Drug Interaction Studies<sup>a</sup> or Predicted Interactions (Information in the table applies to REYATAZ with or without ritonavir, unless otherwise indicated)**

<b>Concomitant Drug Class: Specific Drugs</b>	<b>Effect on Concentration of Atazanavir or Concomitant Drug</b>	<b>Clinical Comment</b>
<i>PDE5 inhibitors:</i> sildenafil, tadalafil, vardenafil	↑ sildenafil ↑ tadalafil ↑ vardenafil	<p>Coadministration with REYATAZ has not been studied but may result in an increase in PDE5 inhibitor-associated adverse reactions, including hypotension, syncope, visual disturbances, and priapism.</p> <p><b>Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH):</b></p> <p>Use of REVATIO® (sildenafil) for the treatment of pulmonary hypertension (PAH) is contraindicated with REYATAZ [see <i>Contraindications (4)</i>].</p> <p>The following dose adjustments are recommended for the use of ADCIRCA® (tadalafil) with REYATAZ:</p> <p>Coadministration of ADCIRCA® in patients on REYATAZ (with or without ritonavir):</p> <ul style="list-style-type: none"> <li>For patients receiving REYATAZ (with or without ritonavir) for at least one week, start ADCIRCA® at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability.</li> </ul> <p>Coadministration of REYATAZ (with or without ritonavir) in patients on ADCIRCA®:</p> <ul style="list-style-type: none"> <li>Avoid the use of ADCIRCA® when starting REYATAZ (with or without ritonavir). Stop ADCIRCA® at least 24 hours before starting REYATAZ (with or without ritonavir). At least one week after starting REYATAZ (with or without ritonavir), resume ADCIRCA® at 20 mg once daily. Increase to 40 mg once daily based on individual tolerability.</li> </ul> <p><b>Use of PDE5 inhibitors for erectile dysfunction:</b></p> <p>Use VIAGRA® (sildenafil) with caution at reduced doses of 25 mg every 48 hours with increased monitoring for adverse events.</p> <p>Use CIALIS® (tadalafil) with caution at reduced doses of 10 mg every 72 hours with increased monitoring for adverse events.</p> <p><b>REYATAZ/ritonavir:</b> Use vardenafil with caution at reduced doses of no more than 2.5 mg every 72 hours with increased monitoring for adverse reactions.</p> <p><b>REYATAZ:</b> Use vardenafil with caution at reduced doses of no more than 2.5 mg every 24 hours with increased monitoring for adverse reactions.</p>
<i>Proton-pump inhibitors:</i> omeprazole	↓ atazanavir	<p>Plasma concentrations of atazanavir were substantially decreased when REYATAZ 400 mg or REYATAZ 300 mg/ritonavir 100 mg once daily was administered with omeprazole 40 mg once daily, which may result in loss of therapeutic effect and development of resistance.</p> <p><b>In treatment-naïve patients:</b></p> <p>The proton-pump inhibitor (PPI) dose should not exceed a dose comparable to omeprazole 20 mg and must be taken approximately 12 hours prior to the REYATAZ 300 mg with ritonavir 100 mg dose.</p> <p><b>In treatment-experienced patients:</b></p> <p>The use of PPIs in treatment-experienced patients receiving REYATAZ is not recommended.</p>

<sup>a</sup> For magnitude of interactions see *Clinical Pharmacology, Tables 21 and 22 (12.3)*.

<sup>b</sup> See *Contraindications (4), Table 6* for orally administered midazolam.



<sup>c</sup> In combination with atazanavir 300 mg and ritonavir 100 mg once daily.

<sup>d</sup> In combination with atazanavir 400 mg once daily.

## 7.4 Drugs with No Observed or Predicted Interactions with REYATAZ

Clinically significant interactions are not expected between atazanavir and substrates of CYP2C19, CYP2C9, CYP2D6, CYP2B6, CYP2A6, CYP1A2, or CYP2E1. Clinically significant interactions are not expected between atazanavir when administered with ritonavir and substrates of CYP2C8. See the complete prescribing information for ritonavir for information on other potential drug interactions with ritonavir.

Based on known metabolic profiles, clinically significant drug interactions are not expected between REYATAZ and dapsone, trimethoprim/sulfamethoxazole, azithromycin, or erythromycin. REYATAZ does not interact with substrates of CYP2D6 (eg, nortriptyline, desipramine, metoprolol). Additionally, no clinically significant drug interactions were observed when REYATAZ was coadministered with methadone, fluconazole, acetaminophen, or atenolol. *[See Clinical Pharmacology, Tables 21 and 22 (12.3).]*

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Pregnancy Category B

***Antiretroviral Pregnancy Registry:*** To monitor maternal-fetal outcomes of pregnant women exposed to REYATAZ, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients by calling 1-800-258-4263.

#### Risk Summary

Atazanavir has been evaluated in a limited number of women during pregnancy and postpartum. Available human and animal data suggest that atazanavir does not increase the risk of major birth defects overall compared to the background rate. However, because the studies in humans cannot rule out the possibility of harm, REYATAZ should be used during pregnancy only if clearly needed.

Cases of lactic acidosis syndrome, sometimes fatal, and symptomatic hyperlactatemia have occurred in pregnant women using REYATAZ in combination with nucleoside analogues. Nucleoside analogues are associated with an increased risk of lactic acidosis syndrome.



Hyperbilirubinemia occurs frequently in patients who take REYATAZ, including pregnant women. All infants, including neonates exposed to REYATAZ *in utero*, should be monitored for the development of severe hyperbilirubinemia during the first few days of life.

## Clinical Considerations

### *Dosing During Pregnancy and the Postpartum Period:*

- REYATAZ should not be administered without ritonavir.
- REYATAZ should only be administered to pregnant women with HIV-1 strains susceptible to atazanavir.
- For pregnant patients, no dose adjustment is required for REYATAZ with the following exceptions:
  - For treatment-experienced pregnant women during the second or third trimester, when REYATAZ is coadministered with either an H<sub>2</sub>-receptor antagonist **or** tenofovir, REYATAZ 400 mg with ritonavir 100 mg once daily is recommended. There are insufficient data to recommend a REYATAZ dose for use with both an H<sub>2</sub>-receptor antagonist *and* tenofovir in treatment-experienced pregnant women.
- No dose adjustment is required for postpartum patients. However, patients should be closely monitored for adverse events because atazanavir exposures could be higher during the first 2 months after delivery. [See *Dosage and Administration* (2.5) and *Clinical Pharmacology* (12.3).]

## Human Data

**Clinical Trials:** In clinical trial AI424-182, REYATAZ/ritonavir (300/100 mg or 400/100 mg) in combination with zidovudine/lamivudine was administered to 41 HIV-infected pregnant women during the second or third trimester. Among the 39 women who completed the study, 38 women achieved an HIV RNA <50 copies/mL at time of delivery. Six of 20 (30%) women on REYATAZ/ritonavir 300/100 mg and 13 of 21 (62%) women on REYATAZ/ritonavir 400/100 mg experienced hyperbilirubinemia (total bilirubin greater than or equal to 2.6 times ULN). There were no cases of lactic acidosis observed in clinical trial AI424-182.

Atazanavir drug concentrations in fetal umbilical cord blood were approximately 12% to 19% of maternal concentrations. Among the 40 infants born to 40 HIV-infected pregnant women, all had test results that were negative for HIV-1 DNA at the time of delivery and/or during the first 6 months postpartum. All 40 infants received antiretroviral prophylactic treatment containing zidovudine. No evidence of severe hyperbilirubinemia (total bilirubin levels greater than 20 mg/dL) or acute or chronic bilirubin encephalopathy was observed among neonates in this

study. However, 10/36 (28%) infants (6 greater than or equal to 38 weeks gestation and 4 less than 38 weeks gestation) had bilirubin levels of 4 mg/dL or greater within the first day of life.

Lack of ethnic diversity was a study limitation. In the study population, 33/40 (83%) infants were Black/African American, who have a lower incidence of neonatal hyperbilirubinemia than Caucasians and Asians. In addition, women with Rh incompatibility were excluded, as well as women who had a previous infant who developed hemolytic disease and/or had neonatal pathologic jaundice (requiring phototherapy).

Additionally, of the 38 infants who had glucose samples collected in the first day of life, 3 had adequately collected serum glucose samples with values of <40 mg/dL that could not be attributed to maternal glucose intolerance, difficult delivery, or sepsis.

**Antiretroviral Pregnancy Registry Data:** As of January 2010, the Antiretroviral Pregnancy Registry (APR) has received prospective reports of 635 exposures to atazanavir-containing regimens (425 exposed in the first trimester and 160 and 50 exposed in second and third trimester, respectively). Birth defects occurred in 9 of 393 (2.3%) live births (first trimester exposure) and 5 of 212 (2.4%) live births (second/third trimester exposure). Among pregnant women in the U.S. reference population, the background rate of birth defects is 2.7%. There was no association between atazanavir and overall birth defects observed in the APR.

## **Pharmacokinetics of Atazanavir in Pregnancy**

*[See Clinical Pharmacology (12.3).]*

### **Animal Data**

In animal reproduction studies, there was no evidence of teratogenicity in offspring born to animals at systemic drug exposure levels (AUC) 0.7 (in rabbits) to 1.2 (in rats) times those observed at the human clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir). In pre- and postnatal development studies in the rat, atazanavir caused body weight loss or weight gain suppression in the animal offspring with maternal drug exposure (AUC) 1.3 times the human exposure at this clinical dose. However, maternal toxicity also occurred at this exposure level.

## **8.3 Nursing Mothers**

**The Centers for Disease Control and Prevention recommend that HIV-infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV.** It is not known whether atazanavir is present in human milk. Because of both the potential for HIV transmission

and the potential for serious adverse reactions in nursing infants, **mothers should be instructed not to breastfeed if they are taking REYATAZ.**

## 8.4 Pediatric Use

REYATAZ is indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection in pediatric patients 3 months of age and older weighing at least 10 kg. REYATAZ is not recommended for use in pediatric patients below the age of 3 months due to the risk of kernicterus [see *Indications and Usage (1)*]. All REYATAZ contraindications, warnings, and precautions apply to pediatric patients [see *Contraindications (4)* and *Warnings and Precautions (5)*].

The safety, pharmacokinetic profile, and virologic response of REYATAZ in pediatric patients at least 3 months of age and older weighing at least 10 kg were established in three open-label, multicenter clinical trials: PACTG 1020A, AI424-451, and AI424-397 [see *Clinical Pharmacology (12.3)* and *Clinical Studies (14.3)*]. The safety profile in pediatric patients was generally similar to that observed in adults [see *Adverse Reactions (6.2)*]. See *Dosage and Administration (2.3, 2.4)* for dosing recommendations for the use of REYATAZ capsules and REYATAZ oral powder in pediatric patients.

## 8.5 Geriatric Use

Clinical studies of REYATAZ did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Based on a comparison of mean single-dose pharmacokinetic values for  $C_{max}$  and AUC, a dose adjustment based upon age is not recommended. In general, appropriate caution should be exercised in the administration and monitoring of REYATAZ in elderly patients reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy.

## 8.6 Age/Gender

A study of the pharmacokinetics of atazanavir was performed in young (n=29; 18-40 years) and elderly (n=30; ≥65 years) healthy subjects. There were no clinically significant pharmacokinetic differences observed due to age or gender.

## 8.7 Impaired Renal Function

REYATAZ is not recommended for use in HIV-treatment-experienced patients with end stage renal disease managed with hemodialysis. [See *Dosage and Administration (2.6)* and *Clinical Pharmacology (12.3)*.]

## 8.8 Impaired Hepatic Function

REYATAZ is not recommended for use in patients with severe hepatic impairment. REYATAZ/ritonavir is not recommended in patients with any degree of hepatic impairment. [See *Dosage and Administration* (2.7) and *Clinical Pharmacology* (12.3).]

## 10 OVERDOSAGE


Human experience of acute overdose with REYATAZ is limited. Single doses up to 1200 mg (three times the 400 mg maximum recommended dose) have been taken by healthy volunteers without symptomatic untoward effects. A single self-administered overdose of 29.2 g of REYATAZ in an HIV-infected patient (73 times the 400-mg recommended dose) was associated with asymptomatic bifascicular block and PR interval prolongation. These events resolved spontaneously. At REYATAZ doses resulting in high atazanavir exposures, jaundice due to indirect (unconjugated) hyperbilirubinemia (without associated liver function test changes) or PR interval prolongation may be observed. [See *Warnings and Precautions* (5.1, 5.3) and *Clinical Pharmacology* (12.2).]

Treatment of overdosage with REYATAZ should consist of general supportive measures, including monitoring of vital signs and ECG, and observations of the patient's clinical status. If indicated, elimination of unabsorbed atazanavir should be achieved by emesis or gastric lavage. Administration of activated charcoal may also be used to aid removal of unabsorbed drug. There is no specific antidote for overdose with REYATAZ. Since atazanavir is extensively metabolized by the liver and is highly protein bound, dialysis is unlikely to be beneficial in significant removal of this medicine.

## 11 DESCRIPTION

The active ingredient in REYATAZ capsules and oral powder is atazanavir sulfate, which is an HIV-1 protease inhibitor.

The chemical name for atazanavir sulfate is (3*S*,8*S*,9*S*,12*S*)-3,12-Bis(1,1-dimethylethyl)-8-hydroxy-4,11-dioxo-9-(phenylmethyl)-6-[[4-(2-pyridinyl)phenyl]methyl]-2,5,6,10,13-pentaazatetradecanedioic acid dimethyl ester, sulfate (1:1). Its molecular formula is C<sub>38</sub>H<sub>52</sub>N<sub>6</sub>O<sub>7</sub>•H<sub>2</sub>SO<sub>4</sub>, which corresponds to a molecular weight of 802.9 (sulfuric acid salt). The free base molecular weight is 704.9. Atazanavir sulfate has the following structural formula:



Atazanavir sulfate is a white to pale-yellow crystalline powder. It is slightly soluble in water (4-5 mg/mL, free base equivalent) with the pH of a saturated solution in water being about 1.9 at  $24 \pm 3^\circ\text{C}$ .

REYATAZ Capsules are available for oral administration in strengths of 150 mg, 200 mg, or 300 mg of atazanavir, which are equivalent to 170.8 mg, 227.8 mg, or 341.69 mg of atazanavir sulfate, respectively. The capsules also contain the following inactive ingredients: crospovidone, lactose monohydrate, and magnesium stearate. The capsule shells contain the following inactive ingredients: gelatin, FD&C Blue No. 2, titanium dioxide, black iron oxide, red iron oxide, and yellow iron oxide. The capsules are printed with ink containing shellac, titanium dioxide, FD&C Blue No. 2, isopropyl alcohol, ammonium hydroxide, propylene glycol, n-butyl alcohol, simethicone, and dehydrated alcohol.

REYATAZ oral powder comes in a packet containing 50 mg of atazanavir equivalent to 56.9 mg of atazanavir sulfate in 1.5 g of powder. The powder is off-white to pale yellow and contains the following inactive ingredients: aspartame, sucrose, and orange-vanilla flavor.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Atazanavir is an HIV-1 antiviral drug [*see Microbiology (12.4)*].

### 12.2 Pharmacodynamics

#### Effects on Electrocardiogram

Concentration- and dose-dependent prolongation of the PR interval in the electrocardiogram has been observed in healthy volunteers receiving atazanavir. In a placebo-controlled study (AI424-076), the mean ( $\pm$ SD) maximum change in PR interval from the predose value was 24 ( $\pm$ 15) msec following oral dosing with 400 mg of atazanavir (n=65) compared to

13 ( $\pm$ 11) msec following dosing with placebo (n=67). The PR interval prolongations in this study were asymptomatic. There is limited information on the potential for a pharmacodynamic interaction in humans between atazanavir and other drugs that prolong the PR interval of the electrocardiogram. [See *Warnings and Precautions* (5.1).]

Electrocardiographic effects of atazanavir were determined in a clinical pharmacology study of 72 healthy subjects. Oral doses of 400 mg (maximum recommended dosage) and 800 mg (twice the maximum recommended dosage) were compared with placebo; there was no concentration-dependent effect of atazanavir on the QTc interval (using Fridericia's correction). In 1793 HIV-infected patients receiving antiretroviral regimens, QTc prolongation was comparable in the atazanavir and comparator regimens. No atazanavir-treated healthy subject or HIV-infected patient in clinical trials had a QTc interval >500 msec. [See *Warnings and Precautions* (5.1).]

## 12.3 Pharmacokinetics

The pharmacokinetics of atazanavir were evaluated in healthy adult volunteers and in HIV-infected patients after administration of REYATAZ 400 mg once daily and after administration of REYATAZ 300 mg with ritonavir 100 mg once daily (see Table 17).

**Table 17: Steady-State Pharmacokinetics of Atazanavir in Healthy Subjects or HIV-Infected Patients in the Fed State**

Parameter	400 mg once daily		300 mg with ritonavir 100 mg once daily	
	Healthy Subjects (n=14)	HIV-Infected Patients (n=13)	Healthy Subjects (n=28)	HIV-Infected Patients (n=10)
$C_{max}$ (ng/mL)				
Geometric mean (CV%)	5199 (26)	2298 (71)	6129 (31)	4422 (58)
Mean (SD)	5358 (1371)	3152 (2231)	6450 (2031)	5233 (3033)
$T_{max}$ (h)				
Median	2.5	2.0	2.7	3.0
AUC (ng•h/mL)				
Geometric mean (CV%)	28132 (28)	14874 (91)	57039 (37)	46073 (66)
Mean (SD)	29303 (8263)	22262 (20159)	61435 (22911)	53761 (35294)
T-half (h)				
Mean (SD)	7.9 (2.9)	6.5 (2.6)	18.1 (6.2) <sup>a</sup>	8.6 (2.3)

**Table 17: Steady-State Pharmacokinetics of Atazanavir in Healthy Subjects or HIV-Infected Patients in the Fed State**


Parameter	400 mg once daily		300 mg with ritonavir 100 mg once daily	
	Healthy Subjects (n=14)	HIV-Infected Patients (n=13)	Healthy Subjects (n=28)	HIV-Infected Patients (n=10)
$C_{min}$ (ng/mL)				
Geometric mean (CV%)	159 (88)	120 (109)	1227 (53)	636 (97)
Mean (SD)	218 (191)	273 (298) <sup>b</sup>	1441 (757)	862 (838)

<sup>a</sup> n=26.

<sup>b</sup> n=12.

Figure 1 displays the mean plasma concentrations of atazanavir at steady state after REYATAZ 400 mg once daily (as two 200-mg capsules) with a light meal and after REYATAZ 300 mg (as two 150-mg capsules) with ritonavir 100 mg once daily with a light meal in HIV-infected adult patients.

**Figure 1: Mean (SD) Steady-State Plasma Concentrations of Atazanavir 400 mg (n=13) and 300 mg with Ritonavir (n=10) for HIV-Infected Adult Patients**





## Absorption

Atazanavir is rapidly absorbed with a  $T_{\max}$  of approximately 2.5 hours. Atazanavir demonstrates nonlinear pharmacokinetics with greater than dose-proportional increases in AUC and  $C_{\max}$  values over the dose range of 200 to 800 mg once daily. Steady state is achieved between Days 4 and 8, with an accumulation of approximately 2.3 fold.

## Food Effect

Administration of REYATAZ with food enhances bioavailability and reduces pharmacokinetic variability. Administration of a single 400-mg dose of REYATAZ with a light meal (357 kcal, 8.2 g fat, 10.6 g protein) resulted in a 70% increase in AUC and 57% increase in  $C_{\max}$  relative to the fasting state. Administration of a single 400-mg dose of REYATAZ with a high-fat meal (721 kcal, 37.3 g fat, 29.4 g protein) resulted in a mean increase in AUC of 35% with no change in  $C_{\max}$  relative to the fasting state. Administration of REYATAZ with either a light meal or high-fat meal decreased the coefficient of variation of AUC and  $C_{\max}$  by approximately one-half compared to the fasting state.

Coadministration of a single 300-mg dose of REYATAZ and a 100-mg dose of ritonavir with a light meal (336 kcal, 5.1 g fat, 9.3 g protein) resulted in a 33% increase in the AUC and a 40% increase in both the  $C_{\max}$  and the 24-hour concentration of atazanavir relative to the fasting state. Coadministration with a high-fat meal (951 kcal, 54.7 g fat, 35.9 g protein) did not affect the AUC of atazanavir relative to fasting conditions and the  $C_{\max}$  was within 11% of fasting values. The 24-hour concentration following a high-fat meal was increased by approximately 33% due to delayed absorption; the median  $T_{\max}$  increased from 2.0 to 5.0 hours. Coadministration of REYATAZ with ritonavir with either a light or a high-fat meal decreased the coefficient of variation of AUC and  $C_{\max}$  by approximately 25% compared to the fasting state.

## Distribution

Atazanavir is 86% bound to human serum proteins and protein binding is independent of concentration. Atazanavir binds to both alpha-1-acid glycoprotein (AAG) and albumin to a similar extent (89% and 86%, respectively). In a multiple-dose study in HIV-infected patients dosed with REYATAZ 400 mg once daily with a light meal for 12 weeks, atazanavir was detected in the cerebrospinal fluid and semen. The cerebrospinal fluid/plasma ratio for atazanavir (n=4) ranged between 0.0021 and 0.0226 and seminal fluid/plasma ratio (n=5) ranged between 0.11 and 4.42.



## Metabolism

Atazanavir is extensively metabolized in humans. The major biotransformation pathways of atazanavir in humans consisted of monooxygenation and dioxygenation. Other minor biotransformation pathways for atazanavir or its metabolites consisted of glucuronidation, N-dealkylation, hydrolysis, and oxygenation with dehydrogenation. Two minor metabolites of atazanavir in plasma have been characterized. Neither metabolite demonstrated *in vitro* antiviral activity. *In vitro* studies using human liver microsomes suggested that atazanavir is metabolized by CYP3A.

## Elimination

Following a single 400-mg dose of  $^{14}\text{C}$ -atazanavir, 79% and 13% of the total radioactivity was recovered in the feces and urine, respectively. Unchanged drug accounted for approximately 20% and 7% of the administered dose in the feces and urine, respectively. The mean elimination half-life of atazanavir in healthy volunteers (n=214) and HIV-infected adult patients (n=13) was approximately 7 hours at steady state following a dose of 400 mg daily with a light meal.

## Specific Populations

### ***Renal Impairment***

In healthy subjects, the renal elimination of unchanged atazanavir was approximately 7% of the administered dose. REYATAZ has been studied in adult subjects with severe renal impairment (n=20), including those on hemodialysis, at multiple doses of 400 mg once daily. The mean atazanavir  $C_{\max}$  was 9% lower, AUC was 19% higher, and  $C_{\min}$  was 96% higher in subjects with severe renal impairment not undergoing hemodialysis (n=10), than in age-, weight-, and gender-matched subjects with normal renal function. In a 4-hour dialysis session, 2.1% of the administered dose was removed. When atazanavir was administered either prior to, or following hemodialysis (n=10), the geometric means for  $C_{\max}$ , AUC, and  $C_{\min}$  were approximately 25% to 43% lower compared to subjects with normal renal function. The mechanism of this decrease is unknown. REYATAZ is not recommended for use in HIV-treatment-experienced patients with end stage renal disease managed with hemodialysis. [See *Dosage and Administration* (2.6).]

## **Hepatic Impairment**

REYATAZ has been studied in adult subjects with moderate-to-severe hepatic impairment (14 Child-Pugh B and 2 Child-Pugh C subjects) after a single 400-mg dose. The mean  $AUC_{(0-\infty)}$  was 42% greater in subjects with impaired hepatic function than in healthy volunteers. The mean half-life of atazanavir in hepatically impaired subjects was 12.1 hours compared to 6.4 hours in healthy volunteers. A dose reduction to 300 mg is recommended for patients with moderate hepatic impairment (Child-Pugh Class B) who have not experienced prior virologic failure as increased concentrations of atazanavir are expected. REYATAZ is not recommended for use in patients with severe hepatic impairment. The pharmacokinetics of REYATAZ in combination with ritonavir has not been studied in subjects with hepatic impairment; thus, coadministration of REYATAZ with ritonavir is not recommended for use in patients with any degree of hepatic impairment. [See *Dosage and Administration* (2.7).]

## **Pediatrics**

The pharmacokinetic parameters for atazanavir at steady state in pediatric patients taking the powder formulation are summarized in Table 18 by weight ranges that correspond to the recommended doses. [See *Dosage and Administration* (2.4).]

**Table 18: Steady-State Pharmacokinetics of Atazanavir (powder formulation) with Ritonavir in HIV-Infected Pediatric Patients**

Body Weight (range in kg) [n]	atazanavir/ritonavir Dose (mg)	C <sub>max</sub> ng/mL Geometric Mean (CV%)	AUC ng•h/mL Geometric Mean (CV%)	C <sub>min</sub> ng/mL Geometric Mean (CV%)
10 to <15 [18]	200/80	5197 (53%)	50305 (67%)	572 (111%)
15 to <25 [31]	250/80	5386 (47%)	55525 (46%)	678 (69%)

The pharmacokinetic parameters for atazanavir at steady state in pediatric patients taking the capsule formulation were predicted by a population pharmacokinetic model and are summarized in Table 19 by weight ranges that correspond to the recommended doses. [See *Dosage and Administration* (2.4).]

**Table 19: Predicted Steady-State Pharmacokinetics of Atazanavir (capsule formulation) with Ritonavir in HIV-Infected Pediatric Patients**

Body Weight (range in kg)	atazanavir/ritonavir Dose (mg)	C <sub>max</sub> ng/mL Geometric Mean (CV%)	AUC ng•h/mL Geometric Mean (CV%)	C <sub>min</sub> ng/mL Geometric Mean (CV%)
15 to <20	150/100	5213 (78.7%)	42902 (77.0%)	504 (99.5%)
20 to <40	200/100	4954 (81.7%)	42999 (78.5%)	562 (98.9%)
≥40	300/100	5040 (84.6%)	46777 (80.6%)	691 (98.5%)

### **Pregnancy**

The pharmacokinetic data from HIV-infected pregnant women receiving REYATAZ Capsules with ritonavir are presented in Table 20.

**Table 20: Steady-State Pharmacokinetics of Atazanavir with Ritonavir in HIV-Infected Pregnant Women in the Fed State**

Pharmacokinetic Parameter	Atazanavir 300 mg with ritonavir 100 mg		
	2nd Trimester (n=5 <sup>a</sup> )	3rd Trimester (n=20)	Postpartum <sup>b</sup> (n=34)
C <sub>max</sub> ng/mL	3078.85	3291.46	5721.21
Geometric mean (CV%)	(50)	(48)	(31)
AUC ng•h/mL	27657.1	34251.5	61990.4
Geometric mean (CV%)	(43)	(43)	(32)
C <sub>min</sub> ng/mL <sup>c</sup>	538.70	668.48	1462.59
Geometric mean (CV%)	(46)	(50)	(45)

<sup>a</sup> Available data during the 2nd trimester are limited.

<sup>b</sup> Atazanavir peak concentrations and AUCs were found to be approximately 28% to 43% higher during the postpartum period (4-12 weeks) than those observed historically in HIV-infected, non-pregnant patients. Atazanavir plasma trough concentrations were approximately 2.2-fold higher during the postpartum period when compared to those observed historically in HIV-infected, non-pregnant patients.

<sup>c</sup> C<sub>min</sub> is concentration 24 hours post-dose.

### **Drug Interaction Data**

Atazanavir is a metabolism-dependent CYP3A inhibitor, with a K<sub>inact</sub> value of 0.05 to 0.06 min<sup>-1</sup> and K<sub>i</sub> value of 0.84 to 1.0 μM. Atazanavir is also a direct inhibitor for UGT1A1 (K<sub>i</sub>=1.9 μM) and CYP2C8 (K<sub>i</sub>=2.1 μM).

Atazanavir has been shown *in vivo* not to induce its own metabolism nor to increase the biotransformation of some drugs metabolized by CYP3A. In a multiple-dose study, REYATAZ decreased the urinary ratio of endogenous 6 $\beta$ -OH cortisol to cortisol versus baseline, indicating that CYP3A production was not induced.

Drug interaction studies were performed with REYATAZ and other drugs likely to be coadministered and some drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration of REYATAZ on the AUC, C<sub>max</sub>, and C<sub>min</sub> are summarized in Tables 21 and 22. For information regarding clinical recommendations, see *Drug Interactions* (7).

**Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
atenolol	50 mg QD, d 7–11 (n=19) and d 19–23	400 mg QD, d 1–11 (n=19)	1.00 (0.89, 1.12)	0.93 (0.85, 1.01)	0.74 (0.65, 0.86)
boceprevir	800 mg TID, d 1–6, 25–31	300 mg QD/ritonavir 100 mg QD, d 10–31	atazanavir: 0.75 (0.64–0.88)	atazanavir: 0.65 (0.55–0.78)	atazanavir: 0.51 (0.44–0.61)
			ritonavir: 0.73 (0.64–0.83)	ritonavir: 0.64 (0.58–0.72)	ritonavir: 0.55 (0.45–0.67)
clarithromycin	500 mg BID, d 7–10 (n=29) and d 18–21	400 mg QD, d 1–10 (n=29)	1.06 (0.93, 1.20)	1.28 (1.16, 1.43)	1.91 (1.66, 2.21)
didanosine (ddI) (buffered tablets) plus stavudine (d4T) <sup>b</sup>	ddI: 200 mg $\times$ 1 dose, d4T: 40 mg $\times$ 1 dose (n=31)	400 mg $\times$ 1 dose simultaneously with ddI and d4T (n=31)	0.11 (0.06, 0.18)	0.13 (0.08, 0.21)	0.16 (0.10, 0.27)
	ddI: 200 mg $\times$ 1 dose, d4T: 40 mg $\times$ 1 dose (n=32)	400 mg $\times$ 1 dose 1 h after ddI + d4T (n=32)	1.12 (0.67, 1.18)	1.03 (0.64, 1.67)	1.03 (0.61, 1.73)
ddI (enteric-coated [EC] capsules) <sup>c</sup>	400 mg d 8 (fed) (n=34)	400 mg QD, d 2–8 (n=34)	1.03 (0.93, 1.14)	0.99 (0.91, 1.08)	0.98 (0.89, 1.08)
	400 mg d 19 (fed) (n=31)	300 mg/ritonavir 100 mg QD, d 9–19 (n=31)	1.04 (1.01, 1.07)	1.00 (0.96, 1.03)	0.87 (0.82, 0.92)
diltiazem	180 mg QD, d 7–11 (n=30) and d 19–23	400 mg QD, d 1–11 (n=30)	1.04 (0.96, 1.11)	1.00 (0.95, 1.05)	0.98 (0.90, 1.07)
efavirenz	600 mg QD, d 7–20 (n=27)	400 mg QD, d 1–20 (n=27)	0.41 (0.33, 0.51)	0.26 (0.22, 0.32)	0.07 (0.05, 0.10)
	600 mg QD, d 7–20 (n=13)	400 mg QD, d 1–6 (n=23) then 300 mg/ritonavir 100 mg QD, 2 h before efavirenz, d 7–20 (n=13)	1.14 (0.83, 1.58)	1.39 (1.02, 1.88)	1.48 (1.24, 1.76)

**Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
	600 mg QD, d 11–24 (pm) (n=14)	300 mg QD/ritonavir 100 mg QD, d 1–10 (pm) (n=22), then 400 mg QD/ritonavir 100 mg QD, d 11–24 (pm), (simultaneously with efavirenz) (n=14)	1.17 (1.08, 1.27)	1.00 (0.91, 1.10)	0.58 (0.49, 0.69)
famotidine	40 mg BID, d 7–12 (n=15)	400 mg QD, d 1–6 (n=45), d 7–12 (simultaneous administration) (n=15)	0.53 (0.34, 0.82)	0.59 (0.40, 0.87)	0.58 (0.37, 0.89)
	40 mg BID, d 7–12 (n=14)	400 mg QD (pm), d 1–6 (n=14), d 7–12 (10 h after, 2 h before famotidine) (n=14)	1.08 (0.82, 1.41)	0.95 (0.74, 1.21)	0.79 (0.60, 1.04)
	40 mg BID, d 11–20 <sup>d</sup> (n=14)	300 mg QD/ritonavir 100 mg QD, d 1–10 (n=46), d 11–20 <sup>d</sup> (simultaneous administration) (n=14)	0.86 (0.79, 0.94)	0.82 (0.75, 0.89)	0.72 (0.64, 0.81)
	20 mg BID, d 11–17 (n=18)	300 mg QD/ritonavir 100 mg QD/tenofovir 300 mg QD, d 1–10 (am) (n=39), d 11–17 (am) (simultaneous administration with am famotidine) (n=18) <sup>e,f</sup>	0.91 (0.84, 0.99)	0.90 (0.82, 0.98)	0.81 (0.69, 0.94)
	40 mg QD (pm), d 18–24 (n=20)	300 mg QD/ritonavir 100 mg QD/tenofovir 300 mg QD, d 1–10 (am) (n=39), d 18–24 (am) (12 h after pm famotidine) (n=20) <sup>f</sup>	0.89 (0.81, 0.97)	0.88 (0.80, 0.96)	0.77 (0.63, 0.93)
	40 mg BID, d 18–24 (n=18)	300 mg QD/ritonavir 100 mg QD/tenofovir 300 mg QD, d 1–10 (am) (n=39), d 18–24 (am) (10 h after pm famotidine and 2 h before am famotidine) (n=18) <sup>f</sup>	0.74 (0.66, 0.84)	0.79 (0.70, 0.88)	0.72 (0.63, 0.83)
	40 mg BID, d 11–20 (n=15)	300 mg QD/ritonavir 100 mg QD, d 1–10 (am) (n=46), then 400 mg QD/ritonavir 100 mg QD, d 11–20 (am) (n=15)	1.02 (0.87, 1.18)	1.03 (0.86, 1.22)	0.86 (0.68, 1.08)

**Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
fluconazole	200 mg QD, d 11–20 (n=29)	300 mg QD/ritonavir 100 mg QD, d 1–10 (n=19), d 11–20 (n=29)	1.03 (0.95, 1.11)	1.04 (0.95, 1.13)	0.98 (0.85, 1.13)
ketoconazole	200 mg QD, d 7–13 (n=14)	400 mg QD, d 1–13 (n=14)	0.99 (0.77, 1.28)	1.10 (0.89, 1.37)	1.03 (0.53, 2.01)
nevirapine <sup>g,h</sup>	200 mg BID, d 1–23 (n=23)	300 mg QD/ritonavir 100 mg QD, d 4–13, then 400 mg QD/ritonavir 100 mg QD, d 14–23 (n=23) <sup>i</sup>	0.72 (0.60, 0.86)	0.58 (0.48, 0.71)	0.28 (0.20, 0.40)
			1.02 (0.85, 1.24)	0.81 (0.65, 1.02)	0.41 (0.27, 0.60)
omeprazole	40 mg QD, d 7–12 (n=16) <sup>j</sup>	400 mg QD, d 1–6 (n=48), d 7–12 (n=16)	0.04 (0.04, 0.05)	0.06 (0.05, 0.07)	0.05 (0.03, 0.07)
	40 mg QD, d 11–20 (n=15) <sup>j</sup>	300 mg QD/ritonavir 100 mg QD, d 1–20 (n=15)	0.28 (0.24, 0.32)	0.24 (0.21, 0.27)	0.22 (0.19, 0.26)
	20 mg QD, d 17–23 (am) (n=13)	300 mg QD/ritonavir 100 mg QD, d 7–16 (pm) (n=27), d 17–23 (pm) (n=13) <sup>k,l</sup>	0.61 (0.46, 0.81)	0.58 (0.44, 0.75)	0.54 (0.41, 0.71)
	20 mg QD, d 17–23 (am) (n=14)	300 mg QD/ritonavir 100 mg QD, d 7–16 (am) (n=27), then 400 mg QD/ritonavir 100 mg QD, d 17–23 (am) (n=14) <sup>m,n</sup>	0.69 (0.58, 0.83)	0.70 (0.57, 0.86)	0.69 (0.54, 0.88)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.13 (0.96, 1.32)	1.06 (0.90, 1.26)	NA
rifabutin	150 mg QD, d 15–28 (n=7)	400 mg QD, d 1–28 (n=7)	1.34 (1.14, 1.59)	1.15 (0.98, 1.34)	1.13 (0.68, 1.87)
rifampin	600 mg QD, d 17–26 (n=16)	300 mg QD/ritonavir 100 mg QD, d 7–16 (n=48), d 17–26 (n=16)	0.47 (0.41, 0.53)	0.28 (0.25, 0.32)	0.02 (0.02, 0.03)
ritonavir <sup>o</sup>	100 mg QD, d 11–20 (n=28)	300 mg QD, d 1–20 (n=28)	1.86 (1.69, 2.05)	3.38 (3.13, 3.63)	11.89 (10.23, 13.82)
telaprevir	750 mg q8h for 10 days (n=7)	300 mg QD/ritonavir 100 mg QD for 20 days (n=7)	0.85 (0.73, 0.98)	1.17 (0.97, 1.43)	1.85 (1.40, 2.44)
tenofovir <sup>p</sup>	300 mg QD, d 9–16 (n=34)	400 mg QD, d 2–16 (n=34)	0.79 (0.73, 0.86)	0.75 (0.70, 0.81)	0.60 (0.52, 0.68)
	300 mg QD, d 15–42 (n=10)	300 mg/ritonavir 100 mg QD, d 1–42 (n=10)	0.72 <sup>q</sup> (0.50, 1.05)	0.75 <sup>q</sup> (0.58, 0.97)	0.77 <sup>q</sup> (0.54, 1.10)

**Table 21: Drug Interactions: Pharmacokinetic Parameters for Atazanavir in the Presence of Coadministered Drugs<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Atazanavir Pharmacokinetic Parameters with/without Coadministered Drug; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
voriconazole (Subjects with at least one functional CYP2C19 allele)	200 mg BID, d 2–3, 22–30; 400 mg BID, d 1, 21 (n=20)	300 mg/ritonavir 100 mg QD, d 11–30 (n=20)	0.87 (0.80, 0.96)	0.88 (0.82, 0.95)	0.80 (0.72, 0.90)
voriconazole (Subjects without a functional CYP2C19 allele)	50 mg BID, d 2–3, 22–30; 100 mg BID, d 1, 21 (n=8)	300 mg/ritonavir 100 mg QD, d 11–30 (n=8)	0.81 (0.66, 1.00)	0.80 (0.65, 0.97)	0.69 (0.54, 0.87)

<sup>a</sup> Data provided are under fed conditions unless otherwise noted.

<sup>b</sup> All drugs were given under fasted conditions.

<sup>c</sup> 400 mg ddi EC and REYATAZ were administered together with food on Days 8 and 19.

<sup>d</sup> REYATAZ 300 mg plus ritonavir 100 mg once daily coadministered with famotidine 40 mg twice daily resulted in atazanavir geometric mean C<sub>max</sub> that was similar and AUC and C<sub>min</sub> values that were 1.79- and 4.46-fold higher relative to REYATAZ 400 mg once daily alone.

<sup>e</sup> Similar results were noted when famotidine 20 mg BID was administered 2 hours after and 10 hours before atazanavir 300 mg and ritonavir 100 mg plus tenofovir 300 mg.

<sup>f</sup> Atazanavir/ritonavir/tenofovir was administered after a light meal.

<sup>g</sup> Study was conducted in HIV-infected individuals.

<sup>h</sup> Compared with atazanavir 400 mg historical data without nevirapine (n=13), the ratio of geometric means (90% confidence intervals) for C<sub>max</sub>, AUC, and C<sub>min</sub> were 1.42 (0.98, 2.05), 1.64 (1.11, 2.42), and 1.25 (0.66, 2.36), respectively, for atazanavir/ritonavir 300/100 mg; and 2.02 (1.42, 2.87), 2.28 (1.54, 3.38), and 1.80 (0.94, 3.45), respectively, for atazanavir/ritonavir 400/100 mg.

<sup>i</sup> Parallel group design; n=23 for atazanavir/ritonavir plus nevirapine, n=22 for atazanavir 300 mg/ritonavir 100 mg without nevirapine. Subjects were treated with nevirapine prior to study entry.

<sup>j</sup> Omeprazole 40 mg was administered on an empty stomach 2 hours before REYATAZ.

<sup>k</sup> Omeprazole 20 mg was administered 30 minutes prior to a light meal in the morning and REYATAZ 300 mg plus ritonavir 100 mg in the evening after a light meal, separated by 12 hours from omeprazole.

<sup>l</sup> REYATAZ 300 mg plus ritonavir 100 mg once daily separated by 12 hours from omeprazole 20 mg daily resulted in increases in atazanavir geometric mean AUC (10%) and C<sub>min</sub> (2.4-fold), with a decrease in C<sub>max</sub> (29%) relative to REYATAZ 400 mg once daily in the absence of omeprazole (study days 1–6).

<sup>m</sup> Omeprazole 20 mg was given 30 minutes prior to a light meal in the morning and REYATAZ 400 mg plus ritonavir 100 mg once daily after a light meal, 1 hour after omeprazole. Effects on atazanavir concentrations were similar when REYATAZ 400 mg plus ritonavir 100 mg was separated from omeprazole 20 mg by 12 hours.

<sup>n</sup> REYATAZ 400 mg plus ritonavir 100 mg once daily administered with omeprazole 20 mg once daily resulted in increases in atazanavir geometric mean AUC (32%) and C<sub>min</sub> (3.3-fold), with a decrease in C<sub>max</sub> (26%) relative to REYATAZ 400 mg once daily in the absence of omeprazole (study days 1–6).

<sup>o</sup> Compared with atazanavir 400 mg QD historical data, administration of atazanavir/ritonavir 300/100 mg QD increased the atazanavir geometric mean values of C<sub>max</sub>, AUC, and C<sub>min</sub> by 18%, 103%, and 671%, respectively.

<sup>p</sup> Note that similar results were observed in studies where administration of tenofovir and REYATAZ was separated by 12 hours.

<sup>q</sup> Ratio of atazanavir plus ritonavir plus tenofovir to atazanavir plus ritonavir. Atazanavir 300 mg plus ritonavir 100 mg results in higher atazanavir exposure than atazanavir 400 mg (see footnote <sup>o</sup>). The geometric mean values of atazanavir

pharmacokinetic parameters when coadministered with ritonavir and tenofovir were:  $C_{\max}$  = 3190 ng/mL,  $AUC = 34459 \text{ ng}\cdot\text{h/mL}$ , and  $C_{\min} = 491 \text{ ng/mL}$ . Study was conducted in HIV-infected individuals. NA = not available.

**Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			$C_{\max}$	AUC	$C_{\min}$
acetaminophen	1 gm BID, d 1–20 (n=10)	300 mg QD/ritonavir 100 mg QD, d 11–20 (n=10)	0.87 (0.77, 0.99)	0.97 (0.91, 1.03)	1.26 (1.08, 1.46)
atenolol	50 mg QD, d 7–11 (n=19) and d 19–23	400 mg QD, d 1–11 (n=19)	1.34 (1.26, 1.42)	1.25 (1.16, 1.34)	1.02 (0.88, 1.19)
boceprevir	800 mg TID, d 1–6, 25–31	300 mg QD/ritonavir 100 mg QD, d 10–31	0.93 (0.80, 1.08)	0.95 (0.87, 1.05)	0.82 (0.68, 0.98)
clarithromycin	500 mg BID, d 7–10 (n=21) and d 18–21	400 mg QD, d 1–10 (n=21)	1.50 (1.32, 1.71) OH-clarithromycin: 0.28 (0.24, 0.33)	1.94 (1.75, 2.16) OH-clarithromycin: 0.30 (0.26, 0.34)	2.60 (2.35, 2.88) OH-clarithromycin: 0.38 (0.34, 0.42)
didanosine (ddI) (buffered tablets) plus stavudine (d4T) <sup>b</sup>	ddI: 200 mg × 1 dose, d4T: 40 mg × 1 dose (n=31)	400 mg × 1 dose simultaneous with ddI and d4T (n=31)	ddI: 0.92 (0.84, 1.02) d4T: 1.08 (0.96, 1.22)	ddI: 0.98 (0.92, 1.05) d4T: 1.00 (0.97, 1.03)	NA  d4T: 1.04 (0.94, 1.16)
ddI (enteric-coated [EC] capsules) <sup>c</sup>	400 mg d 1 (fasted), d 8 (fed) (n=34)	400 mg QD, d 2–8 (n=34)	0.64 (0.55, 0.74)	0.66 (0.60, 0.74)	1.13 (0.91, 1.41)
	400 mg d 1 (fasted), d 19 (fed) (n=31)	300 mg QD/ritonavir 100 mg QD, d 9–19 (n=31)	0.62 (0.52, 0.74)	0.66 (0.59, 0.73)	1.25 (0.92, 1.69)
diltiazem	180 mg QD, d 7–11 (n=28) and d 19–23	400 mg QD, d 1–11 (n=28)	1.98 (1.78, 2.19) desacetyl-diltiazem: 2.72 (2.44, 3.03)	2.25 (2.09, 2.16) desacetyl-diltiazem: 2.65 (2.45, 2.87)	2.42 (2.14, 2.73) desacetyl-diltiazem: 2.21 (2.02, 2.42)
ethinyl estradiol & norethindrone <sup>d</sup>	Ortho-Novum <sup>®</sup> 7/7/7 QD, d 1–29 (n=19)	400 mg QD, d 16–29 (n=19)	ethinyl estradiol: 1.15 (0.99, 1.32) norethindrone: 1.67 (1.42, 1.96)	ethinyl estradiol: 1.48 (1.31, 1.68) norethindrone: 2.10 (1.68, 2.62)	ethinyl estradiol: 1.91 (1.57, 2.33) norethindrone: 3.62 (2.57, 5.09)
ethinyl estradiol & norgestimate <sup>e</sup>	Ortho Tri-Cyclen <sup>®</sup> QD, d 1–28 (n=18), then Ortho Tri-Cyclen <sup>®</sup> LO QD, d 29–42 <sup>f</sup> (n=14)	300 mg QD/ritonavir 100 mg QD, d 29–42 (n=14)	ethinyl estradiol: 0.84 (0.74, 0.95) 17-deacetyl norgestimate: <sup>g</sup> 1.68 (1.51, 1.88)	ethinyl estradiol: 0.81 (0.75, 0.87) 17-deacetyl norgestimate: <sup>g</sup> 1.85 (1.67, 2.05)	ethinyl estradiol: 0.63 (0.55, 0.71) 17-deacetyl norgestimate: <sup>g</sup> 2.02 (1.77, 2.31)



**Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
fluconazole	200 mg QD, d 1–10 (n=11) and 200 mg QD, d 11–20 (n=29)	300 mg QD/ritonavir 100 mg QD, d 11–20 (n=29)	1.05 (0.99, 1.10)	1.08 (1.02, 1.15)	1.07 (1.00, 1.15)
methadone	Stable maintenance dose, d 1–15 (n=16)	400 mg QD, d 2–15 (n=16)	(R)-methadone <sup>h</sup> 0.91 (0.84, 1.0) total: 0.85 (0.78, 0.93)	(R)-methadone <sup>h</sup> 1.03 (0.95, 1.10) total: 0.94 (0.87, 1.02)	(R)-methadone <sup>h</sup> 1.11 (1.02, 1.20) total: 1.02 (0.93, 1.12)
nevirapine <sup>i,j</sup>	200 mg BID, d 1–23 (n=23)	300 mg QD/ritonavir 100 mg QD, d 4–13, then	1.17 (1.09, 1.25)	1.25 (1.17, 1.34)	1.32 (1.22, 1.43)
		400 mg QD/ritonavir 100 mg QD, d 14–23 (n=23)	1.21 (1.11, 1.32)	1.26 (1.17, 1.36)	1.35 (1.25, 1.47)
omeprazole <sup>k</sup>	40 mg single dose, d 7 and d 20 (n=16)	400 mg QD, d 1–12 (n=16)	1.24 (1.04, 1.47)	1.45 (1.20, 1.76)	NA
rifabutin	300 mg QD, d 1–10 then 150 mg QD, d 11–20 (n=3)	600 mg QD, <sup>l</sup> d 11–20 (n=3)	1.18 (0.94, 1.48) 25-O-desacetyl-rifabutin: 8.20 (5.90, 11.40)	2.10 (1.57, 2.79) 25-O-desacetyl-rifabutin: 22.01 (15.97, 30.34)	3.43 (1.98, 5.96) 25-O-desacetyl-rifabutin: 75.6 (30.1, 190.0)
	150 mg twice weekly, d 1–15 (n=7)	300 mg QD/ritonavir 100 mg QD, d 1–17 (n=7)	2.49 <sup>m</sup> (2.03, 3.06) 25-O-desacetyl-rifabutin: 7.77 (6.13, 9.83)	1.48 <sup>m</sup> (1.19, 1.84) 25-O-desacetyl-rifabutin: 10.90 (8.14, 14.61)	1.40 <sup>m</sup> (1.05, 1.87) 25-O-desacetyl-rifabutin: 11.45 (8.15, 16.10)
pitavastatin	4 mg QD for 5 days	300 mg QD for 5 days	1.60 (1.39, 1.85)	1.31 (1.23, 1.39)	NA
rosiglitazone <sup>n</sup>	4 mg single dose, d 1, 7, 17 (n=14)	400 mg QD, d 2–7, then	1.08 (1.03, 1.13)	1.35 (1.26, 1.44)	NA
		300 mg QD/ritonavir 100 mg QD, d 8–17 (n=14)	0.97 (0.91, 1.04)	0.83 (0.77, 0.89)	NA
rosuvastatin	10 mg single dose	300 mg QD/ritonavir 100 mg QD for 7 days	↑ 7-fold <sup>o</sup>	↑ 3-fold <sup>o</sup>	NA
saquinavir <sup>p</sup> (soft gelatin capsules)	1200 mg QD, d 1–13 (n=7)	400 mg QD, d 7–13 (n=7)	4.39 (3.24, 5.95)	5.49 (4.04, 7.47)	6.86 (5.29, 8.91)

**Table 22: Drug Interactions: Pharmacokinetic Parameters for Coadministered Drugs in the Presence of REYATAZ<sup>a</sup>**

Coadministered Drug	Coadministered Drug Dose/Schedule	REYATAZ Dose/Schedule	Ratio (90% Confidence Interval) of Coadministered Drug Pharmacokinetic Parameters with/without REYATAZ; No Effect = 1.00		
			C <sub>max</sub>	AUC	C <sub>min</sub>
telaprevir	750 mg q8h for 10 days (n=14)	300 mg QD/ritonavir 100 mg QD for 20 days (n=14)	0.79 (0.74, 0.84)	0.80 (0.76, 0.85)	0.85 (0.75, 0.98)
tenofovir <sup>q</sup>	300 mg QD, d 9–16 (n=33) and d 24–30 (n=33)	400 mg QD, d 2–16 (n=33)	1.14 (1.08, 1.20)	1.24 (1.21, 1.28)	1.22 (1.15, 1.30)
	300 mg QD, d 1–7 (pm) (n=14) d 25–34 (pm) (n=12)	300 mg QD/ritonavir 100 mg QD, d 25–34 (am) (n=12) <sup>r</sup>	1.34 (1.20, 1.51)	1.37 (1.30, 1.45)	1.29 (1.21, 1.36)
voriconazole (Subjects with at least one functional CYP2C19 allele)	200 mg BID, d 2–3, 22–30; 400 mg BID, d 1, 21 (n=20)	300 mg/ritonavir 100 mg QD, d 11–30 (n=20)	0.90 (0.78, 1.04)	0.67 (0.58, 0.78)	0.61 (0.51, 0.72)
voriconazole (Subjects without a functional CYP2C19 allele)	50 mg BID, d 2–3, 22–30; 100 mg BID, d 1, 21 (n=8)	300 mg/ritonavir 100 mg QD, d 11–30 (n=8)	4.38 (3.55, 5.39)	5.61 (4.51, 6.99)	7.65 (5.71, 10.2)
lamivudine + zidovudine	150 mg lamivudine + 300 mg zidovudine BID, d 1–12 (n=19)	400 mg QD, d 7–12 (n=19)	lamivudine: 1.04 (0.92, 1.16) zidovudine: 1.05 (0.88, 1.24) zidovudine glucuronide: 0.95 (0.88, 1.02)	lamivudine: 1.03 (0.98, 1.08) zidovudine: 1.05 (0.96, 1.14) zidovudine glucuronide: 1.00 (0.97, 1.03)	lamivudine: 1.12 (1.04, 1.21) zidovudine: 0.69 (0.57, 0.84) zidovudine glucuronide: 0.82 (0.62, 1.08)

<sup>a</sup> Data provided are under fed conditions unless otherwise noted.

<sup>b</sup> All drugs were given under fasted conditions.

<sup>c</sup> 400 mg ddi EC and REYATAZ were administered together with food on Days 8 and 19.

<sup>d</sup> Upon further dose normalization of ethinyl estradiol 25 mcg with atazanavir relative to ethinyl estradiol 35 mcg without atazanavir, the ratio of geometric means (90% confidence intervals) for C<sub>max</sub>, AUC, and C<sub>min</sub> were 0.82 (0.73, 0.92), 1.06 (0.95, 1.17), and 1.35 (1.11, 1.63), respectively.

<sup>e</sup> Upon further dose normalization of ethinyl estradiol 35 mcg with atazanavir/ritonavir relative to ethinyl estradiol 25 mcg without atazanavir/ritonavir, the ratio of geometric means (90% confidence intervals) for C<sub>max</sub>, AUC, and C<sub>min</sub> were 1.17 (1.03, 1.34), 1.13 (1.05, 1.22), and 0.88 (0.77, 1.00), respectively.

<sup>f</sup> All subjects were on a 28 day lead-in period; one full cycle of Ortho Tri-Cyclen<sup>®</sup>. Ortho Tri-Cyclen<sup>®</sup> contains 35 mcg of ethinyl estradiol. Ortho Tri-Cyclen<sup>®</sup> LO contains 25 mcg of ethinyl estradiol. Results were dose normalized to an ethinyl estradiol dose of 35 mcg.

<sup>g</sup> 17-deacetyl norgestimate is the active component of norgestimate.

<sup>h</sup> (R)-methadone is the active isomer of methadone.

<sup>i</sup> Study was conducted in HIV-infected individuals.

<sup>j</sup> Subjects were treated with nevirapine prior to study entry.

- <sup>k</sup> Omeprazole was used as a metabolic probe for CYP2C19. Omeprazole was given 2 hours after REYATAZ on Day 7; and was given alone 2 hours after a light meal on Day 20.
- <sup>l</sup> Not the recommended therapeutic dose of atazanavir.
- <sup>m</sup> When compared to rifabutin 150 mg QD alone d1–10 (n=14). Total of Rifabutin + 25-O-desacetyl-rifabutin: AUC 2.19 (1.78, 2.69).
- <sup>n</sup> Rosiglitazone used as a probe substrate for CYP2C8.
- <sup>o</sup> Mean ratio (with/without coadministered drug). ↑ indicates an increase in rosuvastatin exposure.
- <sup>p</sup> The combination of atazanavir and saquinavir 1200 mg QD produced daily saquinavir exposures similar to the values produced by the standard therapeutic dosing of saquinavir at 1200 mg TID. However, the  $C_{max}$  is about 79% higher than that for the standard dosing of saquinavir (soft gelatin capsules) alone at 1200 mg TID.
- <sup>q</sup> Note that similar results were observed in a study where administration of tenofovir and REYATAZ was separated by 12 hours.
- <sup>r</sup> Administration of tenofovir and REYATAZ was temporally separated by 12 hours.
- NA = not available.

## 12.4 Microbiology

### Mechanism of Action

Atazanavir (ATV) is an azapeptide HIV-1 protease inhibitor (PI). The compound selectively inhibits the virus-specific processing of viral Gag and Gag-Pol polyproteins in HIV-1 infected cells, thus preventing formation of mature virions.

### Antiviral Activity in Cell Culture

Atazanavir exhibits anti-HIV-1 activity with a mean 50% effective concentration ( $EC_{50}$ ) in the absence of human serum of 2 to 5 nM against a variety of laboratory and clinical HIV-1 isolates grown in peripheral blood mononuclear cells, macrophages, CEM-SS cells, and MT-2 cells. ATV has activity against HIV-1 Group M subtype viruses A, B, C, D, AE, AG, F, G, and J isolates in cell culture. ATV has variable activity against HIV-2 isolates (1.9–32 nM), with  $EC_{50}$  values above the  $EC_{50}$  values of failure isolates. Two-drug combination antiviral activity studies with ATV showed no antagonism in cell culture with NNRTIs (delavirdine, efavirenz, and nevirapine), PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir), NRTIs (abacavir, didanosine, emtricitabine, lamivudine, stavudine, tenofovir, zalcitabine, and zidovudine), the HIV-1 fusion inhibitor enfuvirtide, and two compounds used in the treatment of viral hepatitis, adefovir and ribavirin, without enhanced cytotoxicity.

### Resistance

*In Cell Culture:* HIV-1 isolates with a decreased susceptibility to ATV have been selected in cell culture and obtained from patients treated with ATV or atazanavir/ritonavir (ATV/RTV). HIV-1 isolates with 93- to 183-fold reduced susceptibility to ATV from three different viral strains were selected in cell culture by 5 months. The substitutions in these HIV-1 viruses that contributed to

ATV resistance include I50L, N88S, I84V, A71V, and M46I. Changes were also observed at the protease cleavage sites following drug selection. Recombinant viruses containing the I50L substitution without other major PI substitutions were growth impaired and displayed increased susceptibility in cell culture to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir). The I50L and I50V substitutions yielded selective resistance to ATV and amprenavir, respectively, and did not appear to be cross-resistant.

*Clinical Studies of Treatment-Naive Patients: Comparison of Ritonavir-Boosted REYATAZ vs. Unboosted REYATAZ:* Study AI424-089 compared REYATAZ 300 mg once daily with ritonavir 100 mg vs. REYATAZ 400 mg once daily when administered with lamivudine and extended-release stavudine in HIV-infected treatment-naive patients. A summary of the number of virologic failures and virologic failure isolates with ATV resistance in each arm is shown in Table 23.

**Table 23: Summary of Virologic Failures<sup>a</sup> at Week 96 in Study AI424-089: Comparison of Ritonavir Boosted REYATAZ vs. Unboosted REYATAZ: Randomized Patients**

	REYATAZ 300 mg + ritonavir 100 mg (n=95)	REYATAZ 400 mg (n=105)
Virologic Failure (≥50 copies/mL) at Week 96	15 (16%)	34 (32%)
Virologic Failure with Genotypes and Phenotypes Data	5	17
Virologic Failure Isolates with ATV-resistance at Week 96	0/5 (0%) <sup>b</sup>	4/17 (24%) <sup>b</sup>
Virologic Failure Isolates with I50L Emergence at Week 96 <sup>c</sup>	0/5 (0%) <sup>b</sup>	2/17 (12%) <sup>b</sup>
Virologic Failure Isolates with Lamivudine Resistance at Week 96	2/5 (40%) <sup>b</sup>	11/17 (65%) <sup>b</sup>

<sup>a</sup> Virologic failure includes patients who were never suppressed through Week 96 and on study at Week 96, had virologic rebound or discontinued due to insufficient viral load response.

<sup>b</sup> Percentage of Virologic Failure Isolates with genotypic and phenotypic data.

<sup>c</sup> Mixture of I50I/L emerged in 2 other ATV 400 mg-treated patients. Neither isolate was phenotypically resistant to ATV.

*Clinical Studies of Treatment-Naive Patients Receiving REYATAZ 300 mg with Ritonavir 100 mg:* In Phase III study AI424-138, an as-treated genotypic and phenotypic analysis was conducted on samples from patients who experienced virologic failure (HIV-1 RNA ≥400 copies/mL) or discontinued before achieving suppression on ATV/RTV (n=39; 9%) and

LPV/RTV (n=39; 9%) through 96 weeks of treatment. In the ATV/RTV arm, one of the virologic failure isolates had a 56-fold decrease in ATV susceptibility emerge on therapy with the development of PI resistance-associated substitutions L10F, V32I, K43T, M46I, A71I, G73S, I85I/V, and L90M. The NRTI resistance-associated substitution M184V also emerged on treatment in this isolate conferring emtricitabine resistance. Two ATV/RTV-virologic failure isolates had baseline phenotypic ATV resistance and IAS-defined major PI resistance-associated substitutions at baseline. The I50L substitution emerged on study in one of these failure isolates and was associated with a 17-fold decrease in ATV susceptibility from baseline and the other failure isolate with baseline ATV resistance and PI substitutions (M46M/I and I84I/V) had additional IAS-defined major PI substitutions (V32I, M46I, and I84V) emerge on ATV treatment associated with a 3-fold decrease in ATV susceptibility from baseline. Five of the treatment failure isolates in the ATV/RTV arm developed phenotypic emtricitabine resistance with the emergence of either the M184I (n=1) or the M184V (n=4) substitution on therapy and none developed phenotypic tenofovir disoproxil resistance. In the LPV/RTV arm, one of the virologic failure patient isolates had a 69-fold decrease in LPV susceptibility emerge on therapy with the development of PI substitutions L10V, V11I, I54V, G73S, and V82A in addition to baseline PI substitutions L10L/I, V32I, I54I/V, A71I, G73G/S, V82V/A, L89V, and L90M. Six LPV/RTV virologic failure isolates developed the M184V substitution and phenotypic emtricitabine resistance and two developed phenotypic tenofovir disoproxil resistance.

*Clinical Studies of Treatment-Naive Patients Receiving REYATAZ 400 mg without Ritonavir:* ATV-resistant clinical isolates from treatment-naive patients who experienced virologic failure on REYATAZ 400 mg treatment without ritonavir often developed an I50L substitution (after an average of 50 weeks of ATV therapy), often in combination with an A71V substitution, but also developed one or more other PI substitutions (eg, V32I, L33F, G73S, V82A, I85V, or N88S) with or without the I50L substitution. In treatment-naive patients, viral isolates that developed the I50L substitution, without other major PI substitutions, showed phenotypic resistance to ATV but retained in cell culture susceptibility to other PIs (amprenavir, indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir); however, there are no clinical data available to demonstrate the effect of the I50L substitution on the efficacy of subsequently administered PIs.

*Clinical Studies of Treatment-Experienced Patients:* In studies of treatment-experienced patients treated with ATV or ATV/RTV, most ATV-resistant isolates from patients who experienced virologic failure developed substitutions that were associated with resistance to multiple PIs and displayed decreased susceptibility to multiple PIs. The most common protease substitutions to develop in the viral isolates of patients who failed treatment with ATV 300 mg once daily and RTV 100 mg once daily (together with tenofovir and an NRTI) included V32I, L33F/V/I,

E35D/G, M46I/L, I50L, F53L/V, I54V, A71V/T/I, G73S/T/C, V82A/T/L, I85V, and L89V/Q/M/T. Other substitutions that developed on ATV/RTV treatment including E34K/A/Q, G48V, I84V, N88S/D/T, and L90M occurred in less than 10% of patient isolates. Generally, if multiple PI resistance substitutions were present in the HIV-1 virus of the patient at baseline, ATV resistance developed through substitutions associated with resistance to other PIs and could include the development of the I50L substitution. The I50L substitution has been detected in treatment-experienced patients experiencing virologic failure after long-term treatment. Protease cleavage site changes also emerged on ATV treatment but their presence did not correlate with the level of ATV resistance.

*Clinical Studies of Pediatric Subjects in AI424-397 (PRINCE I) and AI424-451 (PRINCE II):* No treatment-emergent ATV-associated substitutions were detected among treatment failures in AI424-397, but four known resistance-associated substitutions to other PIs arose in the viruses from one subject each (L19I/R, M36M/I, H69K/R, and I72I/V). None of these viruses acquired phenotypic resistance to ATV, ATV/RTV, or any NNRTI or NRTI. In AI424-451, ATV-associated resistance substitution (I84V) and other PI substitutions arose in the virus of one subject, including M46M/V, V82V/I, I84I/V, and L90L/M; however, these substitutions did not result in phenotypic resistance to ATV (ATV phenotypic fold change: 1.74, using a commercial investigational assay with an ATV cutoff of 2.2 fold change). Secondary PI substitutions also arose in the viruses of one subject each, including V11V/I, G16G/E, D30D/G, E35E/D, K45K/R, L63P/S, and I72I/T. Q61D and Q61E/G emerged in the viruses of two subjects who failed treatment with ATV/RTV. Viruses from three subjects developed M184V in the reverse transcriptase, and all three exhibited phenotypic resistance to emtricitabine and lamivudine.

## **Cross-Resistance**

Cross-resistance among PIs has been observed. Baseline phenotypic and genotypic analyses of clinical isolates from ATV clinical trials of PI-experienced patients showed that isolates cross-resistant to multiple PIs were cross-resistant to ATV. Greater than 90% of the isolates with substitutions that included I84V or G48V were resistant to ATV. Greater than 60% of isolates containing L90M, G73S/T/C, A71V/T, I54V, M46I/L, or a change at V82 were resistant to ATV, and 38% of isolates containing a D30N substitution in addition to other changes were resistant to ATV. Isolates resistant to ATV were also cross-resistant to other PIs with >90% of the isolates resistant to indinavir, lopinavir, nelfinavir, ritonavir, and saquinavir, and 80% resistant to amprenavir. In treatment-experienced patients, PI-resistant viral isolates that developed the I50L substitution in addition to other PI resistance-associated substitution were also cross-resistant to other PIs.

## Baseline Genotype/Phenotype and Virologic Outcome Analyses

Genotypic and/or phenotypic analysis of baseline virus may aid in determining ATV susceptibility before initiation of ATV/RTV therapy. An association between virologic response at 48 weeks and the number and type of primary PI resistance-associated substitutions detected in baseline HIV-1 isolates from antiretroviral-experienced patients receiving ATV/RTV once daily or lopinavir (LPV)/RTV twice daily in Study AI424-045 is shown in Table 24.

Overall, both the number and type of baseline PI substitutions affected response rates in treatment-experienced patients. In the ATV/RTV group, patients had lower response rates when 3 or more baseline PI substitutions, including a substitution at position 36, 71, 77, 82, or 90, were present compared to patients with 1–2 PI substitutions, including one of these substitutions.

**Table 24: HIV RNA Response by Number and Type of Baseline PI Substitution, Antiretroviral-Experienced Patients in Study AI424-045, As-Treated Analysis**

Number and Type of Baseline PI Substitutions <sup>a</sup>	Virologic Response = HIV RNA <400 copies/mL <sup>b</sup>	
	ATV/RTV (n=110)	LPV/RTV (n=113)
<b>3 or more primary PI substitutions including<sup>c</sup>:</b>		
D30N	75% (6/8)	50% (3/6)
M36I/V	19% (3/16)	33% (6/18)
M46I/L/T	24% (4/17)	23% (5/22)
I54V/L/T/M/A	31% (5/16)	31% (5/16)
A71V/T/I/G	34% (10/29)	39% (12/31)
G73S/A/C/T	14% (1/7)	38% (3/8)
V77I	47% (7/15)	44% (7/16)
V82A/F/T/S/I	29% (6/21)	27% (7/26)
I84V/A	11% (1/9)	33% (2/6)
N88D	63% (5/8)	67% (4/6)
L90M	10% (2/21)	44% (11/25)



**Table 24: HIV RNA Response by Number and Type of Baseline PI Substitution, Antiretroviral-Experienced Patients in Study AI424-045, As-Treated Analysis**

Number and Type of Baseline PI Substitutions <sup>a</sup>	Virologic Response = HIV RNA <400 copies/mL <sup>b</sup>	
	ATV/RTV (n=110)	LPV/RTV (n=113)
<b>Number of baseline primary PI substitutions<sup>a</sup></b>		
All patients, as-treated	58% (64/110)	59% (67/113)
0–2 PI substitutions	75% (50/67)	75% (50/67)
3–4 PI substitutions	41% (14/34)	43% (12/28)
5 or more PI substitutions	0% (0/9)	28% (5/18)

<sup>a</sup> Primary substitutions include any change at D30, V32, M36, M46, I47, G48, I50, I54, A71, G73, V77, V82, I84, N88, and L90.

<sup>b</sup> Results should be interpreted with caution because the subgroups were small.

<sup>c</sup> There were insufficient data (n<3) for PI substitutions V32I, I47V, G48V, I50V, and F53L.

The response rates of antiretroviral-experienced patients in Study AI424-045 were analyzed by baseline phenotype (shift in susceptibility in cell culture relative to reference, Table 25). The analyses are based on a select patient population with 62% of patients receiving an NNRTI-based regimen before study entry compared to 35% receiving a PI-based regimen. Additional data are needed to determine clinically relevant break points for REYATAZ.

**Table 25: Baseline Phenotype by Outcome, Antiretroviral-Experienced Patients in Study AI424-045, As-Treated Analysis**

Baseline Phenotype <sup>a</sup>	Virologic Response = HIV RNA <400 copies/mL <sup>b</sup>	
	ATV/RTV (n=111)	LPV/RTV (n=111)
0–2	71% (55/78)	70% (56/80)
>2–5	53% (8/15)	44% (4/9)
>5–10	13% (1/8)	33% (3/9)
>10	10% (1/10)	23% (3/13)

<sup>a</sup> Fold change susceptibility in cell culture relative to the wild-type reference.

<sup>b</sup> Results should be interpreted with caution because the subgroups were small.



## **13 NONCLINICAL TOXICOLOGY**

### **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

#### **Carcinogenesis**

Long-term carcinogenicity studies in mice and rats were carried out with atazanavir for two years. In the mouse study, drug-related increases in hepatocellular adenomas were found in females at 360 mg/kg/day. The systemic drug exposure (AUC) at the NOAEL (no observable adverse effect level) in females, (120 mg/kg/day) was 2.8 times and in males (80 mg/kg/day) was 2.9 times higher than those in humans at the clinical dose (300 mg/day atazanavir boosted with 100 mg/day ritonavir, non-pregnant patients). In the rat study, no drug-related increases in tumor incidence were observed at doses up to 1200 mg/kg/day, for which AUCs were 1.1 (males) or 3.9 (females) times those measured in humans at the clinical dose.

#### **Mutagenesis**

Atazanavir tested positive in an *in vitro* clastogenicity test using primary human lymphocytes, in the absence and presence of metabolic activation. Atazanavir tested negative in the *in vitro* Ames reverse-mutation assay, *in vivo* micronucleus and DNA repair tests in rats, and *in vivo* DNA damage test in rat duodenum (comet assay).

#### **Impairment of Fertility**

At the systemic drug exposure levels (AUC) 0.9 (in male rats) or 2.3 (in female rats) times that of the human clinical dose, (300 mg/day atazanavir boosted with 100 mg/day ritonavir) significant effects on mating, fertility, or early embryonic development were not observed.

## **14 CLINICAL STUDIES**

### **14.1 Adult Patients without Prior Antiretroviral Therapy**

*Study AI424-138: a 96-week study comparing the antiviral efficacy and safety of REYATAZ/ritonavir with lopinavir/ritonavir, each in combination with fixed-dose tenofovir-emtricitabine in HIV-1 infected treatment-naïve subjects.* Study AI424-138 was a 96-week, open-label, randomized, multicenter study, comparing REYATAZ (300 mg once daily) with ritonavir (100 mg once daily) to lopinavir with ritonavir (400/100 mg twice daily), each in combination with fixed-dose tenofovir with emtricitabine (300/200 mg once daily), in 878 antiretroviral treatment-naïve treated patients. Patients had a mean age of 36 years (range: 19-72), 49% were Caucasian, 18% Black, 9% Asian, 23% Hispanic/Mestizo/mixed race, and

68% were male. The median baseline plasma CD4+ cell count was 204 cells/mm<sup>3</sup> (range: 2 to 810 cells/mm<sup>3</sup>) and the mean baseline plasma HIV-1 RNA level was 4.94 log<sub>10</sub> copies/mL (range: 2.60 to 5.88 log<sub>10</sub> copies/mL). Treatment response and outcomes through Week 96 are presented in Table 26.

**Table 26: Outcomes of Treatment Through Week 96 in Treatment-Naive Adults (Study AI424-138)**

Outcome	REYATAZ 300 mg + ritonavir 100 mg (once daily) with tenofovir/emtricitabine (once daily) <sup>a</sup> (n=441)	lopinavir 400 mg + ritonavir 100 mg (twice daily) with tenofovir/emtricitabine (once daily) <sup>a</sup> (n=437)
	96 Weeks	96 Weeks
Responder <sup>b,c,d</sup>	75%	68%
Virologic failure <sup>e</sup>	17%	19%
Rebound	8%	10%
Never suppressed through Week 96	9%	9%
Death	1%	1%
Discontinued due to adverse event	3%	5%
Discontinued for other reasons <sup>f</sup>	4%	7%

<sup>a</sup> As a fixed-dose combination: 300 mg tenofovir, 200 mg emtricitabine once daily.

<sup>b</sup> Patients achieved HIV RNA <50 copies/mL at Week 96. Roche Amplicor®, v1.5 ultra-sensitive assay.

<sup>c</sup> Pre-specified ITT analysis at Week 48 using as-randomized cohort: ATV/RTV 78% and LPV/RTV 76% (difference estimate: 1.7% [95% confidence interval: -3.8%, 7.1%]).

<sup>d</sup> Pre-specified ITT analysis at Week 96 using as-randomized cohort: ATV/RTV 74% and LPV/RTV 68% (difference estimate: 6.1% [95% confidence interval: 0.3%, 12.0%]).

<sup>e</sup> Includes viral rebound and failure to achieve confirmed HIV RNA <50 copies/mL through Week 96.

<sup>f</sup> Includes lost to follow-up, patient's withdrawal, noncompliance, protocol violation, and other reasons.

Through 96 weeks of therapy, the proportion of responders among patients with high viral loads (ie, baseline HIV RNA ≥100,000 copies/mL) was comparable for the REYATAZ/ritonavir (165 of 223 patients, 74%) and lopinavir/ritonavir (148 of 222 patients, 67%) arms. At 96 weeks, the median increase from baseline in CD4+ cell count was 261 cells/mm<sup>3</sup> for the REYATAZ/ritonavir arm and 273 cells/mm<sup>3</sup> for the lopinavir/ritonavir arm.

*Study AI424-034: REYATAZ once daily compared to efavirenz once daily, each in combination with fixed-dose lamivudine + zidovudine twice daily.* Study AI424-034 was a randomized, double-blind, multicenter trial comparing REYATAZ (400 mg once daily) to efavirenz (600 mg

once daily), each in combination with a fixed-dose combination of lamivudine (3TC) (150 mg) and zidovudine (ZDV) (300 mg) given twice daily, in 810 antiretroviral treatment-naïve patients. Patients had a mean age of 34 years (range: 18 to 73), 36% were Hispanic, 33% were Caucasian, and 65% were male. The mean baseline CD4+ cell count was 321 cells/mm<sup>3</sup> (range: 64 to 1424 cells/mm<sup>3</sup>) and the mean baseline plasma HIV-1 RNA level was 4.8 log<sub>10</sub> copies/mL (range: 2.2 to 5.9 log<sub>10</sub> copies/mL). Treatment response and outcomes through Week 48 are presented in Table 27.

**Table 27: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naïve Adults (Study AI424-034)**

Outcome	REYATAZ 400 mg once daily + lamivudine + zidovudine <sup>d</sup> (n=405)	efavirenz 600 mg once daily + lamivudine + zidovudine <sup>d</sup> (n=405)
Responder <sup>a</sup>	67% (32%)	62% (37%)
Virologic failure <sup>b</sup>	20%	21%
Rebound	17%	16%
Never suppressed through Week 48	3%	5%
Death	—	<1%
Discontinued due to adverse event	5%	7%
Discontinued for other reasons <sup>c</sup>	8%	10%

<sup>a</sup> Patients achieved and maintained confirmed HIV RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.0 or 1.5 as geographically appropriate.

<sup>b</sup> Includes viral rebound and failure to achieve confirmed HIV RNA <400 copies/mL through Week 48.

<sup>c</sup> Includes lost to follow-up, patient's withdrawal, noncompliance, protocol violation, and other reasons.

<sup>d</sup> As a fixed-dose combination: 150 mg lamivudine, 300 mg zidovudine twice daily.

Through 48 weeks of therapy, the proportion of responders among patients with high viral loads (ie, baseline HIV RNA ≥100,000 copies/mL) was comparable for the REYATAZ and efavirenz arms. The mean increase from baseline in CD4+ cell count was 176 cells/mm<sup>3</sup> for the REYATAZ arm and 160 cells/mm<sup>3</sup> for the efavirenz arm.

*Study AI424-008: REYATAZ 400 mg once daily compared to REYATAZ 600 mg once daily, and compared to nelfinavir 1250 mg twice daily, each in combination with stavudine and lamivudine twice daily.* Study AI424-008 was a 48-week, randomized, multicenter trial, blinded to dose of REYATAZ, comparing REYATAZ at two dose levels (400 mg and 600 mg once daily) to

nelfinavir (1250 mg twice daily), each in combination with stavudine (40 mg) and lamivudine (150 mg) given twice daily, in 467 antiretroviral treatment-naïve patients. Patients had a mean age of 35 years (range: 18 to 69), 55% were Caucasian, and 63% were male. The mean baseline CD4+ cell count was 295 cells/mm<sup>3</sup> (range: 4 to 1003 cells/mm<sup>3</sup>) and the mean baseline plasma HIV-1 RNA level was 4.7 log<sub>10</sub> copies/mL (range: 1.8 to 5.9 log<sub>10</sub> copies/mL). Treatment response and outcomes through Week 48 are presented in Table 28.

**Table 28: Outcomes of Randomized Treatment Through Week 48 in Treatment-Naïve Adults (Study AI424-008)**

<b>Outcome</b>	<b>REYATAZ 400 mg once daily + lamivudine + stavudine (n=181)</b>	<b>nelfinavir 1250 mg twice daily + lamivudine + stavudine (n=91)</b>
Responder <sup>a</sup>	67% (33%)	59% (38%)
Virologic failure <sup>b</sup>	24%	27%
Rebound	14%	14%
Never suppressed through Week 48	10%	13%
Death	<1%	–
Discontinued due to adverse event	1%	3%
Discontinued for other reasons <sup>c</sup>	7%	10%

<sup>a</sup> Patients achieved and maintained confirmed HIV RNA <400 copies/mL (<50 copies/mL) through Week 48. Roche Amplicor<sup>®</sup> HIV-1 Monitor<sup>™</sup> Assay, test version 1.0 or 1.5 as geographically appropriate.

<sup>b</sup> Includes viral rebound and failure to achieve confirmed HIV RNA <400 copies/mL through Week 48.

<sup>c</sup> Includes lost to follow-up, patient's withdrawal, noncompliance, protocol violation, and other reasons.

Through 48 weeks of therapy, the mean increase from baseline in CD4+ cell count was 234 cells/mm<sup>3</sup> for the REYATAZ 400-mg arm and 211 cells/mm<sup>3</sup> for the nelfinavir arm.

## 14.2 Adult Patients with Prior Antiretroviral Therapy

*Study AI424-045: REYATAZ once daily + ritonavir once daily compared to REYATAZ once daily + saquinavir (soft gelatin capsules) once daily, and compared to lopinavir + ritonavir twice daily, each in combination with tenofovir + one NRTI.* Study AI424-045 was a randomized, multicenter trial comparing REYATAZ (300 mg once daily) with ritonavir (100 mg once daily) to REYATAZ (400 mg once daily) with saquinavir soft gelatin capsules (1200 mg once daily), and to lopinavir + ritonavir (400/100 mg twice daily), each in combination with tenofovir and one NRTI, in 347 (of 358 randomized) patients who experienced virologic failure on HAART

regimens containing PIs, NNRTIs, and NRTIs. The mean time of prior exposure to antiretrovirals was 139 weeks for PIs, 85 weeks for NNRTIs, and 283 weeks for NRTIs. The mean age was 41 years (range: 24 to 74); 60% were Caucasian, and 78% were male. The mean baseline CD4+ cell count was 338 cells/mm<sup>3</sup> (range: 14 to 1543 cells/mm<sup>3</sup>) and the mean baseline plasma HIV-1 RNA level was 4.4 log<sub>10</sub> copies/mL (range: 2.6 to 5.88 log<sub>10</sub> copies/mL).

Treatment outcomes through Week 48 for the REYATAZ/ritonavir and lopinavir/ritonavir treatment arms are presented in Table 29. REYATAZ/ritonavir and lopinavir/ritonavir were similar for the primary efficacy outcome measure of time-averaged difference in change from baseline in HIV RNA level. Study AI424-045 was not large enough to reach a definitive conclusion that REYATAZ/ritonavir and lopinavir/ritonavir are equivalent on the secondary efficacy outcome measure of proportions below the HIV RNA lower limit of quantification. [See Microbiology, Tables 24 and 25 (12.4).]

**Table 29: Outcomes of Treatment Through Week 48 in Study AI424-045 (Patients with Prior Antiretroviral Experience)**

Outcome	REYATAZ 300 mg + ritonavir 100 mg once daily + tenofovir + 1 NRTI (n=119)	lopinavir/ritonavir (400/100 mg) twice daily + tenofovir + 1 NRTI (n=118)	Difference <sup>a</sup> (REYATAZ- lopinavir/ritonavir) (CI)
HIV RNA Change from Baseline (log <sub>10</sub> copies/mL) <sup>b</sup>	-1.58	-1.70	+0.12 <sup>c</sup> (-0.17, 0.41)
CD4+ Change from Baseline (cells/mm <sup>3</sup> ) <sup>d</sup>	116	123	-7 (-67, 52)
Percent of Patients Responding <sup>e</sup>			
HIV RNA <400 copies/mL <sup>b</sup>	55%	57%	-2.2% (-14.8%, 10.5%)
HIV RNA <50 copies/mL <sup>b</sup>	38%	45%	-7.1% (-19.6%, 5.4%)

<sup>a</sup> Time-averaged difference through Week 48 for HIV RNA; Week 48 difference in HIV RNA percentages and CD4+ mean changes, REYATAZ/ritonavir vs lopinavir/ritonavir; CI = 97.5% confidence interval for change in HIV RNA; 95% confidence interval otherwise.

<sup>b</sup> Roche Amplicor® HIV-1 Monitor™ Assay, test version 1.5.

<sup>c</sup> Protocol-defined primary efficacy outcome measure.

<sup>d</sup> Based on patients with baseline and Week 48 CD4+ cell count measurements (REYATAZ/ritonavir, n=85; lopinavir/ritonavir, n=93).

<sup>e</sup> Patients achieved and maintained confirmed HIV-1 RNA <400 copies/mL (<50 copies/mL) through Week 48.

No patients in the REYATAZ/ritonavir treatment arm and three patients in the lopinavir/ritonavir treatment arm experienced a new-onset CDC Category C event during the study.

In Study AI424-045, the mean change from baseline in plasma HIV-1 RNA for REYATAZ 400 mg with saquinavir (n=115) was  $-1.55 \log_{10}$  copies/mL, and the time-averaged difference in change in HIV-1 RNA levels versus lopinavir/ritonavir was 0.33. The corresponding mean increase in CD4+ cell count was 72 cells/mm<sup>3</sup>. Through 48 weeks of treatment, the proportion of patients in this treatment arm with plasma HIV-1 RNA <400 (<50) copies/mL was 38% (26%). In this study, coadministration of REYATAZ and saquinavir did not provide adequate efficacy [see *Drug Interactions* (7)].

Study AI424-045 also compared changes from baseline in lipid values. [See *Adverse Reactions* (6.1).]

*Study AI424-043:* Study AI424-043 was a randomized, open-label, multicenter trial comparing REYATAZ (400 mg once daily) to lopinavir/ritonavir (400/100 mg twice daily), each in combination with two NRTIs, in 300 patients who experienced virologic failure to only one prior PI-containing regimen. Through 48 weeks, the proportion of patients with plasma HIV-1 RNA <400 (<50) copies/mL was 49% (35%) for patients randomized to REYATAZ (n=144) and 69% (53%) for patients randomized to lopinavir/ritonavir (n=146). The mean change from baseline was  $-1.59 \log_{10}$  copies/mL in the REYATAZ treatment arm and  $-2.02 \log_{10}$  copies/mL in the lopinavir/ritonavir arm. Based on the results of this study, REYATAZ without ritonavir was inferior to lopinavir/ritonavir in PI-experienced patients with prior virologic failure and is not recommended for such patients.

## 14.3 Pediatric Patients

### Pediatric Trials with REYATAZ Capsules

Assessment of the pharmacokinetics, safety, tolerability, and virologic response of REYATAZ capsules was based on data from the open-label, multicenter clinical trial PACTG 1020A which included patients from 6 years to 21 years of age. In this study, 105 patients (43 antiretroviral-naïve and 62 antiretroviral-experienced) received once daily REYATAZ capsule formulation, with or without ritonavir, in combination with two NRTIs.

One-hundred five (105) patients (6 to less than 18 years of age) treated with the REYATAZ capsule formulation, with or without ritonavir, were evaluated. Using an ITT analysis, the overall proportions of antiretroviral-naïve and -experienced patients with HIV RNA <400 copies/mL at Week 96 were 51% (22/43) and 34% (21/62), respectively. The overall proportions of

antiretroviral-naïve and -experienced patients with HIV RNA <50 copies/mL at Week 96 were 47% (20/43) and 24% (15/62), respectively. The median increase from baseline in absolute CD4 count at 96 weeks of therapy was 335 cells/mm<sup>3</sup> in antiretroviral-naïve patients and 220 cells/mm<sup>3</sup> in antiretroviral-experienced patients.

### **Pediatric Trials with REYATAZ Oral Powder**

Assessment of the pharmacokinetics, safety, tolerability, and virologic response of REYATAZ oral powder was based on data from two open-label, multicenter clinical trials.

- AI424-397 (PRINCE I): In pediatric patients from 3 months to less than 6 years of age
- AI424-451 (PRINCE II): In pediatric patients from 3 months to less than 11 years of age

In these studies 134 patients (73 antiretroviral-naïve and 61 antiretroviral-experienced) received once daily REYATAZ oral powder and ritonavir, in combination with two NRTIs.

For inclusion in both trials, treatment-naïve patients had to have genotypic sensitivity to REYATAZ and two NRTIs, and treatment-experienced patients had to have documented genotypic and phenotypic sensitivity at screening to REYATAZ and at least 2 NRTIs. Patients exposed only to antiretrovirals *in utero* or intrapartum were considered treatment-naïve. Patients who received REYATAZ or REYATAZ/ritonavir at any time prior to study enrollment or who had a history of treatment failure on two or more protease inhibitors were excluded from the trials.

Sixty-five (65) patients from both studies weighing 10 kg to less than 25 kg treated with REYATAZ oral powder with ritonavir were evaluated. Patients 10 kg to less than 15 kg received 200 mg REYATAZ and 80 mg ritonavir oral solution, and patients 15 kg to less than 25 kg received 250 mg REYATAZ and 80 mg ritonavir oral solution. Using a modified ITT analysis, the overall proportions of antiretroviral-naïve and antiretroviral-experienced patients with HIV RNA <400 copies/mL at Week 48 were 78% (32/41) and 71% (17/24), respectively in patients who received REYATAZ oral powder with ritonavir. The overall proportions of antiretroviral-naïve and antiretroviral-experienced patients with HIV RNA <50 copies/mL at Week 48 were 66% (27/41) and 58% (14/24), respectively in patients who received REYATAZ oral powder with ritonavir. The median increase from baseline in absolute CD4 count (percent) at 48 weeks of therapy was 412 cells/mm<sup>3</sup> (10.5%) in antiretroviral-naïve patients and 228 cells/mm<sup>3</sup> (6%) in antiretroviral-experienced patients who received REYATAZ oral powder with ritonavir.



## 16 HOW SUPPLIED/STORAGE AND HANDLING

### REYATAZ Capsules

REYATAZ<sup>®</sup> (atazanavir) capsules are available in the following strengths and configurations of plastic bottles with child-resistant closures.

Product Strength*	Capsule Shell Color (cap/body)	Markings on Capsule (ink color)		Capsules per Bottle	NDC Number
		cap	body		
150 mg	blue/powder blue	BMS 150 mg (white)	3624 (blue)	60	0003-3624-12
200 mg	blue/blue	BMS 200 mg (white)	3631 (white)	60	0003-3631-12
300 mg	red/blue	BMS 300 mg (white)	3622 (white)	30	0003-3622-12

\* 150 mg atazanavir equivalent to 170.8 mg atazanavir sulfate.

200 mg atazanavir equivalent to 227.8 mg atazanavir sulfate.

300 mg atazanavir equivalent to 341.69 mg atazanavir sulfate.

Store REYATAZ capsules at 25°C (77°F); excursions permitted to 15°C-30°C (59°F-86°F) [see USP Controlled Room Temperature].

### REYATAZ Oral Powder

REYATAZ oral powder is an orange-vanilla flavored powder, packed in child-resistant packets. Each packet contains 50 mg of atazanavir equivalent to 56.9 mg of atazanavir sulfate in 1.5 g of powder. REYATAZ oral powder is supplied in cartons (NDC 0003-3638-10) of 30 packets each. [See *Dosage and Administration* (2.4).]

Store REYATAZ oral powder below 30°C (86°F). Once the REYATAZ oral powder is mixed with food or beverage, it may be kept at room temperature 20°C to 30°C (68°F-86°F) for up to 1 hour prior to administration. Store REYATAZ oral powder in the original packet and do not open until ready to use.

## 17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling (*Patient Information and Instructions for Use*).

A statement to patients and healthcare providers is included on the product's label:

**ALERT: Find out about medicines that should NOT be taken with REYATAZ<sup>®</sup>.**



REYATAZ is not a cure for HIV infection and patients may continue to experience illnesses associated with HIV infection, including opportunistic infections. Patients should remain under the care of a healthcare provider when using REYATAZ.

Patients should be advised to avoid doing things that can spread HIV infection to others.

- **Do not share or reuse needles or other injection equipment.**
- **Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.**
- **Do not have any kind of sex without protection.** Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.
- **Do not breastfeed.** It is not known if REYATAZ can be passed to your baby in your breast milk and whether it could harm your baby. Also, mothers with HIV should not breastfeed because HIV can be passed to the baby in breast milk.

## Dosing Instructions

Patients should be told that sustained decreases in plasma HIV RNA have been associated with a reduced risk of progression to AIDS and death. Patients should remain under the care of a healthcare provider while using REYATAZ. Patients should be advised to take REYATAZ with food every day and take other concomitant antiretroviral therapy as prescribed. REYATAZ must always be used in combination with other antiretroviral drugs. Patients should not alter the dose or discontinue therapy without consulting with their healthcare provider. If a dose of REYATAZ is missed, patients should take the dose as soon as possible and then return to their normal schedule. However, if a dose is skipped the patient should not double the next dose.

REYATAZ oral powder is available for pediatric patients who are 3 months and older weighing 10 kg to less than 25 kg. Caregivers should be advised on how to mix the REYATAZ oral powder with a food or beverage such as milk or water for infants and young children who can take solid foods or drink liquids from a cup. For infants who cannot take solid food or drink from a cup, the powder formulation mixed in liquid infant formula should be given with an oral dosing syringe. Caregivers should carefully follow the *Instructions for Use* and storage of the powder [see *Dosage and Administration (2.4)* and *FDA-approved patient labeling (Patient Information and Instructions for Use)*].

Caregivers of patients with phenylketonuria should be advised that REYATAZ oral powder contains phenylalanine.

Patients or caregivers should call their healthcare provider or pharmacist if they have any questions.

## Drug Interactions

REYATAZ may interact with some drugs; therefore, patients should be advised to report to their healthcare provider the use of any other prescription, nonprescription medication, or herbal products, particularly St. John's wort.

Patients receiving a PDE5 inhibitor and atazanavir should be advised that they may be at an increased risk of PDE5 inhibitor-associated adverse events including hypotension, syncope, visual disturbances, and priapism, and should promptly report any symptoms to their doctor.

Patients should be informed that REVATIO<sup>®</sup> (used to treat pulmonary arterial hypertension) is contraindicated with REYATAZ and that dose adjustments are necessary when REYATAZ is used with CIALIS<sup>®</sup>, LEVITRA<sup>®</sup>, or VIAGRA<sup>®</sup> (used to treat erectile dysfunction), or ADCIRCA<sup>®</sup> (used to treat pulmonary arterial hypertension).

## Cardiac Conduction Abnormalities

Patients should be informed that atazanavir may produce changes in the electrocardiogram (eg, PR prolongation). Patients should consult their healthcare provider if they are experiencing symptoms such as dizziness or lightheadedness.

## Rash

Patients should be informed that mild rashes without other symptoms have been reported with REYATAZ use. These rashes go away within two weeks with no change in treatment. However, there have been reports of severe skin reactions (eg, Stevens-Johnson syndrome, erythema multiforme, and toxic skin eruptions) with REYATAZ use. Patients developing signs or symptoms of severe skin reactions or hypersensitivity reactions (including, but not limited to, severe rash or rash accompanied by one or more of the following: fever, general malaise, muscle or joint aches, blisters, oral lesions, conjunctivitis, facial edema, hepatitis, eosinophilia, granulocytopenia, lymphadenopathy, and renal dysfunction) must discontinue REYATAZ and seek medical evaluation immediately.

## Hyperbilirubinemia

Patients should be informed that asymptomatic elevations in indirect bilirubin have occurred in patients receiving REYATAZ. This may be accompanied by yellowing of the skin or whites of the eyes and alternative antiretroviral therapy may be considered if the patient has cosmetic concerns.

## **Nephrolithiasis and Cholelithiasis**

Patients should be informed that kidney stones and/or gallstones have been reported with REYATAZ use. Some patients with kidney stones and/or gallstones required hospitalization for additional management and some had complications. Discontinuation of REYATAZ may be necessary as part of the medical management of these adverse events.

## **Fat Redistribution**

Patients should be informed that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy including protease inhibitors and that the cause and long-term health effects of these conditions are not known at this time.

## PATIENT INFORMATION

**REYATAZ<sup>®</sup> (RAY-ah-taz)  
(atazanavir)  
capsules**

**REYATAZ<sup>®</sup> (RAY-ah-taz)  
(atazanavir)  
oral powder**

**Important: Ask your healthcare provider or pharmacist about medicines that should not be taken with REYATAZ.** For more information, see “Who should not take REYATAZ?” and “What should I tell my healthcare provider before taking REYATAZ?”

Read this Patient Information before you start taking REYATAZ and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

### **What is REYATAZ?**

REYATAZ is a prescription HIV-1 (Human Immunodeficiency Virus) medicine that is used with other antiretroviral medicines to treat HIV-1 infection in adults and children 3 months of age and older and who weigh at least 22 pounds (10 kg).

HIV is the virus that causes AIDS (Acquired Immunodeficiency Syndrome).

REYATAZ should not be used in children younger than 3 months of age.

When used with other antiretroviral medicines to treat HIV infection, REYATAZ may help:

- reduce the amount of HIV in your blood. This is called viral load.
- increase the number of CD4+ (T) cells in your blood that help fight off other infections.

Reducing the amount of HIV and increasing the CD4+ (T) cells in your blood may help improve your immune system. This may reduce your risk of death or getting infections that can happen when your immune system is weak (opportunistic infections).

**REYATAZ does not cure HIV infection or AIDS. You must stay on continuous HIV therapy to control HIV infection and decrease HIV-related illnesses.**

**Avoid doing things that can spread HIV infection to others:**

- Do not share or reuse needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with any body fluids such as semen, vaginal secretions, or blood.

Ask your healthcare provider if you have any questions about how to prevent passing HIV to other people.

**Who should not take REYATAZ?**

**Do not take REYATAZ if you:**

- are allergic to atazanavir or any of the ingredients in REYATAZ. See the end of this leaflet for a complete list of ingredients in REYATAZ.
- are taking any of the following medicines. Taking REYATAZ with these medicines may affect how REYATAZ works. REYATAZ may cause serious life-threatening side effects or death when used with these medicines:
  - alfuzosin (UROXATRAL<sup>®</sup>)
  - cisapride (PROPULSID<sup>®</sup>)
  - ergot medicines including:
    - ergotamine tartrate (CAFERGOT<sup>®</sup>, MIGERGOT<sup>®</sup>, ERGOMAR<sup>®</sup>, ERGOSTAT<sup>®</sup>, MEDIHALER<sup>®</sup>, Ergotamine, WIGRAINE<sup>®</sup>, WIGRETTES<sup>®</sup>)
    - dihydroergotamine mesylate (D.H.E. 45<sup>®</sup>, MIGRANAL<sup>®</sup>)
    - methylergonovine (METHERGINE<sup>®</sup>)
  - indinavir (CRIXIVAN<sup>®</sup>)
  - irinotecan (CAMPTOSAR<sup>®</sup>)
  - lovastatin (ADVICOR<sup>®</sup>, ALTOPREV<sup>®</sup>, MEVACOR<sup>®</sup>)
  - midazolam (VERSED<sup>®</sup>), when taken by mouth for sedation
  - nevirapine (VIRAMUNE<sup>®</sup>, VIRAMUNE XR<sup>®</sup>)
  - pimozide (ORAP<sup>®</sup>)
  - rifampin (RIMACTANE<sup>®</sup>, RIFADIN<sup>®</sup>, RIFATER<sup>®</sup>, RIFAMATE<sup>®</sup>)

- sildenafil (REVATIO<sup>®</sup>), when used for the treatment of pulmonary arterial hypertension
- simvastatin (ZOCOR<sup>®</sup>, VYTORIN, SIMCOR)
- St. John's wort (*Hypericum perforatum*)
- triazolam (HALCION<sup>®</sup>)

Serious problems can happen if you or your child take any of the medicines listed above with REYATAZ.

### **What should I tell my healthcare provider before taking REYATAZ?**

#### **Before taking REYATAZ, tell your healthcare provider if you:**

- have heart problems
- have liver problems, including hepatitis B or C virus infection
- have phenylketonuria (PKU). REYATAZ oral powder contains phenylalanine as part of the artificial sweetener aspartame. The artificial sweetener can be harmful to people with PKU.
- are receiving dialysis treatment
- have diabetes
- have hemophilia
- have any other medical conditions
- are pregnant or plan to become pregnant. It is not known if REYATAZ will harm your unborn baby. Pregnant women have developed a serious condition called lactic acidosis (a build-up of lactic acid in the blood) when taking REYATAZ with other HIV medicines called nucleoside analogues.
  - **Hormonal forms of birth control, such as injections, vaginal rings or implants, contraceptive patch, and some birth control pills may not work during treatment with REYATAZ.** Talk to your healthcare provider about forms of birth control that may be used during treatment with REYATAZ.
  - **Pregnancy Registry.** There is a pregnancy registry for women who take antiviral medicines during pregnancy. The purpose of this registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.
  - **After your baby is born,** tell your healthcare provider if your baby's skin or the white part of his/her eyes turns yellow.
- are breastfeeding or plan to breastfeed. Do not breastfeed if you are taking REYATAZ. You should not breastfeed if you have HIV because of the risk of passing HIV to your baby. It is not known if REYATAZ passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby.

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

REYATAZ may affect the way other medicines work, and other medicines may affect how REYATAZ works and may cause serious side effects. If you take certain medicines with REYATAZ, the amount of REYATAZ in your body may be too low and it may not work to help control your HIV infection. The HIV virus in your body may become resistant to REYATAZ or other HIV medicines that are like it. Especially tell your healthcare provider if you take any of the medicines below:

**The following medicines should not be taken during your treatment with REYATAZ:**

- boceprevir (VICTRELIS®)
- salmeterol (SEREVENT DISKUS®) and salmeterol with fluticasone (ADVAIR DISKUS®, ADVAIR HFA®)
- voriconazole (VFEND®)

**If you take any of the following medicines during treatment with REYATAZ, your healthcare provider may need to change the dose of the medicine or change the dose of REYATAZ, or your healthcare provider may need to monitor you more closely:**

- medicines used to treat abnormal heart rhythm:
  - amiodarone (CORDARONE®, NEXTERONE®, PACERONE®)
  - lidocaine
  - quinidine
- allergy or asthma medicines (nasal spray or inhaled treatment):
  - fluticasone propionate (FLONASE®, FLOVENT DISKUS®, FLOVENT HFA®)
- an antibiotic used to treat tuberculosis (TB):
  - rifabutin (MYCOBUTIN®)
- the anticoagulant (blood thinner) medicine:
  - warfarin (COUMADIN®, JANTOVEN®)
- antidepressant medicines:
  - amitriptyline
  - desipramine (NORPRAMIN®)
  - doxepin (SILENOR®)
  - trazodone (OLEPTRO®)
  - trimipramine (SURMONTIL®)
  - imipramine (TOFRANIL®, TOFRANIL-PM®)
  - protriptyline (VIVACTIL®)

- the antipsychotic medicine:
  - quetiapine (SEROQUEL®)
- the chest pain medicine:
  - bepridil (VASCOR®)
- cholesterol-lowering medicines:
  - atorvastatin (LIPITOR®)
  - rosuvastatin (CRESTOR®)
- erectile dysfunction medicines:
  - sildenafil (VIAGRA®)
  - tadalafil (CIALIS®)
  - vardenafil (LEVITRA®, STAXYN®)
- a medicine for gout or treatment of familial Mediterranean fever:
  - colchicine (COLCRYS®)
- organ transplant rejection medicines:
  - cyclosporine (SANDIMMUNE®, NEORAL®, GENGRAF®)
  - sirolimus (RAPAMUNE®)
  - tacrolimus (PROGRAF®, ASTAGRAF XL®)
- pain and addiction medicines:
  - buprenorphine or buprenorphine/naloxone (BUPRENEX®, BUTRANS®, ZUBSOLV®, SUBOXONE®)
- pulmonary arterial hypertension medicines:
  - bosentan (TRACLEER®)
  - tadalafil (ADCIRCA®)
- medicine to treat fungal infections
  - ketoconazole (NIZORAL®)
  - itraconazole (SPORANOX®, ONMEL®)

People who take sildenafil (VIAGRA®), tadalafil (CIALIS®, ADCIRCA®) or vardenafil (LEVITRA®, STAXYN®) with REYATAZ may have a higher risk of certain side effects. Tell your healthcare provider right away if you take any of these medicines and have any of the following side effects:

- lightheadedness, especially when standing
- fainting
- changes in your vision
- an erection that lasts more than 4 hours

**If you take any of the following medicines, your healthcare provider may change the dose or the time you take REYATAZ or the other medicine:**



- antiepileptic medicines:
  - carbamazepine (CARBATROL<sup>®</sup>, EPITOL<sup>®</sup>, TEGRETOL<sup>®</sup>, TEGRETOL-XR<sup>®</sup>, EQUETRO<sup>®</sup>, TERIL<sup>®</sup>)
  - phenytoin (DILANTIN<sup>®</sup>)
  - lamotrigine (LAMICTAL<sup>®</sup>, LAMICTAL CD<sup>®</sup>, LAMICTAL XR<sup>®</sup>, LAMICTAL ODT<sup>®</sup>)
- calcium channel blockers such as:
  - diltiazem (CARDIZEM<sup>®</sup>, CARDIZEM CD<sup>®</sup>, CARDIZEM LA<sup>®</sup>, TIAZAC<sup>®</sup>, CARTIA XT<sup>®</sup>, DILACOR XR<sup>®</sup>, DILT-CD<sup>®</sup>, DILTZAC<sup>®</sup>, TAZTIA XT<sup>®</sup>)
  - verapamil (COVERA-HS<sup>®</sup>, CALAN<sup>®</sup>, CALAN SR<sup>®</sup>, VERALAN<sup>®</sup>, VERELAN PM<sup>®</sup>)
  - felodipine (PLENDIL<sup>®</sup>)
  - nifedipine (ADALAT CC<sup>®</sup>, AFEDITAB CR<sup>®</sup>, PROCARDIA<sup>®</sup>, PROCARDIA XL<sup>®</sup>)
  - nicardipine (CARDENE<sup>®</sup>, CARDENE SR<sup>®</sup>)
- indigestion, heartburn, or ulcer medicines:
  - cimetidine (TAGAMET<sup>®</sup>)
  - esomeprazole (NEXIUM<sup>®</sup>)
  - famotidine (PEPCID<sup>®</sup>, PEPCID AC<sup>®</sup>)
  - rabeprazole (ACIPHEX<sup>®</sup>, ACIPHEX SPRINKLE<sup>™</sup>)
  - nizatidine (AXID<sup>®</sup>)
  - lansoprazole (PREVACID<sup>®</sup>)
  - omeprazole (PRILOSEC<sup>®</sup>)
  - pantoprazole (PROTONIX<sup>®</sup>)
  - ranitidine (ZANTAC<sup>®</sup>)
- antacids or buffered medicines
- antibiotics or antiviral medicines
  - clarithromycin (BIAXIN<sup>®</sup>, BIAXIN XL<sup>®</sup>)
  - didanosine (VIDEX<sup>®</sup>, VIDEX EC<sup>®</sup>)
  - efavirenz (SUSTIVA<sup>®</sup>)
  - rifabutin (MYCOBUTIN<sup>®</sup>)
  - ritonavir (NORVIR<sup>®</sup>)
  - saquinavir (INVIRASE<sup>®</sup>)
  - tenofovir disoproxil fumarate (VIREAD<sup>®</sup>)

Ask your healthcare provider or pharmacist if you are not sure if your medicine is one that is listed above.

**Know the medicines you take. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine.**

Your healthcare provider and your pharmacist can tell you if you can take these medicines with REYATAZ. Do not start any new medicines while you are taking REYATAZ without first talking with your healthcare provider or pharmacist. You can ask your healthcare provider or pharmacist for a list of medicines that can interact with REYATAZ.

### **How should I take REYATAZ?**

- **Take REYATAZ exactly as your healthcare provider tells you to.**
- **Do not change your dose or stop taking REYATAZ unless your healthcare provider tells you to.**
- Stay under the care of your healthcare provider during treatment with REYATAZ.
- REYATAZ must be used with other antiretroviral medicines.
- Take REYATAZ 1 time each day.
- REYATAZ comes as capsules and oral powder.
- **Take REYATAZ capsules and oral powder with food.**
- Swallow the capsules whole. **Do not open the capsules.**
- REYATAZ oral powder must be mixed with food or liquid. Your child's healthcare provider will prescribe the right dose of REYATAZ based on your child's weight.
- REYATAZ oral powder must be taken with ritonavir.
- If you miss a dose of REYATAZ, take it as soon as you remember. Then take the next dose at your regular time. Do not take 2 doses at the same time.
- **If you take too much REYATAZ**, call your healthcare provider or go to the nearest hospital emergency room right away.

**When your supply of REYATAZ starts to run low**, get more from your healthcare provider or pharmacy. It is important not to run out of REYATAZ. The amount of HIV-1 in your blood may increase if the medicine is stopped for even a short time. The virus may become resistant to REYATAZ and harder to treat.

### **What are the possible side effects of REYATAZ?**

**REYATAZ can cause serious side effects, including:**

- **A change in the way your heart beats (heart rhythm change).** Tell your healthcare provider right away if you get dizzy or lightheaded. These could be symptoms of a heart problem.
- **Skin rash.** Skin rash is common with REYATAZ but can sometimes be severe. Skin rash usually goes away within 2 weeks without any change in treatment. Severe rash may develop in association with other symptoms which could be serious. **If you develop a severe rash or a rash with any of the following**

**symptoms, stop taking REYATAZ and call your healthcare provider right away:**

- general feeling of discomfort or “flu-like” symptoms
- fever
- muscle or joint aches
- red or inflamed eyes, like “pink eye” (conjunctivitis)
- blisters
- mouth sores
- swelling of your face
- painful, warm, or red lump under your skin
- **Yellowing of your skin or the white part of your eyes** is common with REYATAZ but may be a symptom of a serious problem. These effects may be due to increases in bilirubin levels in your blood (bilirubin is made by the liver). Although these effects may not be damaging to your liver, skin, or eyes, tell your healthcare provider right away if your skin or the white part of your eyes turns yellow.
- **Liver problems.** If you have liver problems, including hepatitis B or C infection, your liver problems may get worse when you take REYATAZ. Your healthcare provider will do blood tests to check your liver before you start REYATAZ and during treatment. Tell your healthcare provider right away if you get any of the following symptoms:
  - your skin or the white part of your eyes turns yellow
  - dark “tea-colored” urine
  - light colored stools
  - nausea
  - itching
  - stomach-area pain
- **Kidney stones** have happened in some people who take REYATAZ. Tell your healthcare provider right away if you get symptoms of kidney stones which may include, pain in your low back or low stomach-area, blood in your urine, or pain when you urinate.
- **Gallbladder problems** have happened in some people who take REYATAZ. Tell your healthcare provider right away if you get symptoms of gallbladder problems which may include:
  - pain in the right or middle upper stomach area
  - fever
  - nausea and vomiting
  - your skin or the white part of your eyes turns yellow
- **Diabetes and high blood sugar (hyperglycemia)** have happened or have worsened in some people who take protease inhibitor medicines like REYATAZ.

Some people have had to start taking medicine to treat diabetes or have had to change their diabetes medicine.

- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Tell your healthcare provider if you start having new symptoms after starting your HIV medicine.
- **Changes in body fat** can happen in people taking HIV medicine. These changes may include increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the main part of your body (trunk). Loss of fat from the legs, arms, and face may also happen. The cause and long-term health effects of these conditions are not known.
- **Increased bleeding problems in people with hemophilia** have happened when taking protease inhibitors like REYATAZ.

The most common side effects of REYATAZ include:

- nausea
- headache
- stomach-area pain
- vomiting
- trouble sleeping
- numbness, tingling, or burning of hands or feet
- dizziness
- muscle pain
- diarrhea
- depression
- fever

Tell your healthcare provider if you have any side effect that bothers you or that does not go away.

These are not all the possible side effects of REYATAZ. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store REYATAZ?**

#### **REYATAZ capsules:**

- Store REYATAZ capsules at room temperature, between 68°F to 77°F (20°C to 25°C).
- Keep capsules in a tightly closed container.

**REYATAZ oral powder:**

- Store REYATAZ oral powder below 86°F (30°C).
- Store REYATAZ oral powder in the original packet. Do not open until ready to use.
- After REYATAZ oral powder is mixed with food or liquid it may be kept at room temperature 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

**Keep REYATAZ and all medicines out of the reach of children.**

**General information about REYATAZ**

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use REYATAZ for a condition for which it was not prescribed. Do not give REYATAZ to other people, even if they have the same symptoms that you have. It may harm them. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about REYATAZ that is written for health professionals.

For more information, go to [www.reyataz.com](http://www.reyataz.com) or call 1-800-321-1335.

**What are the ingredients in REYATAZ?**

**Active ingredient:** atazanavir sulfate

**Inactive ingredients:**

**REYATAZ capsules:** crospovidone, lactose monohydrate, and magnesium stearate. The capsule shells contain gelatin, FD&C Blue No. 2, titanium dioxide, black iron oxide, red iron oxide, and yellow iron oxide. The capsules are printed with ink containing shellac, titanium dioxide, FD&C Blue No. 2, isopropyl alcohol, ammonium hydroxide, propylene glycol, n-butyl alcohol, simethicone, and dehydrated alcohol.

**REYATAZ oral powder:** aspartame, sucrose, and orange-vanilla flavor.

This Patient Information has been approved by the U.S. Food and Drug Administration.

Distributed by:  
Bristol-Myers Squibb Company  
Princeton, NJ 08543 USA

Product of Ireland

[Print Code]

Revised: March 2015

VIDEX<sup>®</sup> and REYATAZ<sup>®</sup> are registered trademarks of Bristol-Myers Squibb Company. COUMADIN<sup>®</sup> and SUSTIVA<sup>®</sup> are registered trademarks of Bristol-Myers Squibb Pharma Company. Other brands listed are the trademarks of their respective owners and are not trademarks of Bristol-Myers Squibb Company.

**Instructions for Use  
REYATAZ® (RAY-ah-taz)  
(atazanavir)  
oral powder**

Read this Instructions for Use before you prepare your child's first dose of REYATAZ oral powder, each time you get a refill, and as needed. There may be new information. This information does not take the place of talking to your child's healthcare provider about their medical condition or treatment. Ask your child's healthcare provider or pharmacist if you have questions about how to mix or give a dose of REYATAZ oral powder.

**Important information:**

- For more information about REYATAZ oral powder, see the Patient Information leaflet.
- REYATAZ oral powder must be mixed with food or liquid. If REYATAZ oral powder is mixed with water, the child must eat food right after taking REYATAZ oral powder.
- REYATAZ oral powder must be taken with ritonavir.
- Talk with your child's healthcare provider to help decide the best schedule for giving your child REYATAZ oral powder.

**Instructions for mixing REYATAZ oral powder:**

REYATAZ oral powder should be mixed with food such as applesauce or yogurt, instead of a liquid (milk, infant formula, or water) in young children and infants who can take food.

- Infants less than 6 months old and who cannot eat solid food or drink from a cup should be given REYATAZ oral powder mixed with infant formula using an oral dosing syringe.
- REYATAZ oral powder that is mixed in infant formula or liquid should not be given using a baby bottle.

When preparing REYATAZ oral powder with either food or liquid, choose a clean, flat work surface. Place a clean paper towel on the work surface. Place the supplies you will need on the paper towel.


Wash and dry your hands before and after preparing REYATAZ oral powder.

### Preparing a dose of REYATAZ oral powder mixed with food:

Before you prepare a dose of REYATAZ oral powder mixed with food, gather the following supplies:


- paper towel
- tablespoon
- **small** clean container (such as a **small** cup or bowl)
- a food such as applesauce or yogurt
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose

**Step 1.** Place at least 1 tablespoon of a food such as applesauce or yogurt in the small container (see Figure A).



**Figure A**


**Step 2.** Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure B).



**Figure B**



**Step 3.** Using a clean pair of scissors, cut open the packet on the dotted line (see Figure C).



**Figure C**

**Step 4.** Empty the contents of the packet into the small container onto the food (see Figure D).



**Figure D**

**Repeat Steps 2 through 4 for each packet of REYATAZ oral powder needed for the total prescribed dose.**

**Step 5.** Use a tablespoon to gently mix the powder and the food together (see Figure E).



**Figure E**

Steps 6 through 8 must be completed **within 1 hour** of mixing the medicine.

**Step 6.** Use the tablespoon or a small spoon to feed the REYATAZ oral powder and food mixture to your child. Look in your child's mouth to make sure that all of the mixture is swallowed.

**Step 7.** Add 1 tablespoon more of food to the empty container and gently stir to mix with any contents that may still be in the container.

**Step 8.** Use the tablespoon or a small spoon to feed your child the mixture, making sure your child has swallowed all of the mixture.


**Step 9.** Give your child ritonavir as prescribed right after taking REYATAZ oral powder.

**Step 10.** Wash the container and tablespoon. Allow the container and spoon to dry. Throw away the paper towel and clean the work surface.

**Preparing a dose of REYATAZ oral powder mixed with liquid in a small drinking cup:**


Before you prepare a dose of REYATAZ oral powder mixed with liquid in a small drinking cup, gather the following supplies:

- paper towel
- spoon
- 30 milliliter (mL) medicine cup (ask your pharmacist for this). See Figure F.
- small drinking cup
- liquid such as milk or water
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose




**Figure F**

**Step 1.** Using the 30 mL medicine cup, pour at least 30 mL of liquid into the small drinking cup (see Figure G).




**Figure G**

**Step 2.** Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure H).



**Figure H**

**Step 3.** Using a clean pair of scissors, cut open the packet on the dotted line (see Figure I).



**Figure I**

**Step 4.** Empty the contents of the packet into the small drinking cup (see Figure J).



**Figure J**

**Repeat Steps 2 through 4 for each packet of REYATAZ oral powder needed for the total prescribed dose.**

**Step 5.** Hold the small drinking cup with one hand. With your other hand, use the spoon to gently mix the powder and the liquid (see Figure K).



**Figure K**

Steps 6 and 7 must be completed **within 1 hour** of mixing the medicine.

**Step 6.** Have your child drink all of the mixture in the small drinking cup.

**Step 7.** To make sure there is no mixture left in the small drinking cup add 15 mL more liquid to the small drinking cup:

- Stir with the spoon.
- **Repeat Step 6 above.**

**If REYATAZ oral powder is mixed with water, your child must eat food right after taking REYATAZ oral powder.**


**Step 8.** Give your child ritonavir as prescribed right after taking REYATAZ oral powder.

**Step 9.** Wash the small drinking cup, medicine cup, and spoon. Allow the small drinking cup, medicine cup, and spoon to dry. Throw away the paper towel and clean the work surface.

**Preparing a dose of REYATAZ oral powder mixed with liquid infant formula using an oral dosing syringe and a small medicine cup:**

Before you prepare a dose of REYATAZ oral powder mixed with infant formula using an oral dosing syringe, gather the following supplies:


- paper towel
- small spoon
- 30 milliliter (mL) medicine cup (ask your pharmacist for this). See Figure L.
- 10 mL oral dosing syringe (ask your pharmacist for this). See Figure L.
- infant formula
- the correct number of packets of REYATAZ oral powder needed for the prescribed dose



**Figure L**


**Step 1.** Prepare the infant formula according to the directions on the infant formula package.

**Step 2.** Pour 10 mL of infant formula into the medicine cup (see Figure M).




**Figure M**

**Step 3.** Tap the packet of REYATAZ oral powder to settle the contents to the bottom of the packet (see Figure N).



**Figure N**

**Step 4.** Using a clean pair of scissors, cut open the packet on the dotted line (see Figure O).



**Figure O**

**Step 5.** Empty the contents of the packet into the medicine cup (see Figure P).



**Figure P**

**Repeat Steps 3 through 5 for each packet of REYATAZ oral powder needed for the total prescribed dose.**

**Step 6.** Hold the medicine cup with one hand. With your other hand, use the small spoon to gently mix the powder and the infant formula (see Figure Q).



**Figure Q**

Steps 7 through 9 must be completed **within 1 hour** of mixing the medicine.

**Step 7.** Draw up the powder and infant formula mixture into the oral dosing syringe as follows:

- Check that the plunger is completely pushed into barrel of the syringe (see Figure R).



**Figure R**

- Place the tip of the syringe into the powder and infant formula mixture in the medicine cup (see Figure S).



**Figure S**

- Slowly pull back on the plunger and draw up 10 mL of the mixture (see Figure T).



Figure T

**Step 8.** Place the tip of the oral dosing syringe in your baby's mouth along the inner cheek on either the right or left side (see Figure U). Slowly push on the plunger to give your baby all of the REYATAZ oral powder and infant formula mixture.

- Draw up any remaining mixture with the oral dosing syringe and repeat until all of the mixture has been given to the baby.




Figure U

**Step 9.** To make sure there is no mixture left in the medicine cup or syringe:

- Repeat Step 1 above to add 10 mL more infant formula to the medicine cup
- Stir with a small spoon.
- **Then repeat Steps 7 through 8 above.**

**To make sure that your baby gets all of the medicine, do not give REYATAZ oral powder in a baby bottle.**

**Step 10.** Give your baby ritonavir as prescribed right after taking REYATAZ oral powder.

**Step 11.** Remove the plunger from the oral dosing syringe. Wash the medicine cup, spoon, and oral dosing syringe. Allow the medicine cup, spoon, and oral dosing syringe to dry. Throw away the paper towel and clean the work surface.



**How should I store REYATAZ oral powder?**

- Store REYATAZ oral powder below 86°F (30°C).
- Store REYATAZ oral powder in the original packet. Do not open until ready to use.
- After REYATAZ oral powder is mixed with food or liquid, it may be kept at room temperature 68°F to 86°F (20°C to 30°C) for up to 1 hour. Take REYATAZ oral powder within 1 hour after mixing with food or liquid.

**Keep REYATAZ oral powder and all medicines out of the reach of children.**

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Distributed by:  
Bristol-Myers Squibb Company  
Princeton, NJ 08543 USA

Product of Ireland

[Print Code]

Revised: March 2015