ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd. Bold Italic
Sb Heads -6 pt Helvetica Cnd. Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 3.5 pt Zaph Dingbats no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule
Hairline - .5 pt rule

Certified by: (b) (4)
APO: 2015-1137
04.27.15 - (b) (4)

8.4251 [214]

1.5550 [39.5]

3.0315 [77]

1.5748 [40]

9.7046 [246.5]

R0.2362 [R6]

0.5118 [13]

15

Claritin® RediTabs® 24 Hour 60 ORALLY DISINTEGRATING TABLETS FOR 60 DAYS OF RELIEF

Original Prescription Strength

Non-Drowsy*

Claritin®

RediTabs®

loratadine 10 mg/antihistamine

Indoor & Outdoor Allergies

24 Hour Relief of:
- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed Melts in Your Mouth

*When taken as directed. See Drug Facts Panel.

60 ORALLY DISINTEGRATING TABLETS

LOT/EXP AREA NO VARNISH

ITM # 918720

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer.
© 2015 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981
Product of Ireland.
XXXXX-XX

Bayer

recyclable carton

0 00000 00000 0

NO VARNISH

GLUE STRIP

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 10 mg	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

1. Peel back outer edge.
2. Gently push tablet out.
3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Follow these directions carefully. Do not attempt to push the tablet through the foil.

• No Water Needed • Melts in your mouth

Claritin® RediTabs® 24 Hour

Drug Facts (continued)

Other information
- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients
anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Reference ID: 3852067

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic
Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.
Headings - 8 pt Helvetica Cnd. Bold Italic
Sb Heads -6 pt Helvetica Cnd. Bold
Text - 6 pt Helvetica Cnd, minimum leading 6.5
Square Bullets - 3.5 pt Zaph Dingbats no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule
Hairline - .5 pt rule

Certified by: (b) (4)
APO: 2015-1985
04.28.15 - (b) (4)

8.4251 [214]

1.5550 [39.5]

3.0315 [77]

1.5748 [40]

9.7046 [246.5]

Claritin® RediTabs® 24 Hour 50 ORALLY DISINTEGRATING TABLETS FOR 50 DAYS OF RELIEF

Original Prescription Strength

Non-Drowsy*
Claritin®
RediTabs®
loratadine 10 mg/antihistamine
Indoor & Outdoor Allergies

24 Hour Relief of:
- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed Melts in Your Mouth

*When taken as directed. See Drug Facts Panel.

50 ORALLY DISINTEGRATING TABLETS

LOT/EXP AREA NO VARNISH

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer.
© 2015 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981
Product of Ireland.
XXXXX-XX

Bayer

recyclable carton

0 00000 00000 0

NO VARNISH

00000000

Claritin® RediTabs® 24 Hour

• No Water Needed • Melts in your mouth

Follow these directions carefully. Do not attempt to push the tablet through the foil.

1. Peel back outer edge.
2. Gently push tablet out.
3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Directions
- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 10 mg	Antihistamine

Drug Facts (continued)

Other information
- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients
anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

GLUE STRIP

R0.2362 [R6]

0.5118 [13]

15

Reference ID: 3852067

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads -6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: (b) (4)
APO #: 2015-1135
04.27.15 (b) (4)

7.7571 [197.03]
1.320 [33.5]
4.764 [121]
3.032 [77.00]
9.547 [242.50]
1.496 [38.00]

Claritin® RediTabs® 24 Hour 30 ORALLY DISINTEGRATING TABLETS FOR 30 DAYS OF RELIEF

Original Prescription Strength

Non-Drowsy*
Claritin®
RediTabs®
loratadine 10 mg/antihistamine
Indoor & Outdoor Allergies

24 Hour
Relief of:
- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed Melts in Your Mouth

*When taken as directed. See Drug Facts Panel.

30 ORALLY DISINTEGRATING TABLETS

LOT/EXP AREA NO VARNISH

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer.
© 2015 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981
Product of Ireland.
XXXXX-XX

Bayer

recyclable carton

0 00000 00000 0

NO VARNISH

00000000

GLUE STRIP

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 10 mg	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

1. Peel back outer edge.
2. Gently push tablet out.
3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Follow these directions carefully. Do not attempt to push the tablet through the foil.

• No Water Needed • Melts in your mouth

Claritin® RediTabs® 24 Hour

Questions or comments?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Inactive ingredients
anhydrous citric acid, gelatin, mannitol, mint flavor

■ complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Other information
- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Drug Facts (continued)

Reference ID: 3852067

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads -6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: (b) (4)
APO #: 2015-1134
04.26.15 (b) (4)

165
121
21.5
77
210.5
22
R2.25
R6
13

Claritin® RediTabs®

24 Hour 10 ORALLY DISINTEGRATING TABLETS FOR 10 DAYS OF RELIEF

Original Prescription Strength

Non-Drowsy*

Claritin®

RediTabs®

loratadine 10 mg/antihistamine

Indoor & Outdoor Allergies

24 Hour Relief of:
- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itchy Throat or Nose

No Water Needed Melts in Your Mouth

BAYER *When taken as directed. See Drug Facts Panel.

10 ORALLY DISINTEGRATING TABLETS

Claritin® RediTabs®

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer.
© 2015 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981
Product of Ireland.
XXXXX-XX

Bayer

recyclable carton

0 00000 00000 0

NO VARNISH

00000000

GLUE STRIP

Drug Facts

Active ingredient (in each tablet) — **Purpose**
Loratadine 10 mg ..... Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

1. Peel back outer edge.
2. Gently push tablet out.
3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Follow these directions carefully. Do not attempt to push the tablet through the foil.

• No Water Needed • Melts in your mouth

Claritin® RediTabs® 24 HOUR

Questions or comments? 1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Inactive ingredients
anhydrous citric acid, gelatin, mannitol, mint flavor

Drug Facts (continued)

Other information
- Safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Drug Facts (continued)
Drug Facts - 8 pt Helvetica Cnd. Bold Italic (continued) - 8 pt Helvetica Cnd.

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads -6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 3.5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

(b) (4)

Certified by:
APO #: 2015-1171
04.27.15 (b) (4)

Reference ID: 3852067



APO #: 2015-1999
04.29.15 - (b) (4)

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads -6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: (b) (4)

APO #: 2015-1091
04.28.15 - (b) (4)

7.7571 [197.03]

1.320 [33.5]

4.764 [121]

9.547 [242.50]

3.032 [77.001]

1.496 [38.001]

Claritin® RediTabs® 12 Hour 30 ORALLY DISINTEGRATING TABLETS

Non-Drowsy*
Claritin®
RediTabs®
loratadine 5 mg/antihistamine
Indoor & Outdoor Allergies
12 Hour
Relief of:
• Sneezing
• Runny Nose
• Itchy, Watery Eyes
• Itchy Throat or Nose
No Water Needed Melts in Your Mouth
* When taken as directed. See Drug Facts Panel.
30 ORALLY DISINTEGRATING TABLETS

LOT/EXP AREA NO VARNISH

Bayer, the Bayer Cross, Claritin, the Claritin Vista and RediTabs are registered trademarks of Bayer.
© 2015 Bayer.
Dist by: Bayer HealthCare LLC
Whippany, NJ 07981
Product of Ireland.
XXXXX-XX

Bayer

0 00000 00000 0

NO VARNISH

00000000

GLUE STRIP

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 5 mg	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings
Do not use if you have ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.
When using this product do not take more than directed. Taking more than directed may cause drowsiness.
Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

1. Peel back outer edge.
2. Gently push tablet out.
3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

Follow these directions carefully. Do not attempt to push the tablet through the foil.

• No Water Needed • Melts in your mouth

Claritin® RediTabs® 12 Hour

Questions or comments?
1-800-CLARITIN (1-800-252-7484) or www.claritin.com

Inactive ingredients
anhydrous citric acid, gelatin, mannitol, mint flavor

Other information
- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Drug Facts (continued)

Reference ID: 3852067

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic
Headings - 8 pt Helvetica Cnd. Bold Italic
Sub Heads -6 pt Helvetica Cnd. Bold
Text - 6 pt Helvetica Cnd. minimum leading 6.5
Square Bullets - 5 pt Zaph Dingbats no compression
Border - 1 pt rule, 3/64" gap all around
Barline - 1 pt rule
Hairline - .5 pt rule

Certified by: (b) (4)
APO #: 2015-1090 (b)
04.29.15 - (4)

ATTACH LEGEND ON ALL DIGITAL MECHANICALS

STANDARD FORMAT
Helvetica Condensed
Variable horizontal scale adheres to 39 characters per inch

Drug Facts - 9 pt Helvetica Cnd. Bold Italic

Headings - 8 pt Helvetica Cnd. Bold Italic

Sub Heads -6 pt Helvetica Cnd. Bold

Text - 6 pt Helvetica Cnd, minimum leading 6.5

Square Bullets - 5 pt Zaph Dingbats no compression

Border - 1 pt rule, 3/64" gap all around

Barline - 1 pt rule

Hairline - .5 pt rule

Certified by: (b) (4)

APO #: 2015-1170
04.28.15 - (b) (4)

Reference ID: 3852067

APO #: 2015-1996
04.29.16 · (b) (4)

Reference ID: 3852067