

Sterile Water for Injection, USP

10 mL Single Dose Vial

Container Label: 462-683-00

Submission 5

7 / 23 / 2015

(b) (4)

WARNINGS: NOT ISOTONIC. HEMOLYTIC.
NDC 0641-6147-01  only
Contains no antimicrobial or other added substance. Sterile, nonpyrogenic. **Do not give intravenously unless rendered nearly isotonic.**

Sterile Water for Injection, USP

FOR DRUG DILUENT USE ONLY

10 mL Single Dose Vial

USUAL DOSAGE: See insert. Store at 20° to 25°C (68° to 77°F) [See USP]. DO NOT FREEZE.
Mid. by:  WESTWARD
Eatontown, NJ 07724 USA
462-683-00

LOT:  (01)0030641 6147019
EXP: 

Sterile Water for Injection, USP

25 x 10 mL Single Dose Vials

Shelf Pack: 462-684-00

Submission 5

7 / 23 / 2015

(b) (4)

LOT: _____
EXP: _____

NDC 0641-6147-25 **Rx only**

**Sterile Water
for Injection, USP**

FOR DRUG DILUENT USE ONLY

25 x 10 mL Single Dose Vials

Mfd. by:  WEST-WARD
Eatontown, NJ 07724 USA 462-684-00

Contains no antimicrobial or other added substance.
Sterile, nonpyrogenic. Use only if clear and seal is intact and undamaged.
Do not give intravenously unless rendered nearly isotonic. Use promptly; discard unused portion.
USE ASEPTIC TECHNIQUE. Remove cover from vial and cleanse stopper with antiseptic.

With Sterile Syringe and Needle:

1. Aspirate desired portion of vial contents and add to suitable container.
2. Discard any remaining fluid in vial.

Store at 20° to 25°C (68° to 77°F)
[See USP Controlled Room Temperature].
DO NOT FREEZE.

(01)0030041614725





068200

462-682-00

Sterile Water for Injection, USP

Rx only

462-682-00

Sterile Water
for Injection, USPSubmission 1
7-23-2015Drawing: 6INS577B (1)
Size: 4 1/2" x 16 1/2"

All Print Black

DESCRIPTION

This preparation is designed solely for parenteral use only after addition of drugs that require dilution or must be dissolved in an aqueous vehicle prior to injection.

Sterile Water for Injection, USP is a sterile, nonpyrogenic preparation of water for injection which contains no bacteriostat, antimicrobial agent or added buffer and is supplied only in single-dose containers to dilute or dissolve drugs for injection. For intravenous injection, add sufficient solute to make an approximately isotonic solution.

Water for Injection, USP is chemically designated H₂O.

The glass vial is a Type I borosilicate glass and meets the requirements according to the USP standards.

CLINICAL PHARMACOLOGY

Water is an essential constituent of all body tissues and accounts for approximately 70% of total body weight. Average normal adult daily requirement ranges from two to three liters (1.0 to 1.5 liters each for insensible water loss by perspiration and urine production).

Water balance is maintained by various regulatory mechanisms. Water for distribution depends primarily on the concentration of electrolytes in the body compartments and sodium (Na⁺) plays a major role in maintaining physiologic equilibrium.

The small volume of fluid provided by Sterile Water for Injection, USP when used only as a pharmaceutical aid for diluting or dissolving drugs for parenteral injection, is unlikely to exert a significant effect on fluid balance except possibly in neonates or very small infants.

INDICATIONS AND USAGE

This parenteral preparation is indicated only for diluting or dissolving drugs for intravenous, intramuscular or subcutaneous injection, according to instructions of the manufacturer of the drug to be administered.

CONTRAINDICATIONS

Sterile Water for Injection, USP must be made approximately isotonic prior to use.

WARNINGS

Intravenous administration of Sterile Water for Injection without a solute may result in hemolysis.

PRECAUTIONS

Do not use for intravenous injection unless the osmolar concentration of additives results in an approximate isotonic admixture.

Consult the manufacturer's instructions for choice of vehicle, appropriate dilution or volume for dissolving the drugs to be injected, including the route and rate of injection.

Inspect reconstituted (diluted or dissolved) drugs for clarity (if soluble) and freedom from unexpected precipitation or discoloration prior to administration.

Pregnancy Category C

Animal reproduction studies have not been conducted with Sterile Water for Injection. It is also not known whether sterile water containing additives can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sterile Water for Injection with additives should be given to a pregnant woman only if clearly needed.

Pediatric Use

Safety and effectiveness have been established in pediatric patients. However, in neonates or very small infants the volume of fluid may affect fluid and electrolyte balance.

Drug Interactions

Some drugs for injection may be incompatible in a given vehicle, or when combined in the same vehicle or in a vehicle containing benzyl alcohol. Consult with pharmacist, if available.

Use aseptic technique for single or multiple entry and withdrawal from all containers.

When diluting or dissolving drugs, mix thoroughly and use promptly.

Do not store reconstituted solutions of drugs for injection unless otherwise directed by the manufacturer of the solute.

Do not use unless the solution is clear and seal intact. Do not reuse single-dose containers. Discard unused portion.

ADVERSE REACTIONS

Reactions which may occur because of this solution, added drugs or the technique of reconstitution or administration include febrile response, local tenderness, abscess, tissue necrosis or infection at the site of injection, venous thrombosis or phlebitis extending from the site of injection and extravasation.

If an adverse reaction does occur, discontinue the infusion, evaluate the patient, institute appropriate countermeasures, and if possible, retrieve and save the remainder of the unused vehicle for examination.

OVERDOSAGE

Use only as a diluent or solvent. This parenteral preparation is unlikely to pose a threat of fluid overload except possibly in neonates or very small infants. In the event these should occur, re-evaluate the patient and institute appropriate corrective measures. See **WARNINGS**, **PRECAUTIONS** and **ADVERSE REACTIONS**.

DOSAGE AND ADMINISTRATION

The volume of the preparation to be used for diluting or dissolving any drug for injection is dependent on the vehicle concentration, dose and route of administration as recommended by the manufacturer.

This parenteral should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED

Sterile Water for Injection, USP is supplied in the following:

10 mL single dose vials packaged in carton of 25 vials (NDC 0641-6147-25)

Storage

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature.] DO NOT FREEZE.

To report SUSPECTED ADVERSE REACTIONS, contact West-Ward Pharmaceuticals Corp. at 1-877-845-0689, or the FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

For Product Inquiry call 1-877-845-0689.

Manufactured by



WEST-WARD
PHARMACEUTICALS

Eatontown, NJ 07724 USA

Appear this way
on original

