New Information: Read Entire Label
NDC 50580-110-02
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
50 Coated Caplets*
actual size
*Capsule-Shaped Tablets
Drug Facts (continued)
Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, sha lac, stearic acid, titanium dioxide
Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8255 (collect)
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© J&J CI 2016
www.motrin.com
Drug Facts (continued)
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
■ do not take more than directed
■ the smallest effective dose should be used
adults and children 12 years and older
■ take 1 caplet every 4 to 6 hours while symptoms persist
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years
■ ask a doctor
Other information
■ store between 20-25°C (68-77°F)
■ do not use if neck wrap or foil inner seal is imprinted "SAFETY SEAL@" is broken or missing
Drug Facts (continued)
Important: Read all product information before using. Keep this box for important information.
Active ingredient (in each caplet) Purpose
ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug
Uses
■ temporarily relieves minor aches and pains due to
■ headache
■ minor pain of arthritis
■ backache
■ menstrual cramps
■ temporarily reduces fever
■ muscular aches
■ toothache
■ the common cold
Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin
Symptoms may include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have stomach pain that does not get better
■ feel faint ■ vomit blood
■ have bloody or black stools
Skin use and ask a doctor if
■ you experience any of the following signs of stomach bleeding
Stop use and ask a doctor if
■ take with food or milk if stomach upset occurs
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
When using this product
■ under a doctor's care for any serious condition
■ taking any other drug
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you take asthma medicine ■ you are taking a diuretic
Do not use
■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery
■ take more or for a longer time than directed
■ take 3 or more alcoholic drinks every day while using this product
LOT
EXP
3 0045-0481-02 3
00000000
00000000
00000000

(b) (4)

(b) (4)

New Information: Read Entire Label
NDC 50580 110 01
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
100 Coated Caplets*
*Capsule-Shaped Tablets
Drug Facts (continued)
Drug Facts
Drug Facts (continued)
Drug Facts (continued)
Directions
Other information
Inactive ingredients
LOT:
EXP:
3 0045-0481-016
00000000

(b) (4)

New Information: Read Entire Label
NDC 50580 110 10
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
100 Coated Caplets*
*Capsule-Shaped Tablets
Drug Facts (continued)
Drug Facts
Drug Facts (continued)
Drug Facts (continued)
Direct ons
Other information
Inactive ingredients
LOT:
EXP:
3 0045-0481-016
00000000
0 00 00 000
00000000

(b) (4)

Motrin IB
Ibuprofen Tablets, USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
100 Coated Caplets*
*Liquid-Filled Tablets

(b) (4)


LOT:
EXP.
Important: Read all product information before using. Keep the box for important information.
Active ingredient (in each caplet) Purpose
Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug
Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches ■ minor pain of arthritis ■ toothache
■ backache ■ the common cold ■ menstrual cramps
■ temporarily reduces fever
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug. When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain ■ that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (800 222-1222)
Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor
Other information ■ store between 20°-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL" is broken or missing. Inactive ingredients carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments? Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)
Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.motrin.com
NDC 50580-710-10
Motrin® IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
100 Coated Caplets* *Capsule-Shaped Tablets




BASE LABEL,
PAGE 5
GLUE APPLIED
HERE




PAGE 2

PAGE 3

PAGE 4
(GLUED TO BASE)

COVER, PAGE 1

(b) (4)

Drug Facts
Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug
Purpose
Pain reliever/fever reducer
Uses
■ temporarily relieves minor aches and pains due to:
■ muscular aches
■ minor pain of arthritis
■ backache
■ toothache
■ the common cold
■ menstrual cramps
■ temporarily reduces fever
Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives
■ facial swelling
■ asthma (wheezing)
■ shock
■ skin reddening
■ rash
■ dizziness
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Do not use
■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
New Information: Read Entire Label
NDC 50580-110-37
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300 Coated Caplets*
*Capsule-Shaped Tablets
New Information: Read Entire Label
NDC 50580-110-37
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300 Coated Caplets*
*Capsule-Shaped Tablets
Drug Facts (continued)
■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver disease, or kidney disease
■ you have asthma
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug
When using this product
■ take with food or milk if stomach upset occurs
■ the smallest effective dose should be used
■ if you use more than directed or for longer than directed
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during birth.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
■ do not take more than directed
■ the smallest effective dose should be used
adults and children 12 years and older
■ take 1 caplet every 4 to 6 hours while symptoms last
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years
■ ask a doctor
Drug Facts (continued)
Other information
■ store between 20-25°C (68-77°F)
■ do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL®" is broken or missing
Inactive ingredients
carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments?
Call 1-877-895-3665 (toll free) or 215-273-8755 (collect)
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2018
www.motrin.com
EXP:
LOT:
3 0045-0781-37 5

(b) (4)

Drug Facts
Purpose
Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)* ..... Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug
Uses
■ temporarily relieves minor aches and pains due to:
■ headache ■ muscular aches
■ minor pain of arthritis ■ toothache
■ backache ■ the common cold
■ menstrual cramps
■ temporarily reduces fever
Warnings
Allergy alert: ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling
■ asthma (wheezing) ■ shock
■ shortness of breath ■ rash
■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Do not use
■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
New Information: Read Entire Label
NDC 50580-110-38
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300 Coated Caplets*
*Capsule-Shaped Tablets
New Information: Read Entire Label
NDC 50580-110-38
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300 Coated Caplets*
*Capsule-Shaped Tablets
Drug Facts (continued)
■ the stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver disease, or kidney disease
■ you have asthma
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug
When using this product
■ take with food or milk if stomach upset occurs
■ the more you smoke while using this product, the greater your risk for getting stomach problems
■ if you use more than directed or for longer than directed
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding
■ feel faint ■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during birth.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
■ do not take more than directed
■ the smallest effective dose should be used
adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms last ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
children under 12 years: ■ ask a doctor
Drug Facts (continued)
Other information
■ store between 20-25°C (68-77°F)
■ do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL" is broken or missing
Inactive ingredients
carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments?
Call 1-877-895-3665 (toll free) or 215-273-8755 (collect)
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2018
www.motrin.com
00000000
EXP:
LOT:
3 0045-0781-37 5

(b) (4)

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug
Purpose
Pain reliever/fever reducer
Uses ■ temporarily relieves minor aches and pains due to
■ headache ■ muscular aches ■ minor pain of arthritis
■ toothache ■ backache ■ the common cold
■ menstrual cramps
■ temporarily reduces fever
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor
Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL" is broken or missing
Inactive ingredients: carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments? Call 1-877-895-3665 (toll free) or 215-273-8755 (collect)
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.motrin.com
NDC 50580-110-37
Motrin® IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300
Coated Caplets*
*capsule-shaped tablets

(b) (4)

Active Ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug
Purpose
Pain reliever/fever reducer
Uses ■ temporarily relieves minor aches and pains due to
■ headache ■ muscular aches ■ minor pain of arthritis
■ toothache ■ backache ■ the common cold
■ menstrual cramps
■ temporarily reduces fever
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed. Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery. Ask a doctor before use if you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma ■ you are taking a diuretic. Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug
When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed. Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear. If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions ■ do not take more than directed ■ the smallest effective dose should be used. Adults and children 12 years and older: ■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor. Children under 12 years: ■ ask a doctor
Other information ■ store between 20-25°C (68-77°F) ■ do not use if neck wrap or foil inner seal imprinted "SAFETY SEAL" is broken or missing
Inactive ingredients: carnauba wax, colloidal silicon dioxide, FD&C yellow no. 6, hypromellose, iron oxide, magnesium stearate, modified starch, polydextrose, polyethylene glycol, pregelatinized starch, propylene glycol, shellac, stearic acid, titanium dioxide
Questions or comments? Call 1-877-895-3665 (toll free) or 215-273-8755 (collect)
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
©J&JCI 2016
www.motrin.com
NDC 50580-110-38
Motrin® IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
300
Coated Caplets*
*capsule-shaped tablets

(b) (4)

00000000
New Information: Read Entire Label
For Hospital and Government Use Only
NDC 50580 110 60
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever/Fever Reducer (NSAID)
60 Coated Caplets
Transfer Shape Tablets
EXP
LOT
50580 11060 7
00000000

(b) (4)

Motrin IB
Rob Bellows
Fever Reducer (NSA 2)
60 cm
Copyright
Reproduction
Keep the Insert Important Information

(b) (4)




(b) (4)

Drug Facts
Active ingredient (in each caplet)
Purpose
Pain reliever/ fever reducer
Use
Warnings
Drug Facts (continued)
Directions
Other information
Inactive ingredients
Questions or comments?
New Information: Read Entire Label
NDC 50580 116 52
Motrin IB
Ibuprofen Tablets USP, 200 mg
Pain Reliever / Fever Reducer (NSAID)
Convenient Take-Along Packs
Coated Caplets
50 Pouches of 2 Coated Caplets* Each
*Capsule Shaped Tablets

(b) (4)

---

**This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.**

---

/s/

---

VALERIE S PRATT
08/18/2016