





PAGE 2

PAGE 7



PAGE 3




PAGE 4



PDP = 2.41 square inches Anti-Diarrheal





PAGE 6

Reference ID: 3912606








COVER, PAGE 1

PAGE 5






Base Label

BOOKLET PAGES:



Cover Page 1



Page 2

Page 7



PDP = 2.41 square inches



Fraction - Decimal Conversion Chart		
	Inches	
16	.0625	
18	.125	
3 16	.1875	







Drug Facts (continued)

Other information

- each 30 mL contains: sodium 16 mg
 store between 20-25°C (68-77°F)
 do not use if printed inner or outer n is broken or missing

nactive ingredients hydrous citric acid, caramel color, D&C yellow no. 10, b&C blue no. 1, flavor, glycerin, mic ocrystalline cellulos d ca boxymethyclellulose sodium, propylene glycol, rified water, simethicone emulsion, sodium benzoate, itanium dioxide, xanthan gum

Page 6

Drug Facts (continued)

/hen using this product tiredness, d owsiness or izziness may occur. Be careful when driving or perating machinery.

Stop use and ask a doctor if


- Stop use and ask a vocuor in symptoms get worse diarrhea lasts for more than 2 days you get abdominal swelling or bulging. These may be signs of a serious condition
- f pregnant or breast-feeding, ask a hea th
- onal before use. Keep out of reach of children. In case of overdo
- t medica help or contact a Poison Control Cente ht away. (1-800-222-1222)


Page 3

Drug Facts (continued) Directions 4 drink plenty of clear fluids to help prevent dehydration caused by diarrhea Find ight dose on chart. If possible, use weight to dose; otherwise use age. shake well before using use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device. mL = milliliter		Drug Facts (continued)	
		children 9-11 years (60-95 lbs) to 5 mL after first loose stool; 7 5 mL after each subsequent loose stool; but no more than 45 mL in 24 hours children 6-8 years (48-59 lbs) (48-59 lbs) to mL after first loose stool; 7 5 mL after each subsequent loose stool; but no more than 30 mL in 24 hours	
			7 5 mL after each subsequent loose stool; but no more than
	30 mL after the first loose stool; 15 mL after each subsequent loose stool; but no more than 60 mL in 24 hours	children under 6 years (up to 47 lbs)	ask a doctor

Page 4


Page 5





PAGE 2

PAGE 7




PAGE 3

Drug Facts (continued) Directions If of class of class fluids to help prevent dehydration caused by diarrhea lind right dose on chart. If possible, use weight to dose; othe wise use age. shake well before using use only enclosed dosing cup specifically designed fo with this p oduct. Do not use any other dosing device. ml = milliliter adults and children 30 mL after the first loose stool; 12 years and over 15 mL after each subsequent 60 mL in 24 hours

PAGE 4







PAGE 6


Reference ID: 3912606








COVER, PAGE 1



PAGE 5





Base Label

BOOKLET PAGES:



Cover Page 1



Page 2

Inches

-.0625

.125

-1875

16

 $\begin{pmatrix} 1\\ 8 \end{pmatrix}$

3

Page 7









Drug Facts (continued)

Other information

- each 30 mL contains: sodium 16 mg
 store between 20-25°C (68-77°F) do not use if printed inner or outer is broken or missing

active ingredients hydrous citric acid, caramel color, D&C yellow no. 10, &C blue no. 1, flavor, glycerin, mic ocrystalline cellulo ca boxymethylcellulose sodium, propylene glycol, ed water, simethicone emulsion, sodium benzoate itanium dioxide, xanthan gum

Page 6

Drug Facts (continued)

hen using this product tiredness, d owsiness or zziness may occur. Be careful when driving or erating machinery.


Stop use and ask a doctor if

symptoms get worse
diarrhea lasts for more than 2 days

- you get abdominal swelling or bulging.
 These may be signs of a serious conditi
- f pregnant or breast-feeding, ask a hea th
- onal before use. Keep out of reach of children. In case of overdo


et medica help or contact a Poison Control Center ght away. (1-800-222-1222)

Page 3



Page 4

Page 5



Anti-Diarrheal

Controls the symptoms of diarrhea


Mint Flay

COVER,

PAGE 1


Oral Solution

8 fl oz (240 mL)



Drug Facts	Drug Facts (continued)	
Active ingredient (in each 7.5 mL) Purpose Loperamide HCl 1 mgAnti-diarrhea	Stop use and ask a doctor if symptoms get wors diarrhea lasts for more than 2 days you get abdominal swelling or bulging. These may be	
Use controls symptoms of diarrhea, including Travelers' Diarrhea	signs of a serious condition.	
	If pregnant or breast-feeding, ask a health professional before use.	
Warnings Allergy alert Do not use if you have ever had a rash or other allergic reaction to loperamide HCI	Keep out of reach of children. In case of ove dose, get medical help or c_ntact a Pois_n Control Center right away. (1 800 222-1222	
Do not use if you have bloody or black stool	Directions	
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease	drink plenty of clear fluids to help prevent dehydration caused by diarrhea find right dose on chart. If poss ble, use weight to dose; otherwise use age. shake well before using	
Ask a doctor or pharmacist before use if you are taking ant biotics		
When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.	only use enclosed dosing up specifically designed for use with his product. Do not use any o her dosing device.	





use if pri and is b




PAGE 4

LABEL


GLUED TO BASE


	Anti-Diarrheal Oral Solution	Fraction - Decimal Conversion Chart
.073"	Controls the symptoms of diarrhea 8 floz	16 .0625
	(240 mL)NN Mint Flavor	31875


BASE LABEL, PAGE 5 GLUE APPLIED HERE



PAGE 3







PAGE 4


COVER, PAGE 1

Drug Facts	Drug Facts (continued)	
Active ingredient (in each 7.5 mL) Purpose Loperamide HCl 1 mgAnti-diarrheal		
Use controls symptoms of diarrhea, including Travelers'		
Diarrhea	If pregnant or breast-feeding, ask a health professional	
Warnings Allergy alert Do not use if you have ever had a rash or other a lergic reaction to loperamide HCI	before use. Keep out of reach of children . In case of ove dose, get medical help or contact a Poison Control Center right away. (800 222-1222)	
Do not use if you have bloody or black stool	Directions	
Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of iver disease	 drink plenty of clear fluids to help prevent dehydration caused by diarrhea find right dose on chart. If possible, use weight to dose; otherwise use age. shake well before using only use enclosed dosing cup specifically designed for use with his conduct. Do not use any o her dosing device. 	
Ask a doctor or pharmacist before use if you are taking antibiotics		
When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.		

PAGE 2



1 Page(s) of Draft Labeling have been Withheld in Full as B4 (CCI/TS) immediately following this page



PAGE 3

(b) (4