Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	70%
Headers:	8.00 pt	Leading:	6.50 pt	Barlines:	2.50 pt		
Subheaders:	6.00 pt	Drug Facts (continued):	8.00 pt	Hairlines:	0.50 pt	Average Tracking:	-25

McNeil Submission Artwork

(b) (4)

Labeling Format Information:	Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 75%	
Headers: 4.50 pt	Leading: 4.50 pt	Barlines: NA pt		
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: -25	

McNeil Submission Artwork

(b) (4)

NDC 50580-783-10

TO OPEN: 1. PUSH DOWN 2. TURN CAP

See New Warning

TYLENOL® 8HR

ARTHRITIS PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain

*Capsule-Shaped Bi-Layer Tablets

Actual Size

100 Caplets*
650 mg each

OPEN TO READ DIRECTIONS AND ADDITIONAL DRUG FACTS INFORMATION

Do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing.

3 0045-0838-21 6

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

Contains No Aspirin

Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg.....Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

COVER, PAGE 1

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 6 caplets in 24 hours, which is the maximum daily amount
- w th other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medica help right away.

Do not use
- w th any other drug containing acetaminophen (prescription or nonpresc iption). If you are not sure whether a drug contains acetaminophen, ask a doctor or pha macist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfa in

Stop use and ask a doctor if
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur ■ redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a hea th professiona before use.
Keep out of reach of children.
Overdose warning: In case of ove dose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

BACK OF COVER, PAGE 2

Drug Facts (continued)

Directions
- **do not take more than directed (see overdose warning)**

adults	■ take 2 caplets every 8 hours w th water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
under 18 years of age	■ ask a doctor

Other information
- store between 20-25°C (68-77°F)
- **do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Drug Facts (continued)

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

TYLENOL® 8 HR ARTHRITIS PAIN
Acetaminophen extended-release tablets
100 Caplets – 650 mg each

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

BASE LABEL, PAGE 3

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	10.00 pt	Body Text:	6.5-7.0 pt	Bullets:	5.0-6.0 pt	Horizontal Scale:	85%
Headers:	8.0-9.0 pt	Leading:	6.5-7.5 pt	Barlines:	2.50 pt		
Subheaders:	7.0 pt	Drug Facts (cotinued):	9.00 pt	Hairlines:	0.50 pt	Average Kerning:	-25

McNeil Submission Artwork

(b) (4)

Reference ID: 3965345

Drug Facts

Active ingredient (in each caplet) — **Purpose**

Acetaminophen 650 mg ..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults	■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
under 18 years of age	■ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if carton is opened, or if foil inner seal imprinted with "SAFETY SEAL®" is broken**

Inactive ingredients

carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

00000000

3 0045-0838-377

EXP.

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

Contains No Aspirin

Care To Recycle™

TYLENOL® 8HR ARTHRITIS PAIN

HERITAGE OF CARE • TYLENOL • HERITAGE OF CARE

How can we help?

1-877-895-3665

00000000

NDC 50580-783-25

See New Warning

TYLENOL® 8HR ARTHRITIS PAIN

Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer
For The Temporary Relief Of Minor Arthritis Pain

*Capsule-Shaped Bi-Layer Tablets

Actual Size

225 Caplets* 650 mg each

00000000

Labeling Format Information: Fonts: HelveticaDF Medium, Bold, Bold Italic

Drug Facts:	10.00 pt	Body Text:	7.00 pt	Bullets:	5.00 pt	Horizontal Scale:	85%
Headers:	9.00 pt	Leading:	7.50 pt	Barlines:	2.50 pt		
Subheaders:	7.00 pt	Drug Facts (cotinued):	9.00 pt	Hairlines:	0.50 pt	Average Kerning:	-25

McNeil

(b) (4)

Labeling Format Information:	Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 6.00 pt	Bullets: 5.00 pt	Horizontal Scale: 90%	
Headers: 8.00 pt	Leading: 6.70 pt	Barlines: NA pt		
Subheaders: 6.00 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: -25	

McNeil Submission Artwork

(b) (4)

00000000

Drug Facts

Active ingredient (in each caplet) **Purpose**

Acetaminophen 650 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

HERITAGE OF CARE • HERITAGE OF CARE

TYLENOL

How can we help?

1-877-895-3665

00000000

NDC 50580-783-29

EXP.

See New Warning

TYLENOL® 8HR ARTHRITIS PAIN

Acetaminophen Extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain

*Capsule-Shaped Bi-Layer Tablets

Actual Size

290 Caplets* 650 mg each

TYLENOL® 8HR ARTHRITIS PAIN

Acetaminophen Extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain

Contains No Aspirin

3 0045-0838-29 2

00000000

Drug Facts (continued)

Directions

- do not take more than directed (see overdose warning)

adults	■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
under 18 years of age	■ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened, or if foil inner seal imprinted with "SAFETY SEAL@" is broken

Inactive ingredients

carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Care To Recycle™

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

00000000

Labeling Format Information: Fonts: HelveticaDF Medium, Bold, Bold Italic

Drug Facts:	10.00 pt	Body Text:	7.00 pt	Bullets:	5.00 pt	Horizontal Scale:	85%
Headers:	9.00 pt	Leading:	7.50 pt	Barlines:	2.50 pt		
Subheaders:	7.00 pt	Drug Facts (cotinued):	9.00 pt	Hairlines:	0.50 pt	Average Kerning:	0

McNeil Submission Artwork

(b) (4)

Active ingredient (in each caplet) **Purpose**

Acetaminophen 650 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - muscular aches
  - backache
  - premenstrual and menstrual cramps
  - the common cold
  - headache
  - toothache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medica help right away.

Do not use ■w th any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

EXP.

LOT:

TO OPEN: PUSH AND TURN CAP

READ THE LABEL

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

NDC 50580-783-29

TYLENOL® 8HR

ARTHRITIS PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For The Temporary Relief Of Minor Arthritis Pain

*Capsule-Shaped Bi-Layer Tablets

290 Caplets*
650 mg each

- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a hea th professional before use.

Keep out of reach of children.

Overdose warning: In case of ove dose, get medica help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

Adults:

- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

Under 18 years of age: ■ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- **do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients

ca nauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	90%
Headers:	8.00 pt	Leading:	7.80 pt	Barlines:	NA pt		
Subheaders:	6.00 pt	Drug Facts (cotinued):	NA pt	Hairlines:	NA pt	Average Kerning:	-25

McNeil Submission Artwork

(b) (4)

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	70%
Headers:	8.00 pt	Leading:	6.50 pt	Barlines:	2.50 pt		
Subheaders:	6.00 pt	Drug Facts (continued):	8.00 pt	Hairlines:	0.50 pt	Average Kerning:	-25

McNeil Submission Artwork

(b) (4)



Labeling Format Information:	Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic			
Drug Facts: NA pt	Body Text: 4.50 pt	Bullets: 3.50 pt	Horizontal Scale: 80%	
Headers: 4.50 pt	Leading: 4.70 pt	Barlines: NA pt		
Subheaders: 4.50 pt	Drug Facts (cotinued): NA pt	Hairlines: NA pt	Average Kerning: -25	

McNeil Submission Artwork

(b) (4)

Drug Facts

Active ingredient (in each caplet) — **Purpose**

Acetaminophen 650 mg ..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - backache
  - minor pain of arthritis
  - toothache
  - premenstrual and menstrual cramps
  - headache
  - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	■ do not use

Other information

- store between 20-25°C (68-77°F)
- **do not use if carton is opened, or if foil inner seal imprinted with "SAFETY SEAL@" is broken**

Inactive ingredients

carnauba wax, hydroxyethylcellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

EXP

00000000

Contains No Aspirin

Care To Recycle™

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

TYLENOL® 8HR MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

HERITAGE OF CARE • TYLENOL • HERITAGE OF CARE

How can we help?
1-877-895-3665

00000000

See New Warning

TYLENOL® 8HR
MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets

00000000

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	80%
Headers:	8.00 pt	Leading:	6.50 pt	Barlines:	2.50 pt		
Subheaders:	6.00 pt	Drug Facts (continued):	8.00 pt	Hairlines:	0.50 pt	Average Kerning:	0

McNeil Submission Artwork

(b) (4)

Reference ID: 3965345

Active ingredient (in each caplet) **Purpose**

Acetaminophen 650 mg..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - backache
  - minor pain of arthritis
  - toothache
  - premenstrual and menstrual cramps
  - headache
  - the common cold
- temporarily reduces fever

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take
- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product. **Ask a doctor before use if you have** liver disease. **Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin. **Stop use and ask a doctor if**
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

TO OPEN: PUSH AND TURN CAP

READ THE LABEL

NDC 50580-309-02

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

TYLENOL® 8HR

MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

*Capsule-Shaped Bi-Layer Tablets

Keep out of reach of children.

Overdose warning In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
- **do not take more than directed (see overdose warning)**

Adults and children 12 years and over
- take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use for more than 10 days unless directed by a doctor

Children under 12 years
- do not use

Other information
- store between 20-25°C (68-77°F)
- **do not use if foil inner seal imprinted with, "SAFETY SEAL®" is broken or missing**

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA ©McNEIL-PPC, INC. 2016

LOT:

EXP.

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	75%
Headers:	6.00 pt	Leading:	6.50 pt	Barlines:	NA pt		
Subheaders:	6.00 pt	Drug Facts (cotinued):	NA pt	Hairlines:	NA pt	Average Kerning:	-25

McNeil Submission Artwork

(b) (4)

Drug Facts

Active ingredient (in each caplet) **Purpose**

Acetaminophen 650 mg..... Pain reliever/fever reducer

Uses
- temporarily relieves minor aches and pains due to:
  - muscular aches
  - backache
  - minor pain of arthritis
  - toothache
  - premenstrual and menstrual cramps
  - headache
  - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	■ take 2 caplets every 8 hours with water ■ swallow whole; do not crush, chew, split or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	■ do not use

Other information

- store between 20-25°C (68-77°F)
- **do not use if carton is opened, or if foil inner seal imprinted with "SAFETY SEAL®" is broken**

Inactive ingredients

ca nauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by:
McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

Contains No Aspirin

Care To Recycle™

TYLENOL® 8HR MUSCLE ACHES & PAIN
Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer
For Up to 8 Hours Relief of Minor Muscle Aches & Pain

Exp:

00000000

How can we help?
1-877-895-3665

NDC 50580-309-03

See New Warning

TYLENOL® 8HR
MUSCLE ACHES & PAIN

Acetaminophen
Extended-release tablets
Pain Reliever / Fever Reducer
For Up to 8 Hours Relief of Minor Muscle Aches & Pain
*Capsule-Shaped Bi-Layer Tablets

Actual Size

225 Caplets*
650 mg each

00000000

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	9.00 pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	90%
Headers:	8.00 pt	Leading:	7.00 pt	Barlines:	2.50 pt		
Subheaders:	6.00 pt	Drug Facts (continued):	8.00 pt	Hairlines:	0.50 pt	Average Kerning:	0

McNeil Submission Artwork

(b) (4)

Reference ID: 3955345

00000000

EXP.
LOT:

Active ingredient (in each caplet) — Purpose

Acetaminophen 650 mg ..... Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
  - muscular aches
  - backache
  - mino pain of arthritis
  - toothache
  - premenstrual and menstrual cramps
  - headache
  - the common cold
- temporarily reduces fever

Warnings

Liver warning This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 6 caplets in 24 hours, which is the maximum daily amount ■ w th other drugs containing acetaminophen ■ 3 o more alcoholic drinks every day while using this p oduct. **Allergy alert** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. **Do not use** ■ with any other drug containing acetaminophen (presc iption or nonpresc iption). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this p oduct. **Ask a doctor before use if you have** liver disease. **Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfa in. **Stop use and ask a doctor if** ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

TO OPEN: PUSH AND TURN CAP

READ THE LABEL

NDC 50580-309-03

DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN

TYLENOL® 8HR

MUSCLE ACHES & PAIN

Acetaminophen Extended-release tablets

Pain Reliever / Fever Reducer

For Up to 8 Hours Relief of Minor Muscle Aches & Pain

225 Caplets*
650 mg each

*Capsule-Shaped Bi-Layer Tablets

If pregnant or breast-feeding, ask a hea th profess ona be ore use.

Keep out of reach of children.

Overdose warning In case of overdose, get medica help or contact a Poison Cont ol Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

Adults and children 12 years and over ■ take 2 caplets every 8 hours with water
- swallow whole; do not crush, chew, split or dissolve
- do not take more than 6 caplets in 24 hours
- do not use fo more than 10 days unless directed by a doctor

Children under 12 years ■ do not use

Other information

- store between 20-25°C (68-77°F)
- **do not use if foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients carnauba wax, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, modified starch, povidone powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments? call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

Distributed by: McNeil Consumer Healthcare
Division of McNEIL-PPC, Inc.
Fort Washington, PA 19034 USA
©McNEIL-PPC, INC. 2016

Labeling Format Information: Fonts: HelveticaDF Cond Medium, Cond Bold, Cond Bold Italic

Drug Facts:	NA pt	Body Text:	6.00 pt	Bullets:	5.00 pt	Horizontal Scale:	90%
Headers:	8.00 pt	Leading:	6.50 pt	Barlines:	NA pt		
Subheaders:	6.00 pt	Drug Facts (cotinued):	NA pt	Hairlines:	NA pt	Average Kerning:	0

McNeil Submission Artwork

(b) (4)

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

KAREN M MAHONEY
08/26/2016