Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg

SOI font: 22 pt Helvetica Bold Condesned Oblique

00000000

(b) (4)

Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg

SOI font: 19.35 pt Helvetica Bold Condesned Oblique

(b) (4)

Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg
Famotidine Tablets 10 mg

SOI font: 19.35 pt Helvetica Bold Condesned Oblique

NDC 16837-872-22

ORIGINAL STRENGTH

Pepcid® AC

Famotidine Tablets 10 mg
Acid Reducer

Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

90 Tablets

7 16837 87290 0

ORIGINAL STRENGTH

Pepcid® AC

90 Tablets

Famotidine ©active ingredient) made in Republic of Ireland.
Processed in Canada.

Distributed by: JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19034 USA
© McNeil Consumer Pharmaceuticals Co. 2016
PEPCID® and PEPCID AC® are registered trademarks of
McNeil QPC

LOT:
EXP.

Drug Facts

Active ingredient (in each tablet) **Purpose**
Famotidine 10 mg ..... Acid reducer

Uses
- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Drug Facts (continued)

Do not use
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness

Drug Facts (continued)

- eat pain or sh outder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- **frequent chest pain**
- **frequent** wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant for breastfeeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Other information
- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients
carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, red iron oxide, talc, titanium dioxide

Drug Facts

Questions or comments?
1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Tips for Managing Heartburn
- Do not lie flat or bend over after eating
- Do not wear tight-fitting clothing around the stomach
- Do not eat before bedtime
- Raise the head of your bed
- Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables
- Eat slowly and avoid big meals
- If overweight, lose weight
- Quit smoking

JUST ONE TABLET prevents and relieves heartburn due to add indigestion brought on by eating and drinking certain foods and beverages.

Do not use if carton is open or if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

(b) (4)

Do not use if this seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

Active ingredient (in each tablet) **Purpose**

Famotidine 10 mg ..... Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

00000000

Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • hear burn with **light headedness, sweating, or dizziness** • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders or lightheadedness • frequent chest pain • frequest wheezing, particularly with heartburn • unexplained weight loss • nausea or vom ting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of ch ldren.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swa low 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Famotidine (active ingredient) made in Republic of Ireland. Processed in Canada

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Hea thcare Division, Fort Washington, PA 19034 USA

©McNeil CPC, 2016 PEPCID® and PEPCID AC® are registered tradema ks of McNeil CPC

ORIGINAL STRENGTH

NDC 16837-672-22

Pepcid® AC

Famotidine Tablets 10 mg Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn Due to Acid Indigestion

90 Tablets

(b) (4)

Do not use if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.
Active ingredient (in each tablet) Purpose
Famotidine 10 mg ..... Acid reducer
Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers. Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn
ORIGINAL STRENGTH
NDC 16837-872-22
Pepcid®
AC
Famotidine Tablets 10 mg Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn Due to Acid Indigestion
90 Tablets
• unexplained weight loss • nausea or vomiting • stomach pain. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor Famotidine (active ingredient) made in Republic of Ireland, Processed in Canada 00000000/000000
EXP.
LOT:
Distributed by: JOHNSON & JOHNSON CONSUMER INC. Skillman, NJ 08558 USA
©J&JCHC, USA 2015
PEPCID and PEPCID AC are registered trademarks

(b) (4)

Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg

SOI font: 22 pt Helvetica Bold Condesned Oblique

(b) (4)

Reference ID: 3942868

Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg

SOI font: 22 pt Helvetica Bold Condesned Oblique

(b) (4)

Reference ID: 3942868

Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg

SOI font: 19.35 pt Helvetica Bold Condesned Oblique

NDC 16837-855-16

MAXIMUM STRENGTH

Pepcid® AC

Famotidine Tablets 20 mg
Acid Reducer

Just One Tablet!
Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

50 Tablets

7 16837 85550 7

Drug Facts (continued)

Other information
- read the directions and warning before use
- keep the carton, It contains sim portant inforrma tion.
- store at20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients
carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?
1-800-755-4008 (toll-free) or 215-273-8755 (collect)

Drug Facts (continued)

- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

frequent chest pain
- frequent heezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting - stomach pain

Stop use and ask a doctor if
- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast feeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Do not use
- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness

Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine 20mg	Acid reducer

Uses
- relieves heartburn associated with acid indigestion an sour stomach
- prevents heartburn associated with acid indigestion an sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Famotidine (active ingredient) made in Republic of Ireland.
Processed in Canada.

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
Skillman, NJ 08558 USA
Fort Washington, PA 19034 USA
©Mchal Consumer Pharmaceuticals Co. 2016
PEPCID® and PEPCID AC® are registered trademarks of McNeil CPC

LOT:
EXP:

Pepcid® AC
MAXIMUM STRENGTH
50 Tablets

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Do not use if carton is open or if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

Tips for Managing Heartburn
- Do not lie flat or bend over after eating - Do not wear tight-fitting clothing around the stomach - Do not eat before bedtime - Raise the head of your bed - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables - Eat slowly and avoid big meals - If overweight, lose weight - Quit smoking

(b) (4)

Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg
Famotidine Tablets 20 mg

SOI font: 19.35 pt Helvetica Bold Condesned Oblique

NDC 16837-855-16

MAXIMUM STRENGTH

Pepcid® AC

Famotidine Tablets 20 mg
Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn
Due to Acid Indigestion

actual size

50 Tablets

Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine 20mg	Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- if you have kidney disease, except under the advice and supervision of a doctor
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness

Drug Facts (continued)

- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- **frequent chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting • stomach pain

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breastfeeding, ask a health professional before use.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- adults and children 12 years and over:
- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 0 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Drug Facts (continued)

Other information

- read the directions and warning before use
- keep the carton. It contains important information.
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, pregelatinized starch, talc, titanium dioxide

Questions or comments?

1-800-755-4008 (toll-free) or 215-273-8755 (collect)

7 16837 85550 7

Famotidine (active ingredient) made in Republic of Ireland.
Processed in Canada.
Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
199 Grandview Road, Skillman, NJ 08558
Fort Washington, PA 19034 USA
©McNeil Consumer Pharmaceuticals Co. 2016
PEPCID® and PEPCID AC® are registered trademarks of McNeil CPC

LOT:
EXP:

MAXIMUM STRENGTH
Pepcid® AC
50 Tablets

JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.

Do not use if carton is open or if foil seal under bottle cap printed "SAFETY SEAL®" is broken or missing.

Tips for Managing Heartburn

- Do not lie flat or bend over after eating • Do not wear tight-fitting clothing around the stomach • Do not eat before bedtime • Raise the head of your bed • Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables • Eat slowly and avoid big meals
- If overweight, lose weight • Quit smoking

(b) (4)

Do not use if foil seal under bottle cap printed "SAFETY SEAL" is broken or missing.

Active ingredient (in each tablet) **Purpose**

Famotidine 20 mg ..... Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings **Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers

Do not use • if you have trouble or pain swallowing food, vomiting with blood or blood (or black) stools. These may be signs of a serious condition. See your doctor. • if you have kidney disease, except under the advice and supervision of a doctor • with other acid reducers. **Ask a doctor before use if you have** • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or high headedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. **Stop use and ask a doctor if** • your heartburn continues or worsens • you need to take this product for more than 14 days. **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 1 hr to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor

Famotidine (active ingredient) made in Republic of Ireland, Processed in Canada

Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**

McNeil Consumer Healthcare Division, Fort Washington, PA 19034 USA

©McNeil CPC 2016 PEPCID® and PEPCID A ® are registered trademarks of McNeil CPC

00000000

MAXIMUM STRENGTH

NDC 16837-855-16

Pepcid®

AC

Famotidine Tablets 20 mg Acid Reducer

Just One Tablet!

Prevents & Relieves Heartburn Due to Acid Indigestion

50 Tablets

(b) (4)

Do not use if foil seal under bottle cap printed "SAFETY SEALS" is broken or missing.
Active ingredient (in each tablet) Purpose
Famotidine 20 mg ..... Acid reducer
Uses • relieves heartburn associated with acid indigestion and sour stomach • prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers
Do not use • if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. • if you have kidney disease, except under the advice and supervision of a doctor • with other acid reducers. Ask a doctor before use if you have • had heartburn over 3 months. This may be a sign of a more serious condition. • heartburn with lightheadedness, sweating, or dizziness • chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness • frequent chest pain • frequent wheezing, particularly with heartburn • unexplained weight loss • nausea or vomiting • stomach pain. Stop use and ask a doctor if • your heartburn continues or worsens • you need to take this product for more than 14 days
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Directions • adults and children 12 years and over: • to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. • to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn • do not use more than 2 tablets in 24 hours • children under 12 years: ask a doctor
Famotidine (active ingredient) made in Republic of Ireland, Processed in Canada
000000000/000000
MAXIMUM STRENGTH
NDC 16837-855-16
Pepcid® AC
Famotidine Tablets 20 mg Acid Reducer
Just One Tablet!
Prevents & Relieves Heartburn Due to Acid Indigestion
50 Tablets
EXP.
LOT:
Distributed by: JOHNSON & JOHNSON CONSUMER INC. Skillman, NJ 08558 USA
©J&JCHC, 2015
Pepcid and PEPCID AC are registered trademarks of J&JCHC.

(b) (4)

[redacted]

Blister @ 100% Scale

Blister @ 200% Scale



[redacted]

(b) (4)

(b) (4)

[redacted]

Blister @ 100% Scale



Blister @ 200% Scale



(b) (4)

[redacted]

(b) (4)

[redacted]

Blister @ 100% Scale



Blister @ 200% Scale



(b) (4)

[redacted]

This is a representation of an electronic record that was signed electronically and this page is the manifestation of the electronic signature.

/s/

VALERIE S PRATT
06/08/2016