



Drug Facts (continued)

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters.
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you child:
 has had stomach ulcers or bleeding problems
 takes a blood thinning (anticoagulant) or steroid drug
 takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use
 if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer right before or after heart surgery
Ask a doctor before use if
 stomach bleeding warning applies to your child
 child has a history of stomach problems, such as heartburn
 child has problems or serious side effects from taking pain relievers or fever reducers
 child has not been drinking fluids child has lost a lot of fluid due to vomiting or diarrhea
 child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
 child has asthma child is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 under a doctor's care for any serious condition taking any other drug
When using this product
 take with food or milk if stomach upset occurs
 the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 child experiences any of the following signs of stomach bleeding
 feels faint, vomits blood, has bloody or black stools
 has stomach pain that does not get better
 the child does not get any relief within first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days
 redness or swelling is present in the painful area any new symptoms appear

Drug Facts (continued)

Directions
 this product does not contain directions or complete warnings for adult use
 do not give more than directed
 shake well before using
 find right dose on chart below. If possible, use weight to dose; otherwise use age.
 mL = milliliter
 measure with the dosing device provided. Do not use with any other device.
 dispense liquid slowly into the child's mouth, toward the inner cheek
 if needed, repeat dose every 6-8 hours
 do not use more than 4 times a day
Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL









Other information
 store between 20-25°C (68-77°F)
 do not use if carton top is imprinted "SAFETY SEAL" or bottle wrap is imprinted "SAFETY SEAL" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing
Inactive ingredients: anhydrous citric acid, carmellose, FD&C red no. 40, flavors, glycerin, polyethylene glycol, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum
Questions or comments?
 Call 1-877-495-3665 (toll-free) or 215-273-8755 (local)

New Information. Read Entire Label
 NDC 50580-100-18
For Ages 6 Mos. to 23 Mos.
Concentrated Motrin® Infants' Drops
 contains **Ibuprofen Oral Suspension (NSAID)**
50 mg per 1.25 mL
Pain Reliever/Fever Reducer
Lasts up to 8 hours
Original Berry Flavor
1/2 fl oz (15 ml)
 Use only with enclosed syringe.



LOT: _____
 EXP: _____
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2016
 www.motrin.com

USA
 000000
 891



(b) (4)



New Information. Read Entire Label
 NDC 50580-198-50
For Ages 6 Mos. to 23 Mos.
Concentrated Motrin®
Infants' Drops
DYE-FREE
 contains **Ibuprofen**
Oral Suspension (NSAID)
50 mg per 1.25 mL
Pain Reliever/Fever Reducer
Lasts up to 8 hours
 Use only with enclosed syringe.
1 fl oz (30 mL)

Concentrated Motrin®
Infants' Drops
DYE-FREE
Concentrated Motrin®
Infants' Drops
 contains **Ibuprofen**
Oral Suspension (NSAID)
50 mg per 1.25 mL
Pain Reliever/Fever Reducer
Non-Staining
Original Berry Flavor
1 fl oz (30 mL)
 Actual Size

Drug Facts (continued)
Warnings
 Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
 Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
 has had stomach ulcers or bleeding problems
 takes a blood thinning (anticoagulant) or steroid drug
 takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use
 if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 right before or after heart surgery
Ask a doctor before use if stomach bleeding/warning applies to your child
 child has a history of stomach problems, such as heartburn
 child has problems or serious side effects from taking pain relievers or fever reducers
 child has not been drinking fluids child has lost a lot of fluid due to vomiting or diarrhea
 child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
 child has asthma child is taking a diuretic
Ask a doctor or pharmacist before use if the child is
 under a doctor's care for any serious condition taking any other drug
When using this product take with food or milk if stomach upset occurs
 the risk of heart attack or stroke may increase if you use more than directed or for longer than directed
Stop use and ask a doctor if
 child experiences any of the following signs of stomach bleeding
 feels faint vomits blood has bloody or black stools
 has stomach pain that does not get better
 the child does not get any relief within first day (24 hours) of treatment
 fever or pain gets worse or lasts more than 3 days
 redness or swelling is present in the painful area any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
Directions
 this product does not contain directions or complete warnings for adult use
 do not give more than directed shake well before using
 find right dose on chart. If possible, use weight to dose; otherwise use age.

Drug Facts (continued)
 mL = mL for
 measure with the dosing device provided. Do not use with any other device.
 dispense liquid slowly into the child's mouth, toward the inner cheek
 if needed, repeat dose every 6-8 hours
 do not use more than 4 times a day
Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
8-23 lbs	12-23 mos	1.875 mL

Other information
 store between 20-25°C (68-77°F)
 do not use if carton is opened or carton tape imprinted "SAFETY SEAL®" or bottle wrap imprinted "SAFETY SEAL®" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing
Inactive ingredients
 anhydrous citric acid, caramel, flavors, glycerin, polyorbital 80, polyethylene glycol, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum
Questions or comments?
 Call 1-877-885-3665 (to free) or 215-273-8755 (toll)
 3 0045-0198-013

LOT: _____
 EXP: _____
 Distributed by: **JOHNSON & JOHNSON CONSUMER INC.**
 McNeil Consumer Healthcare Division
 Fort Washington, PA 19034 USA ©J&JCI 2016
 www.motrin.com

USA
 000000
 887

54,0 mm 111,0 mm 57,0 mm

51,0 mm

THE DOSING CHART AND DIRECTIONS APPEARING ON THE FIRST AND LAST PAGE ARE REPEATED FOR YOUR CONVENIENCE. PLEASE READ AND KEEP ALL WARNINGS AND DIRECTIONS TO ENSURE THE PROPER USE OF THIS CONCENTRATED INFANTS' MOTRIN® PRODUCT.

Important: Read all product information before using. Keep the box for important information. This product is intended for use in children ages 6 months to 23 months of age.

Active Ingredient (In each 1.25 mL)	Purpose
ibuprofen 50 mg (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ mL = milliliter ■ measure with the dosing device provided. Do not use with any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Concentrated Motrin® Infants' Drops
contains ibuprofen Oral Suspension (NSAID) 50 mg per 1.25 mL Pain Reliever/Fever Reducer
Non-Staining Original Berry Flavor
1 fl oz (30 mL)

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Page 2 **Page 7**

57,0 mm 111,0 mm 54,0 mm

51,0 mm

OPEN TO read all Warnings and Directions before use

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ mL = milliliter ■ measure with the dosing device provided. Do not use with any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Concentrated Motrin® Infants' Drops
contains ibuprofen Oral Suspension (NSAID) 50 mg per 1.25 mL Pain Reliever/Fever Reducer
Non-Staining Original Berry Flavor
1 fl oz (30 mL)

Glue Area (glued to base label if only 7 pages needed)

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Page 1 (COVER)

54,0 mm 108,0 mm 54,0 mm

51,0 mm

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ find right dose on chart. If possible, use weight to dose; otherwise use age. ■ mL = milliliter

Dosing Chart

Weight (lb)	Age (mos)	Dose (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Page 4 **Page 5**

54,0 mm 108,0 mm 54,0 mm

51,0 mm

■ measure with the dosing device provided. Do not use with any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek. ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day

Other Information

- store between 20-25°C (68-77°F)
- do not use if bottle wrap imprinted "SAFETY SEAL®" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing

Inactive Ingredients anhydrous citric acid, caramel, flavors, glycerin, polyorbate 80, pregelatinized starch, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?
Call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

Distributed by:
JOHNSON & JOHNSON CONSUMER INC.
McNeil Consumer Healthcare Division
Fort Washington, PA 19094 USA
©J&JCI 2016 www.motrin.com 00000000

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include:
hives ■ facial swelling ■ asthma (wheezing)
shock ■ skin redness ■ rash ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:
has had stomach ulcers or bleeding problems
takes a blood thinning (anticoagulant) or steroid drug (aspirin, ibuprofen, naproxen, or others)
takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
takes more or for a longer time than directed
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Page 6 **Page 3**

13,0 mm 90,5 mm 57,0 mm 20,5 mm

51 mm < 9,5 mm

Cover Label FRONT

LOT: 000000 EXP:

20,5 mm 90,5 mm 57,0 mm 13,0 mm

51 mm < 9,5 mm

Cover Label BACK

54,0 mm 74,5 mm 20,5 mm

51,0 mm

Base Label



(b) (4)