HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use LEXIVA safely and effectively. See full prescribing information for LEXIVA.

LEXIVA (fosamprenavir calcium) tablets, for oral use
LEXIVA (fosamprenavir calcium) oral suspension
Initial U.S. Approval: 2003

RECENT MAJOR CHANGES

Contraindications (4) 09/2016

INDICATIONS AND USAGE

LEXIVA is an HIV protease inhibitor indicated in combination with other antiretroviral agents for the treatment of HIV-1 infection. (1)

DOSAGE AND ADMINISTRATION

- Therapy-naive Adults: LEXIVA 1,400 mg twice daily; LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily; LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily; LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily. (2.1)
- Protease Inhibitor-experienced Adults: LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily. (2.1)
- Pediatric Patients (aged at least 4 weeks to 18 years): Dosage should be calculated based on body weight (kg) and should not exceed adult dose. (2.2)
- Hepatic Impairment: Recommended adjustments for patients with mild, moderate, or severe hepatic impairment. (2.3)

Dosing Considerations

- LEXIVA tablets may be taken with or without food. (2)
- LEXIVA suspension: Adults should take without food; pediatric patients should take with food. (2)

DOSAGE FORMS AND STRENGTHS

- 700-mg tablets(3)
- 50-mg-per-mL oral suspension (3)

CONTRAINDICATIONS

- Hypersensitivity to LEXIVA or amprenavir (e.g., Stevens-Johnson syndrome). (4)
- Drugs highly dependent on CYP3A4 for clearance and for which elevated plasma levels may result in serious and/or life-threatening events. (4)
- Review ritonavir contraindications when used in combination. (4)

WARNINGS AND PRECAUTIONS

- The concomitant use of LEXIVA with ritonavir and certain other drugs may result in known or potentially significant drug interactions. Consult the full prescribing information prior to and during treatment for potential drug interactions. (5.1, 7.3)
- LEXIVA should be discontinued for severe skin reactions including Stevens-Johnson syndrome. (5.2)
- LEXIVA should be used with caution in patients with a known sulfonamide allergy. (5.3)
- Use of higher than approved doses may lead to transaminase elevations. Patients with hepatitis B or C are at increased risk of transaminase elevations. (5.4)
- Patients receiving LEXIVA may develop new onset or exacerbations of diabetes mellitus, hyperglycemia (5.5), immune reconstitution syndrome (5.6), redistribution/accumulation of body fat (5.7), and elevated triglyceride and cholesterol concentrations (5.8). Monitor cholesterol and triglycerides prior to therapy and periodically thereafter.
- Acute hemolytic anemia has been reported with amprenavir. (5.9)
- Hemophilia: Spontaneous bleeding may occur, and additional factor VIII may be required. (5.10)
- Nephrolithiasis: Cases of nephrolithiasis have been reported with fosamprenavir. (5.11)

ADVERSE REACTIONS

- In adults the most common adverse reactions (incidence greater than or equal to 4%) are diarrhea, rash, nausea, vomiting, and headache. (6.1)
- Vomiting and neutropenia were more frequent in pediatrics than in adults. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact ViiV Healthcare at 1-877-844-8872 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Coadministration of LEXIVA with drugs that induce CYP3A4 may decrease amprenavir (active metabolite) concentrations leading to potential loss of virologic activity. (7, 12.3)
- Coadministration with drugs that inhibit CYP3A4 may increase amprenavir concentrations. (7, 12.3)
- Coadministration of LEXIVA or LEXIVA and ritonavir may result in clinically significant interactions with drugs metabolized by CYP3A4. (7)
- Coadministration of LEXIVA and ritonavir may result in clinically significant interactions with drugs metabolized by CYP2D6. (7)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 09/2016

FULL PRESCRIBING INFORMATION: CONTENTS*

FULL PRESCRIBING INFORMATION
1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Adults
2.2 Pediatric Patients (Aged at Least 4 Weeks to 18 Years)
2.3 Patients with Hepatic Impairment
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Risk of Serious Adverse Reactions Due to Drug Interactions
5.2 Skin Reactions
5.3 Sulfa Allergy
5.4 Hepatic Toxicity
5.5 Diabetes/Hyperglycemia
5.6 Immune Reconstitution Syndrome
5.7 Fat Redistribution
5.8 Lipid Elevations
5.9 Hemolytic Anemia
5.10 Patients with Hemophilia
5.11 Nephrolithiasis
5.12 Resistance/Cross-Resistance
6 ADVERSE REACTIONS
6.1 Clinical Trials
6.2 Postmarketing Experience
7 DRUG INTERACTIONS
7.1 Cytochrome P450 Inhibitors and Inducers
7.2 Drugs that Should Not Be Coadministered with LEXIVA
7.3 Established and Other Potentially Significant Drug Interactions
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.3 Nursing Mothers
8.4 Pediatric Use
8.5 Geriatric Use
8.6 Hepatic Impairment
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
12.4 Microbiology
13 NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
14 CLINICAL STUDIES
14.1 Therapy-Naive Adult Trials
14.2 Protease Inhibitor-Experienced Adult Trials
14.3 Pediatric Trials
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

Reference ID: 3986026

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

LEXIVA® is indicated in combination with other antiretroviral agents for the treatment of human immunodeficiency virus (HIV-1) infection.

The following points should be considered when initiating therapy with LEXIVA plus ritonavir in protease inhibitor-experienced patients:

- The protease inhibitor-experienced patient trial was not large enough to reach a definitive conclusion that LEXIVA plus ritonavir and lopinavir plus ritonavir are clinically equivalent *[see Clinical Studies (14.2)]*.
- Once-daily administration of LEXIVA plus ritonavir is not recommended for adult protease inhibitor-experienced patients or any pediatric patients *[see Dosage and Administration (2.1, 2.2), Clinical Studies (14.2, 14.3)]*.
- Dosing of LEXIVA plus ritonavir is not recommended for protease inhibitor-experienced pediatric patients younger than 6 months *[see Clinical Pharmacology (12.3)]*.

2 DOSAGE AND ADMINISTRATION

LEXIVA tablets may be taken with or without food.

Adults should take LEXIVA oral suspension without food. Pediatric patients should take LEXIVA oral suspension with food *[see Clinical Pharmacology (12.3)]*. If emesis occurs within 30 minutes after dosing, re-dosing of LEXIVA oral suspension should occur.

Higher-than-approved dose combinations of LEXIVA plus ritonavir are not recommended due to an increased risk of transaminase elevations *[see Overdosage (10)]*.

When LEXIVA is used in combination with ritonavir, prescribers should consult the full prescribing information for ritonavir.

2.1 Adults

<u>Therapy-Naive Adults</u>

- LEXIVA 1,400 mg twice daily (without ritonavir).
- LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily.
- LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily.
    - Dosing of LEXIVA 1,400 mg once daily plus ritonavir 100 mg once daily is supported by pharmacokinetic data *[see Clinical Pharmacology (12.3)]*.
- LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

Reference ID: 3986026

- Dosing of LEXIVA 700 mg twice daily plus 100 mg ritonavir twice daily is supported by pharmacokinetic and safety data *[see Clinical Pharmacology (12.3)]*.

Protease Inhibitor-Experienced Adults

- LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

2.2 Pediatric Patients (Aged at Least 4 Weeks to 18 Years)

The recommended dosage of LEXIVA in patients aged at least 4 weeks to 18 years should be calculated based on body weight (kg) and should not exceed the recommended adult dose (Table 1).

Table 1. Twice-Daily Dosage Regimens by Weight for Protease Inhibitor-Naive Pediatric Patients (Aged 4 Weeks and Older) and for Protease Inhibitor-Experienced Pediatric Patients (Aged 6 Months and Older) Using LEXIVA Oral Suspension with Concurrent Ritonavir

Weight	Twice-daily Dosage Regimen
<11 kg	LEXIVA 45 mg/kg plus ritonavir 7 mg/kg[a]
11 kg - <15 kg	LEXIVA 30 mg/kg plus ritonavir 3 mg/kg[a]
15 kg - <20 kg	LEXIVA 23 mg/kg plus ritonavir 3 mg/kg[a]
≥20 kg	LEXIVA 18 mg/kg plus ritonavir 3 mg/kg[a]

[a] When dosing with ritonavir, do not exceed the adult dose of LEXIVA 700 mg/ ritonavir 100 mg twice-daily dose.

Alternatively, protease inhibitor-naive children aged 2 years and older can be administered LEXIVA (without ritonavir) 30 mg per kg twice daily.

LEXIVA should only be administered to infants born at 38 weeks gestation or greater and who have attained a post-natal age of 28 days.

For pediatric patients, pharmacokinetic and clinical data:

- do not support once-daily dosing of LEXIVA alone or in combination with ritonavir *[see Clinical Studies (14.3)]*.
- do not support administration of LEXIVA alone or in combination with ritonavir for protease inhibitor-experienced children younger than 6 months *[see Clinical Pharmacology (12.3)]*.
- do not support twice-daily dosing of LEXIVA without ritonavir in pediatric patients younger than 2 years *[see Clinical Pharmacology (12.3)]*.

Other Dosing Considerations

- When administered without ritonavir, the adult regimen of LEXIVA tablets 1,400 mg twice daily may be used for pediatric patients weighing at least 47 kg.

Reference ID: 3986026

- When administered in combination with ritonavir, LEXIVA tablets may be used for pediatric patients weighing at least 39 kg; ritonavir capsules may be used for pediatric patients weighing at least 33 kg.

2.3 Patients with Hepatic Impairment

See Clinical Pharmacology (12.3).

Mild Hepatic Impairment (Child-Pugh Score Ranging from 5 to 6)

LEXIVA should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naive) or 700 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

Moderate Hepatic Impairment (Child-Pugh Score Ranging from 7 to 9)

LEXIVA should be used with caution at a reduced dosage of 700 mg twice daily without ritonavir (therapy-naive), or 450 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

Severe Hepatic Impairment (Child-Pugh Score Ranging from 10 to 15)

LEXIVA should be used with caution at a reduced dosage of 350 mg twice daily without ritonavir (therapy-naive) or 300 mg twice daily plus ritonavir 100 mg once daily (therapy-naive or protease inhibitor-experienced).

There are no data to support dosing recommendations for pediatric patients with hepatic impairment.

3 DOSAGE FORMS AND STRENGTHS

LEXIVA tablets, 700 mg, are pink, film-coated, capsule-shaped, biconvex tablets with "GX LL7" debossed on one face.

LEXIVA oral suspension, 50 mg per mL, is a white to off-white suspension that has a characteristic grape-bubblegum-peppermint flavor.

4 CONTRAINDICATIONS

LEXIVA is contraindicated:

- in patients with previously demonstrated clinically significant hypersensitivity (e.g., Stevens-Johnson syndrome) to any of the components of this product or to amprenavir.
- when coadministered with drugs that are highly dependent on cytochrome P450 3A4 (CYP3A4) for clearance and for which elevated plasma concentrations are associated with serious and/or life-threatening events (Table 2).

Reference ID: 3986026

Table 2. Drugs Contraindicated with LEXIVA (Information in the table applies to LEXIVA with or without ritonavir, unless otherwise indicated.)

Drug Class/Drug Name	Clinical Comment
Alpha 1-adrenoreceptor antagonist: Alfuzosin	Potentially increased alfuzosin concentrations can result in hypotension.
Antiarrhythmics: Flecainide, propafenone	**POTENTIAL** for serious and/or life-threatening reactions such as cardiac arrhythmias secondary to increases in plasma concentrations of antiarrhythmics if LEXIVA is co-prescribed with **ritonavir**.
Antimycobacterials: Rifampin[a]	May lead to loss of virologic response and possible resistance to LEXIVA or to the class of protease inhibitors.
Antipsychotics: Lurasidone Pimozide	**POTENTIAL** for serious and/or life-threatening reactions if LEXIVA is co-administered with **ritonavir**. **POTENTIAL** for serious and/or life-threatening reactions such as cardiac arrhythmias.
Ergot derivatives: Dihydroergotamine, ergonovine, ergotamine, methylergonovine	**POTENTIAL** for serious and/or life-threatening reactions such as acute ergot toxicity characterized by peripheral vasospasm and ischemia of the extremities and other tissues.
GI motility agents: Cisapride	**POTENTIAL** for serious and/or life-threatening reactions such as cardiac arrhythmias.
Herbal products: St. John's wort (*Hypericum perforatum*)	May lead to loss of virologic response and possible resistance to LEXIVA or to the class of protease inhibitors.
HMG co-reductase inhibitors: Lovastatin, simvastatin	**POTENTIAL** for serious reactions such as risk of myopathy including rhabdomyolysis.
Non-nucleoside reverse transcriptase inhibitor: Delavirdine[a]	May lead to loss of virologic response and possible resistance to delavirdine.
PDE5 inhibitor: Sildenafil (REVATIO®) (for treatment of pulmonary arterial hypertension)	A safe and effective dose has not been established when used with LEXIVA. There is increased potential for sildenafil-associated adverse events (which include visual disturbances, hypotension, prolonged erection, and syncope).

Reference ID: 3986026

Sedative/hypnotics: Midazolam, triazolam	**POTENTIAL** for serious and/or life-threatening reactions such as prolonged or increased sedation or respiratory depression.

[a] *See Clinical Pharmacology (12.3) Tables 10, 11, 12, or 13 for magnitude of interaction.*

- when coadministered with ritonavir in patients receiving the antiarrhythmic agents, flecainide and propafenone. If LEXIVA is coadministered with ritonavir, reference should be made to the full prescribing information for ritonavir for additional contraindications.

5 WARNINGS AND PRECAUTIONS

5.1 Risk of Serious Adverse Reactions Due to Drug Interactions

Initiation of LEXIVA/ritonavir, a CYP3A inhibitor, in patients receiving medications metabolized by CYP3A or initiation of medications metabolized by CYP3A in patients already receiving LEXIVA/ritonavir, may increase plasma concentrations of medications metabolized by CYP3A. Initiation of medications that inhibit or induce CYP3A may increase or decrease concentrations of LEXIVA/ritonavir, respectively. These interactions may lead to:

- Clinically significant adverse reactions, potentially leading to severe, life-threatening, or fatal events from greater exposures of concomitant medications.
- Clinically significant adverse reactions from greater exposures of LEXIVA/ritonavir.
- Loss of therapeutic effect of LEXIVA/ritonavir and possible development of resistance.

See Table 7 for steps to prevent or manage these possible and known significant drug interactions, including dosing recommendations *[see Drug Interactions (7)]*. Consider the potential for drug interactions prior to and during LEXIVA/ritonavir therapy; review concomitant medications during LEXIVA/ritonavir therapy; and monitor for the adverse reactions associated with the concomitant medications *[see Contraindications (4), Drug Interactions (7)]*.

5.2 Skin Reactions

Severe and life-threatening skin reactions, including 1 case of Stevens-Johnson syndrome among 700 subjects treated with LEXIVA in clinical trials. Treatment with LEXIVA should be discontinued for severe or life-threatening rashes and for moderate rashes accompanied by systemic symptoms *[see Adverse Reactions (6)]*.

5.3 Sulfa Allergy

LEXIVA should be used with caution in patients with a known sulfonamide allergy. Fosamprenavir contains a sulfonamide moiety. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown. In a clinical trial of LEXIVA used as the sole protease inhibitor, rash occurred in 2 of 10 subjects (20%) with a history of sulfonamide allergy compared with 42 of 126 subjects (33%) with no history of sulfonamide allergy. In 2 clinical trials of LEXIVA plus low-dose ritonavir, rash occurred in 8 of 50 subjects (16%) with

Reference ID: 3986026

a history of sulfonamide allergy compared with 50 of 412 subjects (12%) with no history of sulfonamide allergy.

5.4 Hepatic Toxicity

Use of LEXIVA with ritonavir at higher-than-recommended dosages may result in transaminase elevations and should not be used *[see Dosage and Administration (2), Overdosage (10)]*. Patients with underlying hepatitis B or C or marked elevations in transaminases prior to treatment may be at increased risk for developing or worsening of transaminase elevations. Appropriate laboratory testing should be conducted prior to initiating therapy with LEXIVA and patients should be monitored closely during treatment.

5.5 Diabetes/Hyperglycemia

New onset diabetes mellitus, exacerbation of pre-existing diabetes mellitus, and hyperglycemia have been reported during postmarketing surveillance in HIV-1-infected patients receiving protease inhibitor therapy. Some patients required either initiation or dose adjustments of insulin or oral hypoglycemic agents for treatment of these events. In some cases, diabetic ketoacidosis has occurred. In those patients who discontinued protease inhibitor therapy, hyperglycemia persisted in some cases. Because these events have been reported voluntarily during clinical practice, estimates of frequency cannot be made and causal relationships between protease inhibitor therapy and these events have not been established.

5.6 Immune Reconstitution Syndrome

Immune reconstitution syndrome has been reported in patients treated with combination antiretroviral therapy, including LEXIVA. During the initial phase of combination antiretroviral treatment, patients whose immune systems respond may develop an inflammatory response to indolent or residual opportunistic infections (such as *Mycobacterium avium* infection, cytomegalovirus, *Pneumocystis jirovecii* pneumonia [PCP], or tuberculosis), which may necessitate further evaluation and treatment.

Autoimmune disorders (such as Graves' disease, polymyositis, and Guillain-Barré syndrome) have also been reported to occur in the setting of immune reconstitution; however, the time to onset is more variable, and can occur many months after initiation of treatment.

5.7 Fat Redistribution

Redistribution/accumulation of body fat, including central obesity, dorsocervical fat enlargement (buffalo hump), peripheral wasting, facial wasting, breast enlargement, and "cushingoid appearance," have been observed in patients receiving antiretroviral therapy, including LEXIVA. The mechanism and long-term consequences of these events are currently unknown. A causal relationship has not been established.

Reference ID: 3986026

5.8 Lipid Elevations

Treatment with LEXIVA plus ritonavir has resulted in increases in the concentration of triglycerides and cholesterol *[see Adverse Reactions (6)]*. Triglyceride and cholesterol testing should be performed prior to initiating therapy with LEXIVA and at periodic intervals during therapy. Lipid disorders should be managed as clinically appropriate *[see Drug Interactions (7)]*.

5.9 Hemolytic Anemia

Acute hemolytic anemia has been reported in a patient treated with amprenavir.

5.10 Patients with Hemophilia

There have been reports of spontaneous bleeding in patients with hemophilia A and B treated with protease inhibitors. In some patients, additional factor VIII was required. In many of the reported cases, treatment with protease inhibitors was continued or restarted. A causal relationship between protease inhibitor therapy and these episodes has not been established.

5.11 Nephrolithiasis

Cases of nephrolithiasis were reported during postmarketing surveillance in HIV-1-infected patients receiving LEXIVA. Because these events were reported voluntarily during clinical practice, estimates of frequency cannot be made. If signs or symptoms of nephrolithiasis occur, temporary interruption or discontinuation of therapy may be considered.

5.12 Resistance/Cross-Resistance

Because the potential for HIV cross-resistance among protease inhibitors has not been fully explored, it is unknown what effect therapy with LEXIVA will have on the activity of subsequently administered protease inhibitors. LEXIVA has been studied in patients who have experienced treatment failure with protease inhibitors *[see Clinical Studies (14.2)]*.

6 ADVERSE REACTIONS

- Severe or life-threatening skin reactions have been reported with the use of LEXIVA *[see Warnings and Precautions (5.2)]*.
- The most common moderate to severe adverse reactions in clinical trials of LEXIVA were diarrhea, rash, nausea, vomiting, and headache.
- Treatment discontinuation due to adverse events occurred in 6.4% of subjects receiving LEXIVA and in 5.9% of subjects receiving comparator treatments. The most common adverse reactions leading to discontinuation of LEXIVA (incidence less than or equal to 1% of subjects) included diarrhea, nausea, vomiting, AST increased, ALT increased, and rash.

Reference ID: 3986026

6.1 Clinical Trials

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Adult Trials

The data for the 3 active-controlled clinical trials described below reflect exposure of 700 HIV-1–infected subjects to LEXIVA tablets, including 599 subjects exposed to LEXIVA for greater than 24 weeks, and 409 subjects exposed for greater than 48 weeks. The population age ranged from 17 to 72 years. Of these subjects, 26% were female, 51% white, 31% black, 16% American Hispanic, and 70% were antiretroviral-naive. Sixty-one percent received LEXIVA 1,400 mg once daily plus ritonavir 200 mg once daily; 24% received LEXIVA 1,400 mg twice daily; and 15% received LEXIVA 700 mg twice daily plus ritonavir 100 mg twice daily.

Selected adverse reactions reported during the clinical efficacy trials of LEXIVA are shown in Tables 3 and 4. Each table presents adverse reactions of moderate or severe intensity in subjects treated with combination therapy for up to 48 weeks.

Table 3. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater than or Equal to 2% of Antiretroviral-Naive Adult Subjects

	APV30001[a]		APV30002[a]	
Adverse Reaction	**LEXIVA 1,400 mg b.i.d. (n = 166)**	**Nelfinavir 1,250 mg b.i.d. (n = 83)**	**LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)**	**Nelfinavir 1,250 mg b.i.d. (n = 327)**
Gastrointestinal				
Diarrhea	5%	18%	10%	18%
Nausea	7%	4%	7%	5%
Vomiting	2%	4%	6%	4%
Abdominal pain	1%	0%	2%	2%
Skin				
Rash	8%	2%	3%	2%
General disorders				
Fatigue	2%	1%	4%	2%
Nervous system				
Headache	2%	4%	3%	3%

[a] All subjects also received abacavir and lamivudine twice daily.

Reference ID: 3986026

Table 4. Selected Moderate/Severe Clinical Adverse Reactions Reported in Greater than or Equal to 2% of Protease Inhibitor-Experienced Adult Subjects (Trial APV30003)

Adverse Reaction	LEXIVA 700 mg b.i.d./ Ritonavir 100 mg b.i.d.[a] (n = 106)	Lopinavir 400 mg b.i.d./ Ritonavir 100 mg b.i.d.[a] (n = 103)
Gastrointestinal		
Diarrhea	13%	11%
Nausea	3%	9%
Vomiting	3%	5%
Abdominal pain	<1%	2%
Skin		
Rash	3%	0%
Nervous system		
Headache	4%	2%

[a] All subjects also received 2 reverse transcriptase inhibitors.

Skin rash (without regard to causality) occurred in approximately 19% of subjects treated with LEXIVA in the pivotal efficacy trials. Rashes were usually maculopapular and of mild or moderate intensity, some with pruritus. Rash had a median onset of 11 days after initiation of LEXIVA and had a median duration of 13 days. Skin rash led to discontinuation of LEXIVA in less than 1% of subjects. In some subjects with mild or moderate rash, dosing with LEXIVA was often continued without interruption; if interrupted, reintroduction of LEXIVA generally did not result in rash recurrence.

The percentages of subjects with Grade 3 or 4 laboratory abnormalities in the clinical efficacy trials of LEXIVA are presented in Tables 5 and 6.

Table 5. Grade 3/4 Laboratory Abnormalities Reported in Greater than or Equal to 2% of Antiretroviral-Naive Adult Subjects in Trials APV30001 and APV30002

	APV30001[a]		APV30002[a]	
Laboratory Abnormality	LEXIVA 1,400 mg b.i.d. (n = 166)	Nelfinavir 1,250 mg b.i.d. (n = 83)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)	Nelfinavir 1,250 mg b.i.d. (n = 327)
ALT (>5 x ULN)	6%	5%	8%	8%
AST (>5 x ULN)	6%	6%	6%	7%
Serum lipase (>2 x ULN)	8%	4%	6%	4%
Triglycerides[b] (>750 mg/dL)	0%	1%	6%	2%

Reference ID: 3986026

Neutrophil count, absolute (<750 cells/mm^3)	3%	6%	3%	4%

[a] All subjects also received abacavir and lamivudine twice daily.
[b] Fasting specimens.
ULN = Upper limit of normal.

The incidence of Grade 3 or 4 hyperglycemia in antiretroviral-naive subjects who received LEXIVA in the pivotal trials was less than 1%.

Table 6. Grade 3/4 Laboratory Abnormalities Reported in Greater than or Equal to 2% of Protease Inhibitor-Experienced Adult Subjects in Trial APV30003

Laboratory Abnormality	LEXIVA 700 mg b.i.d./ Ritonavir 100 mg b.i.d.[a] (n = 104)	Lopinavir 400 mg b.i.d./ Ritonavir 100 mg b.i.d.[a] (n = 103)
Triglycerides[b] (>750 mg/dL)	11%[c]	6%[c]
Serum lipase (>2 x ULN)	5%	12%
ALT (>5 x ULN)	4%	4%
AST (>5 x ULN)	4%	2%
Glucose (>251 mg/dL)	2%[c]	2%[c]

[a] All subjects also received 2 reverse transcriptase inhibitors.
[b] Fasting specimens.
[c] n = 100 for LEXIVA plus ritonavir, n = 98 for lopinavir plus ritonavir.
ULN = Upper limit of normal.

Pediatric Trials

LEXIVA with and without ritonavir was studied in 237 HIV-1–infected pediatric subjects aged at least 4 weeks to 18 years in 3 open-label trials; APV20002, APV20003, and APV29005 *[see Clinical Studies (14.3)]*. Vomiting and neutropenia occurred more frequently in pediatric subjects compared with adults. Other adverse events occurred with similar frequency in pediatric subjects compared with adults.

The frequency of vomiting among pediatric subjects receiving LEXIVA twice daily with ritonavir was 20% in subjects aged at least 4 weeks to younger than 2 years and 36% in subjects aged 2 to 18 years compared with 10% in adults. The frequency of vomiting among pediatric subjects receiving LEXIVA twice daily without ritonavir was 60% in subjects aged 2 to 5 years compared with 16% in adults.

The median duration of drug-related vomiting episodes in APV29005 was 1 day (range: 1 to 3 days), in APV20003 was 16 days (range: 1 to 38 days), and in APV20002 was 9 days (range: 4 to 13 days). Vomiting was treatment limiting in 4 pediatric subjects across all 3 trials.

The incidence of Grade 3 or 4 neutropenia (neutrophils less than 750 cells per mm^3) seen in pediatric subjects treated with LEXIVA with and without ritonavir was higher (15%) than the

Reference ID: 3986026

incidence seen in adult subjects (3%). Grade 3/4 neutropenia occurred in 10% (5 of 51) of subjects aged at least 4 weeks to younger than 2 years and 16% (28 of 170) of subjects aged 2 to 18 years.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of LEXIVA. Because these reactions are reported voluntarily from a population of unknown size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. These reactions have been chosen for inclusion due to a combination of their seriousness, frequency of reporting, or potential causal connection to LEXIVA.

Cardiac Disorders

Myocardial infarction.

Metabolism and Nutrition Disorders

Hypercholesterolemia.

Nervous System Disorders

Oral paresthesia.

Skin and Subcutaneous Tissue Disorders

Angioedema.

Urogenital

Nephrolithiasis.

7 DRUG INTERACTIONS

See also Contraindications (4), Clinical Pharmacology (12.3).

If LEXIVA is used in combination with ritonavir, see full prescribing information for ritonavir for additional information on drug interactions.

7.1 Cytochrome P450 Inhibitors and Inducers

Amprenavir, the active metabolite of fosamprenavir, is an inhibitor of CYP3A4 metabolism and therefore should not be administered concurrently with medications with narrow therapeutic windows that are substrates of CYP3A4. Data also suggest that amprenavir induces CYP3A4.

Amprenavir is metabolized by CYP3A4. Coadministration of LEXIVA and drugs that induce CYP3A4, such as rifampin, may decrease amprenavir concentrations and reduce its therapeutic effect. Coadministration of LEXIVA and drugs that inhibit CYP3A4 may increase amprenavir concentrations and increase the incidence of adverse effects.

The potential for drug interactions with LEXIVA changes when LEXIVA is coadministered with the potent CYP3A4 inhibitor ritonavir. The magnitude of CYP3A4-mediated drug interactions

Reference ID: 3986026

(effect on amprenavir or effect on coadministered drug) may change when LEXIVA is coadministered with ritonavir. Because ritonavir is a CYP2D6 inhibitor, clinically significant interactions with drugs metabolized by CYP2D6 are possible when coadministered with LEXIVA plus ritonavir.

There are other agents that may result in serious and/or life-threatening drug interactions *[see Contraindications (4)]*.

7.2 Drugs that Should Not Be Coadministered with LEXIVA

See Contraindications (4).

7.3 Established and Other Potentially Significant Drug Interactions

Table 7 provides a listing of established or potentially clinically significant drug interactions. Information in the table applies to LEXIVA with or without ritonavir, unless otherwise indicated.

Table 7. Established and Other Potentially Significant Drug Interactions

Concomitant Drug Class: Drug Name	Effect on Concentration of Amprenavir or Concomitant Drug	Clinical Comment
HCV/HIV-Antiviral Agents		
HCV protease inhibitor: Boceprevir	**LEXIVA:** ↓Amprenavir (predicted) ↔ or ↓Boceprevir (predicted) **LEXIVA/ritonavir:** ↓Amprenavir (predicted) ↓Boceprevir (predicted)	Coadministration of LEXIVA or LEXIVA/ritonavir and boceprevir is not recommended.

Reference ID: 3986026

HCV protease inhibitor: Simeprevir	**LEXIVA:** ↔Amprenavir (predicted) ↑ or ↓Simeprevir (predicted) **LEXIVA/ritonavir:** ↔Amprenavir (predicted) ↑Simeprevir (predicted)	Coadministration of LEXIVA or LEXIVA/ritonavir and simeprevir is not recommended.
HCV protease inhibitor: Paritaprevir (coformulated with ritonavir and ombitasvir and coadministered with dasabuvir)	**LEXIVA:** ↑Amprenavir (predicted) ↑ or ↔Paritaprevir (predicted) **LEXIVA/ritonavir:** ↑ or ↔Amprenavir (predicted) ↑Paritaprevir (predicted)	Appropriate doses of the combinations with respect to safety and efficacy have not been established. LEXIVA 1400 mg once daily may be considered when coadministered with paritaprevir/ritonavir/ombitasvir/ dasabuvir. Coadministration of LEXIVA/ritonavir and paritaprevir/ritonavir/ombitasvir/ dasabuvir is not recommended.
Non-nucleoside reverse transcriptase inhibitor: Efavirenz[a]	**LEXIVA:** ↓Amprenavir **LEXIVA/ritonavir:** ↓Amprenavir	Appropriate doses of the combinations with respect to safety and efficacy have not been established. An additional 100 mg/day (300 mg total) of ritonavir is recommended when efavirenz is administered with LEXIVA/ritonavir once daily. No change in the ritonavir dose is required when efavirenz is administered with LEXIVA plus ritonavir twice daily.
Non-nucleoside reverse transcriptase inhibitor: Nevirapine[a]	**LEXIVA:** ↓Amprenavir ↑Nevirapine **LEXIVA/ritonavir:** ↓Amprenavir ↑Nevirapine	Coadministration of nevirapine and LEXIVA without ritonavir is not recommended. No dosage adjustment required when nevirapine is administered with LEXIVA/ritonavir twice daily. The combination of nevirapine

Reference ID: 3986026

		administered with LEXIVA/ritonavir once-daily regimen has not been studied.
HIV protease inhibitor: Atazanavir[a]	**LEXIVA:** Interaction has not been evaluated. **LEXIVA/ritonavir:** ↓Atazanavir ↔Amprenavir	Appropriate doses of the combinations with respect to safety and efficacy have not been established.
HIV protease inhibitors: Indinavir[a], nelfinavir[a]	**LEXIVA:** ↑Amprenavir Effect on indinavir and nelfinavir is not well established. **LEXIVA/ritonavir:** Interaction has not been evaluated.	Appropriate doses of the combinations with respect to safety and efficacy have not been established.
HIV protease inhibitors: Lopinavir/ritonavir[a]	↓Amprenavir ↓Lopinavir	An increased rate of adverse events has been observed. Appropriate doses of the combinations with respect to safety and efficacy have not been established.
HIV protease inhibitor: Saquinavir[a]	**LEXIVA:** ↓Amprenavir Effect on saquinavir is not well established. **LEXIVA/ritonavir:** Interaction has not been evaluated.	Appropriate doses of the combination with respect to safety and efficacy have not been established.
HIV integrase inhibitor: Raltegravir[a]	**LEXIVA:** ↓Amprenavir ↓Raltegravir **LEXIVA/ritonavir:** ↓Amprenavir ↓Raltegravir	Appropriate doses of the combination with respect to safety and efficacy have not been established.
HIV integrase inhibitor: Dolutegravir[a]	**LEXIVA/ritonavir:** ↓Dolutegravir	The recommended dose of dolutegravir is 50 mg twice daily when coadministered with LEXIVA/ritonavir.

Reference ID: 3986026

		Use an alternative combination where possible in patients with known or suspected integrase inhibitor resistance.
HIV CCR5 co-receptor antagonist: Maraviroc[a]	**LEXIVA/ritonavir:** ↓Amprenavir ↑Maraviroc	No dosage adjustment required for LEXIVA/ritonavir. The recommended dose of maraviroc is 150 mg twice daily when coadministered with LEXIVA/ritonavir. LEXIVA should be given with ritonavir when coadministered with maraviroc.
Other Agents		
Antiarrhythmics: Amiodarone, lidocaine (systemic), and quinidine	↑Antiarrhythmics	Use with caution. Increased exposure may be associated with life-threatening reactions such as cardiac arrhythmias. Therapeutic concentration monitoring, if available, is recommended for antiarrhythmics.
Anticoagulant: Warfarin		Concentrations of warfarin may be affected. It is recommended that INR (international normalized ratio) be monitored.
Anticonvulsants: Carbamazepine, phenobarbital, phenytoin	**LEXIVA:** ↓Amprenavir	Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations in patients taking these agents concomitantly.
Phenytoin[a]	**LEXIVA/ritonavir:** ↑Amprenavir ↓Phenytoin	Plasma phenytoin concentrations should be monitored and phenytoin dose should be increased as appropriate. No change in LEXIVA/ritonavir dose is recommended.
Antidepressant: Paroxetine, trazodone	↓Paroxetine	Any paroxetine dose adjustment should be guided by clinical effect (tolerability and efficacy).
	↑Trazodone	Adverse events of nausea, dizziness, hypotension, and syncope have been observed following coadministration of trazodone and ritonavir. If trazodone is used with a CYP3A4 inhibitor such as LEXIVA, the combination should be used with caution and a lower dose of

Reference ID: 3986026

		trazodone should be considered.
Antifungals: Ketoconazole[a], itraconazole	↑Ketoconazole ↑Itraconazole	Increase monitoring for adverse events. **LEXIVA:** Dose reduction of ketoconazole or itraconazole may be needed for patients receiving more than 400 mg ketoconazole or itraconazole per day. **LEXIVA/ritonavir:** High doses of ketoconazole or itraconazole (greater than 200 mg/day) are not recommended.
Anti-gout: Colchicine	↑Colchicine	Patients with renal or hepatic impairment should not be given colchicine with LEXIVA/ritonavir. **LEXIVA/ritonavir and coadministration of colchicine:** **Treatment of gout flares:** 0.6 mg (1 tablet) x 1 dose, followed by 0.3 mg (half tablet) 1 hour later. Dose to be repeated no earlier than 3 days. **Prophylaxis of gout flares:** If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once every other day. **Treatment of familial Mediterranean fever (FMF):** Maximum daily dose of 0.6 mg (may be given as 0.3 mg twice a day). **LEXIVA and coadministration of colchicine:** **Treatment of gout flares:** 1.2 mg (2 tablets) x 1 dose. Dose to be repeated no earlier than 3 days.

Reference ID: 3986026

		Prophylaxis of gout flares: If the original regimen was 0.6 mg twice a day, the regimen should be adjusted to 0.3 mg twice a day or 0.6 mg once a day. If the original regimen was 0.6 mg once a day, the regimen should be adjusted to 0.3 mg once a day. **Treatment of FMF:** Maximum daily dose of 1.2 mg (may be given as 0.6 mg twice a day).
Antimycobacterial: Rifabutin[a]	↑Rifabutin and rifabutin metabolite	A complete blood count should be performed weekly and as clinically indicated to monitor for neutropenia. **LEXIVA:** A dosage reduction of rifabutin by at least half the recommended dose is required. **LEXIVA/ritonavir:** Dosage reduction of rifabutin by at least 75% of the usual dose of 300 mg/day is recommended (a maximum dose of 150 mg every other day or 3 times per week).
Antipsychotics: Quetiapine	**LEXIVA/ritonavir:** ↑Quetiapine	Initiation of LEXIVA with ritonavir in patients taking quetiapine: Consider alternative antiretroviral therapy to avoid increases in quetiapine drug exposures. If coadministration is necessary, reduce the quetiapine dose to 1/6 of the current dose and monitor for quetiapine-associated adverse reactions. Refer to the quetiapine prescribing information for recommendations on adverse reaction monitoring. Initiation of quetiapine in patients taking LEXIVA with ritonavir: Refer to the quetiapine prescribing information for initial dosing and titration

Reference ID: 3986026

Lurasidone	↑Lurasidone	of quetiapine. **LEXIVA:** If coadministration is necessary, reduce the lurasidone dose. Refer to the lurasidone prescribing information for concomitant use with moderate CYP3A4 inhibitors. **LEXIVA/ritonavir**: Use of lurasidone is contraindicated.
Benzodiazepines: Alprazolam, clorazepate, diazepam, flurazepam	↑Benzodiazepines	Clinical significance is unknown. A decrease in benzodiazepine dose may be needed.
Calcium channel blockers: Diltiazem, felodipine, nifedipine, nicardipine, nimodipine, verapamil, amlodipine, nisoldipine, isradipine	↑Calcium channel blockers	Use with caution. Clinical monitoring of patients is recommended.
Corticosteroid: Dexamethasone	↓Amprenavir	Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations.
Endothelin-receptor antagonists: Bosentan	↑Bosentan	Coadministration of bosentan in patients on LEXIVA: In patients who have been receiving LEXIVA for at least 10 days, start bosentan at 62.5 mg once daily or every other day based upon individual tolerability. Coadministration of LEXIVA in patients on bosentan: Discontinue use of bosentan at least 36 hours prior to initiation of LEXIVA. After at least 10 days following the initiation of LEXIVA, resume bosentan at 62.5 mg once daily or every other day based upon individual tolerability.

Reference ID: 3986026

Histamine H_2-receptor antagonists: Cimetidine, famotidine, nizatidine, ranitidine[a]	**LEXIVA:** ↓Amprenavir **LEXIVA/ritonavir:** Interaction not evaluated	Use with caution. LEXIVA may be less effective due to decreased amprenavir plasma concentrations.
HMG-CoA reductase inhibitors: Atorvastatin[a]	↑Atorvastatin	Titrate atorvastatin dose carefully and use the lowest necessary dose; do not exceed atorvastatin 20 mg/day.
Immunosuppressants: Cyclosporine, tacrolimus, sirolimus	↑Immunosuppressants	Therapeutic concentration monitoring is recommended for immunosuppressant agents.
Inhaled beta-agonist: Salmeterol	↑Salmeterol	Concurrent administration of salmeterol with LEXIVA is not recommended. The combination may result in increased risk of cardiovascular adverse events associated with salmeterol, including QT prolongation, palpitations, and sinus tachycardia.
Inhaled/nasal steroid: Fluticasone	**LEXIVA:** ↑Fluticasone **LEXIVA/ritonavir:** ↑Fluticasone	Use with caution. Consider alternatives to fluticasone, particularly for long-term use. May result in significantly reduced serum cortisol concentrations. Systemic corticosteroid effects including Cushing's syndrome and adrenal suppression have been reported during postmarketing use in patients receiving ritonavir and inhaled or intranasally administered fluticasone. Coadministration of fluticasone and LEXIVA/ritonavir is not recommended unless the potential benefit to the patient outweighs the risk of systemic corticosteroid side effects.
Narcotic analgesic: Methadone	↓Methadone	Data suggest that the interaction is not clinically relevant; however, patients should be monitored for opiate withdrawal symptoms.
Oral contraceptives: Ethinyl estradiol/ norethindrone[a]	**LEXIVA:**	Alternative methods of non-hormonal contraception are recommended. May lead to loss of virologic response.[a]

Reference ID: 3986026

	↓Amprenavir ↓Ethinyl estradiol **LEXIVA/ritonavir:** ↓Ethinyl estradiol	Increased risk of transaminase elevations. No data are available on the use of LEXIVA/ritonavir with other hormonal therapies, such as hormone replacement therapy (HRT) for postmenopausal women.
PDE5 inhibitors: Sildenafil, tadalafil, vardenafil	↑Sildenafil ↑Tadalafil ↑Vardenafil	May result in an increase in PDE5 inhibitor-associated adverse events, including hypotension, syncope, visual disturbances, and priapism. Use of PDE5 inhibitors for pulmonary arterial hypertension (PAH): • Use of sildenafil (REVATIO) is contraindicated when used for the treatment of PAH *[see Contraindications (4)]*. • The following dose adjustments are recommended for use of tadalafil (ADCIRCA®) with LEXIVA: Coadministration of ADCIRCA in patients on LEXIVA: In patients receiving LEXIVA for at least one week, start ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual tolerability. Coadministration of LEXIVA in patients on ADCIRCA: Avoid use of ADCIRCA during the initiation of LEXIVA. Stop ADCIRCA at least 24 hours prior to starting LEXIVA. After at least one week following the initiation of LEXIVA, resume ADCIRCA at 20 mg once daily. Increase to 40 mg once daily based upon individual

Reference ID: 3986026

		tolerability. <u>Use of PDE5 inhibitors for erectile dysfunction:</u> **LEXIVA:** Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 24 hours. **LEXIVA/ritonavir:** Sildenafil: 25 mg every 48 hours. Tadalafil: no more than 10 mg every 72 hours. Vardenafil: no more than 2.5 mg every 72 hours. Use with increased monitoring for adverse events.
Proton pump inhibitors: Esomeprazole[a], lansoprazole, omeprazole, pantoprazole, rabeprazole	**LEXIVA:** ↔Amprenavir ↑Esomeprazole **LEXIVA/ritonavir:** ↔Amprenavir ↔Esomeprazole	Proton pump inhibitors can be administered at the same time as a dose of LEXIVA with no change in plasma amprenavir concentrations.
Tricyclic antidepressants: Amitriptyline, imipramine	↑Tricyclics	Therapeutic concentration monitoring is recommended for tricyclic antidepressants.

[a] *See Clinical Pharmacology (12.3) Tables 10, 11, 12, or 13 for magnitude of interaction.*

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C. Embryo/fetal development studies were conducted in rats (dosed from Day 6 to Day 17 of gestation) and rabbits (dosed from Day 7 to Day 20 of gestation). Administration of fosamprenavir to pregnant rats and rabbits produced no major effects on embryo-fetal development; however, the incidence of abortion was increased in rabbits that were administered fosamprenavir. Systemic exposures ($AUC_{0\text{-}24\ h}$) to amprenavir at these dosages were 0.8 (rabbits) to 2 (rats) times the exposures in humans following administration of the maximum recommended human dose (MRHD) of fosamprenavir alone or 0.3 (rabbits) to 0.7 (rats) times the exposures in humans following administration of the MRHD of fosamprenavir in

Reference ID: 3986026

combination with ritonavir. In contrast, administration of amprenavir was associated with abortions and an increased incidence of minor skeletal variations resulting from deficient ossification of the femur, humerus, and trochlea, in pregnant rabbits at the tested dose approximately one-twentieth the exposure seen at the recommended human dose.

The mating and fertility of the F_1 generation born to female rats given fosamprenavir was not different from control animals; however, fosamprenavir did cause a reduction in both pup survival and body weights. Surviving F_1 female rats showed an increased time to successful mating, an increased length of gestation, a reduced number of uterine implantation sites per litter, and reduced gestational body weights compared with control animals. Systemic exposure ($AUC_{0\text{-}24\,h}$) to amprenavir in the F_0 pregnant rats was approximately 2 times higher than exposures in humans following administration of the MRHD of fosamprenavir alone or approximately the same as those seen in humans following administration of the MRHD of fosamprenavir in combination with ritonavir.

There are no adequate and well-controlled studies in pregnant women. LEXIVA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Antiretroviral Pregnancy Registry

To monitor maternal-fetal outcomes of pregnant women exposed to LEXIVA, an Antiretroviral Pregnancy Registry has been established. Physicians are encouraged to register patients by calling 1-800-258-4263.

8.3 Nursing Mothers

The Centers for Disease Control and Prevention recommend that HIV-1-infected mothers not breastfeed their infants to avoid risking postnatal transmission of HIV-1 infection. Although it is not known if amprenavir is excreted in human milk, amprenavir is secreted into the milk of lactating rats. Because of both the potential for HIV-1 transmission and the potential for serious adverse reactions in nursing infants, mothers should be instructed not to breastfeed if they are receiving LEXIVA.

8.4 Pediatric Use

The safety, pharmacokinetic profile, virologic, and immunologic responses of LEXIVA with and without ritonavir were evaluated in protease inhibitor-naive and -experienced HIV-1–infected pediatric subjects aged at least 4 weeks to younger than 18 years and weighing at least 3 kg in 3 open-label trials *[see Adverse Reactions (6.1), Clinical Pharmacology (12.3), Clinical Studies (14.3)]*.

Treatment with LEXIVA is not recommended in protease inhibitor-experienced pediatric patients younger than 6 months. The pharmacokinetics, safety, tolerability, and efficacy of LEXIVA in pediatric patients younger than 4 weeks have not been established *[see Clinical Pharmacology (12.3)]*. Available pharmacokinetic and clinical data do not support once-daily dosing of LEXIVA alone or in combination with ritonavir for any pediatrics or twice-daily

Reference ID: 3986026

dosing without ritonavir in pediatric patients younger than 2 years *[see Clinical Pharmacology (12.3), Clinical Studies (14.3)]*. See *Dosage and Administration (2.2)* for dosing recommendations for pediatric patients.

8.5 Geriatric Use

Clinical studies of LEXIVA did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger adults. In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

8.6 Hepatic Impairment

Amprenavir is principally metabolized by the liver; therefore, caution should be exercised when administering LEXIVA to patients with hepatic impairment because amprenavir concentrations may be increased *[see Clinical Pharmacology (12.3)]*. Patients with impaired hepatic function receiving LEXIVA with or without concurrent ritonavir require dose reduction *[see Dosage and Administration (2.3)]*.

There are no data to support dosing recommendations for pediatric subjects with hepatic impairment.

10 OVERDOSAGE

In a healthy volunteer repeat-dose pharmacokinetic trial evaluating high-dose combinations of LEXIVA plus ritonavir, an increased frequency of Grade 2/3 ALT elevations (greater than 2.5 x ULN) was observed with LEXIVA 1,400 mg twice daily plus ritonavir 200 mg twice daily (4 of 25 subjects). Concurrent Grade 1/2 elevations in AST (greater than 1.25 x ULN) were noted in 3 of these 4 subjects. These transaminase elevations resolved following discontinuation of dosing.

There is no known antidote for LEXIVA. It is not known whether amprenavir can be removed by peritoneal dialysis or hemodialysis, although it is unlikely as amprenavir is highly protein bound. If overdosage occurs, the patient should be monitored for evidence of toxicity and standard supportive treatment applied as necessary.

11 DESCRIPTION

LEXIVA (fosamprenavir calcium) is a prodrug of amprenavir, an inhibitor of HIV protease. The chemical name of fosamprenavir calcium is (3*S*)-tetrahydrofuran-3-yl (1*S*,2*R*)-3-[[(4-aminophenyl) sulfonyl](isobutyl)amino]-1-benzyl-2-(phosphonooxy) propylcarbamate monocalcium salt. Fosamprenavir calcium is a single stereoisomer with the (3*S*)(1*S*,2*R*) configuration. It has a molecular formula of $C_{25}H_{34}CaN_3O_9PS$ and a molecular weight of 623.7. It has the following structural formula:

Reference ID: 3986026



Fosamprenavir calcium is a white to cream-colored solid with a solubility of approximately 0.31 mg per mL in water at 25°C.

LEXIVA tablets are available for oral administration in a strength of 700 mg of fosamprenavir as fosamprenavir calcium (equivalent to approximately 600 mg of amprenavir). Each 700-mg tablet contains the inactive ingredients colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone K30. The tablet film-coating contains the inactive ingredients hypromellose, iron oxide red, titanium dioxide, and triacetin.

LEXIVA oral suspension is available in a strength of 50 mg per mL of fosamprenavir as fosamprenavir calcium equivalent to approximately 43 mg of amprenavir. LEXIVA oral suspension is a white to off-white suspension with a grape-bubblegum-peppermint flavor. Each one milliliter (1 mL) contains the inactive ingredients artificial grape-bubblegum flavor, calcium chloride dihydrate, hypromellose, methylparaben, natural peppermint flavor, polysorbate 80, propylene glycol, propylparaben, purified water, and sucralose.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Fosamprenavir is an antiviral agent *[see Microbiology (12.4)]*.

12.3 Pharmacokinetics

The pharmacokinetic properties of amprenavir after administration of LEXIVA, with or without ritonavir, have been evaluated in both healthy adult volunteers and in HIV-1-infected subjects; no substantial differences in steady-state amprenavir concentrations were observed between the 2 populations.

The pharmacokinetic parameters of amprenavir after administration of LEXIVA (with and without concomitant ritonavir) are shown in Table 8.

Reference ID: 3986026

Table 8. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters in Adults

Regimen	C_{max} (mcg/mL)	T_{max} (hours)[a]	AUC_{24} (mcg•h/mL)	C_{min} (mcg/mL)
LEXIVA 1,400 mg b.i.d.	4.82 (4.06-5.72)	1.3 (0.8-4.0)	33.0 (27.6-39.2)	0.35 (0.27-0.46)
LEXIVA 1,400 mg q.d. plus Ritonavir 200 mg q.d.	7.24 (6.32-8.28)	2.1 (0.8-5.0)	69.4 (59.7-80.8)	1.45 (1.16-1.81)
LEXIVA 1,400 mg q.d. plus Ritonavir 100 mg q.d.	7.93 (7.25-8.68)	1.5 (0.75-5.0)	66.4 (61.1-72.1)	0.86 (0.74-1.01)
LEXIVA 700 mg b.i.d. plus Ritonavir 100 mg b.i.d.	6.08 (5.38-6.86)	1.5 (0.75-5.0)	79.2 (69.0-90.6)	2.12 (1.77-2.54)

[a] Data shown are median (range).

The mean plasma amprenavir concentrations of the dosing regimens over the dosing intervals are displayed in Figure 1.

Figure 1. Mean (±SD) Steady-State Plasma Amprenavir Concentrations and Mean EC_{50} Values against HIV from Protease Inhibitor-Naive Subjects (in the Absence of Human Serum)



Reference ID: 3986026

Absorption and Bioavailability

After administration of a single dose of LEXIVA to HIV-1–infected subjects, the time to peak amprenavir concentration (T_{max}) occurred between 1.5 and 4 hours (median 2.5 hours). The absolute oral bioavailability of amprenavir after administration of LEXIVA in humans has not been established.

After administration of a single 1,400-mg dose in the fasted state, LEXIVA oral suspension (50 mg per mL) and LEXIVA tablets (700 mg) provided similar amprenavir exposures (AUC); however, the C_{max} of amprenavir after administration of the suspension formulation was 14.5% higher compared with the tablet.

Amprenavir is both a substrate for and inducer of P-glycoprotein.

Effects of Food on Oral Absorption

Administration of a single 1,400-mg dose of LEXIVA tablets in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with no significant changes in amprenavir C_{max}, T_{max}, or $AUC_{0-\infty}$ *[see Dosage and Administration (2)]*.

Administration of a single 1,400-mg dose of LEXIVA oral suspension in the fed state (standardized high-fat meal: 967 kcal, 67 grams fat, 33 grams protein, 58 grams carbohydrate) compared with the fasted state was associated with a 46% reduction in C_{max}, a 0.72-hour delay in T_{max}, and a 28% reduction in amprenavir $AUC_{0-\infty}$.

Distribution

In vitro, amprenavir is approximately 90% bound to plasma proteins, primarily to $alpha_1$-acid glycoprotein. In vitro, concentration-dependent binding was observed over the concentration range of 1 to 10 mcg per mL, with decreased binding at higher concentrations. The partitioning of amprenavir into erythrocytes is low, but increases as amprenavir concentrations increase, reflecting the higher amount of unbound drug at higher concentrations.

Metabolism

After oral administration, fosamprenavir is rapidly and almost completely hydrolyzed to amprenavir and inorganic phosphate prior to reaching the systemic circulation. This occurs in the gut epithelium during absorption. Amprenavir is metabolized in the liver by the CYP3A4 enzyme system. The 2 major metabolites result from oxidation of the tetrahydrofuran and aniline moieties. Glucuronide conjugates of oxidized metabolites have been identified as minor metabolites in urine and feces.

Elimination

Excretion of unchanged amprenavir in urine and feces is minimal. Unchanged amprenavir in urine accounts for approximately 1% of the dose; unchanged amprenavir was not detectable in feces. Approximately 14% and 75% of an administered single dose of ^{14}C-amprenavir can be

Reference ID: 3986026

accounted for as metabolites in urine and feces, respectively. Two metabolites accounted for greater than 90% of the radiocarbon in fecal samples. The plasma elimination half-life of amprenavir is approximately 7.7 hours.

Special Populations

Hepatic Impairment: The pharmacokinetics of amprenavir have been studied after the administration of LEXIVA in combination with ritonavir to adult HIV-1–infected subjects with mild, moderate, and severe hepatic impairment. Following 2 weeks of dosing with LEXIVA plus ritonavir, the AUC of amprenavir was increased by approximately 22% in subjects with mild hepatic impairment, by approximately 70% in subjects with moderate hepatic impairment, and by approximately 80% in subjects with severe hepatic impairment compared with HIV-1–infected subjects with normal hepatic function. Protein binding of amprenavir was decreased in subjects with hepatic impairment. The unbound fraction at 2 hours (approximate C_{max}) ranged between a decrease of -7% to an increase of 57% while the unbound fraction at the end of the dosing interval (C_{min}) increased from 50% to 102% *[see Dosage and Administration (2.3)]*.

The pharmacokinetics of amprenavir have been studied after administration of amprenavir given as AGENERASE® capsules to adult subjects with hepatic impairment. Following administration of a single 600-mg oral dose, the AUC of amprenavir was increased by approximately 2.5-fold in subjects with moderate cirrhosis and by approximately 4.5-fold in subjects with severe cirrhosis compared with healthy volunteers *[see Dosage and Administration (2.3)]*.

Renal Impairment: The impact of renal impairment on amprenavir elimination in adults has not been studied. The renal elimination of unchanged amprenavir represents approximately 1% of the administered dose; therefore, renal impairment is not expected to significantly impact the elimination of amprenavir.

Pediatric Patients: The pharmacokinetics of amprenavir following administration of LEXIVA oral suspension and LEXIVA tablets, with or without ritonavir, have been studied in a total of 212 HIV-1–infected pediatric subjects enrolled in 3 trials. LEXIVA without ritonavir was administered as 30 or 40 mg per kg twice daily to children aged 2 to 5 years. LEXIVA with ritonavir was administered as LEXIVA 30 mg per kg plus ritonavir 6 mg per kg once daily to children aged 2 to 18 years and as LEXIVA 18 to 60 mg per kg plus ritonavir 3 to 10 mg per kg twice daily to children aged at least 4 weeks to 18 years; body weights ranged from 3 to 103 kg.

Amprenavir apparent clearance decreased with increasing weight. Weight-adjusted apparent clearance was higher in children younger than 4 years, suggesting that younger children require higher mg-per-kg dosing of LEXIVA.

The pharmacokinetics of LEXIVA oral suspension in protease inhibitor-naive infants younger than 6 months (n = 9) receiving LEXIVA 45 mg per kg plus ritonavir 10 mg per kg twice daily generally demonstrated lower AUC_{12} and C_{min} than adults receiving twice-daily LEXIVA 700 mg plus ritonavir 100 mg, the dose recommended for protease-experienced adults. The mean

Reference ID: 3986026

steady-state amprenavir AUC_{12}, C_{max}, and C_{min} were 26.6 mcg•hour per mL, 6.25 mcg per mL, and 0.86 mcg per mL, respectively. Because of expected low amprenavir exposure and a requirement for large volume of drug, twice-daily dosing of LEXIVA alone (without ritonavir) in pediatric subjects younger than 2 years was not studied.

Pharmacokinetic parameters for LEXIVA administered with food and with ritonavir in this patient population at the recommended weight-band–based dosage regimens are provided in Table 9.

Table 9. Geometric Mean (95% CI) Steady-State Plasma Amprenavir Pharmacokinetic Parameters by Weight in Pediatric and Adolescent Subjects Aged at Least 4 Weeks to 18 Years Receiving LEXIVA with Ritonavir

Weight	Recommended Dosage Regimen	C_{max}		AUC_{24}		C_{min}	
		n	(mcg/mL)	n	(mcg•h/mL)	n	(mcg/mL)
<11 kg	LEXIVA 45 mg/kg plus Ritonavir 7 mg/kg b.i.d.	12	6.00 (3.88, 9.29)	12	57.3 (34.1, 96.2)	27	1.65 (1.22, 2.24)
11 kg - <15 kg	LEXIVA 30 mg/kg plus Ritonavir 3 mg/kg b.i.d.	Not studied[a]					
15 kg - <20 kg	LEXIVA 23 mg/kg plus Ritonavir 3 mg/kg b.i.d.	5	9.54 (4.63, 19.7)	5	121 (54.2, 269)	9	3.56 (2.33, 5.43)
20 kg - <39 kg	LEXIVA 18 mg/kg plus Ritonavir 3 mg/kg b.i.d.	13	6.24 (5.01, 7.77)	12	97.9 (77.0, 124)	23	2.54 (2.11, 3.06)
≥39 kg	LEXIVA 700 mg plus Ritonavir 100 mg b.i.d.	15	5.03 (4.04, 6.26)	15	72.3 (59.6, 87.6)	42	1.98 (1.72, 2.29)

[a] Recommended dose for pediatric patients weighing 11 kg to less than 15 kg is based on population pharmacokinetic analysis.

Subjects aged 2 to younger than 6 years receiving LEXIVA 30 mg per kg twice daily without ritonavir achieved geometric mean (95% CI) amprenavir C_{max} (n = 9), AUC_{12} (n = 9), and C_{min} (n = 19) of 7.15 (5.05, 10.1), 22.3 (15.3, 32.6), and 0.513 (0.384, 0.686), respectively.

Geriatric Patients: The pharmacokinetics of amprenavir after administration of LEXIVA to patients older than 65 years have not been studied *[see Use in Specific Populations (8.5)]*.

Gender: The pharmacokinetics of amprenavir after administration of LEXIVA do not differ between males and females.

Race: The pharmacokinetics of amprenavir after administration of LEXIVA do not differ between blacks and non-blacks.

Drug Interactions

[See Contraindications (4), Warnings and Precautions (5.1), Drug Interactions (7).]

Reference ID: 3986026

Amprenavir, the active metabolite of fosamprenavir, is metabolized in the liver by the cytochrome P450 enzyme system. Amprenavir inhibits CYP3A4. Data also suggest that amprenavir induces CYP3A4. Caution should be used when coadministering medications that are substrates, inhibitors, or inducers of CYP3A4, or potentially toxic medications that are metabolized by CYP3A4. Amprenavir does not inhibit CYP2D6, CYP1A2, CYP2C9, CYP2C19, CYP2E1, or uridine glucuronosyltransferase (UDPGT). Amprenavir is both a substrate for and inducer of P-glycoprotein.

Drug interaction trials were performed with LEXIVA and other drugs likely to be coadministered or drugs commonly used as probes for pharmacokinetic interactions. The effects of coadministration on AUC, C_{max}, and C_{min} values are summarized in Table 10 (effect of other drugs on amprenavir) and Table 12 (effect of LEXIVA on other drugs). In addition, since LEXIVA delivers comparable amprenavir plasma concentrations as AGENERASE, drug interaction data derived from trials with AGENERASE are provided in Tables 11 and 13. For information regarding clinical recommendations, *[see Drug Interactions (7)]*.

Table 10. Drug Interactions: Pharmacokinetic Parameters for Amprenavir after Administration of LEXIVA in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	Dose of LEXIVA[a]	n	% Change in Amprenavir Pharmacokinetic Parameters (90% CI)		
			C_{max}	AUC	C_{min}
Antacid (MAALOX TC®) 30 mL single dose	1,400 mg single dose	30	↓35 (↓24 to ↓42)	↓18 (↓9 to ↓26)	↑14 (↓7 to ↑39)
Atazanavir 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	22	↔	↔	↔
Atorvastatin 10 mg q.d. for 4 days	1,400 mg b.i.d. for 2 weeks	16	↓18 (↓34 to ↑1)	↓27 (↓41 to ↓12)	↓12 (↓27 to ↓6)
Atorvastatin 10 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↔	↔	↔
Efavirenz 600 mg q.d. for 2 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 2 weeks	16	↔	↓13 (↓30 to ↑7)	↓36 (↓8 to ↓56)
Efavirenz 600 mg q.d. plus additional ritonavir 100 mg q.d. for 2 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 2 weeks	16	↑18 (↑1 to ↑38)	↑11 (0 to ↑24)	↔

Reference ID: 3986026

Efavirenz 600 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↔	↔	↓17 (↓4 to ↓29)
Esomeprazole 20 mg q.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	25	↔	↔	↔
Esomeprazole 20 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	23	↔	↔	↔
Ethinyl estradiol/ norethindrone 0.035 mg/0.5 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir[b] 100 mg b.i.d. for 21 days	25	↔[c]	↔[c]	↔[c]
Ketoconazole[d] 200 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 days	15	↔	↔	↔
Lopinavir/ritonavir 533 mg/133 mg b.i.d.	1,400 mg b.i.d. for 2 weeks	18	↓13[e]	↓26[e]	↓42[e]
Lopinavir/ritonavir 400 mg/100 mg b.i.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	18	↓58 (↓42 to ↓70)	↓63 (↓51 to ↓72)	↓65 (↓54 to ↓73)
Maraviroc 300 mg b.i.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 20 days	14	↓34 (↓25 to ↓41)	↓35 (↓29 to ↓41)	↓36 (↓27 to ↓43)
Maraviroc 300 mg q.d. for 10 days	1,400 mg q.d. plus ritonavir 100 mg q.d. for 20 days	14	↓29 (↓20 to ↓38)	↓30 (↓23 to ↓36)	↓15 (↓3 to ↓25)
Methadone 70 to 120 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	19	↔[c]	↔[c]	↔[c]
Nevirapine 200 mg b.i.d. for 2 weeks[f]	1,400 mg b.i.d. for 2 weeks	17	↓25 (↓37 to ↓10)	↓33 (↓45 to ↓20)	↓35 (↓50 to ↓15)

Reference ID: 3986026

Nevirapine 200 mg b.i.d. for 2 weeks[f]	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	17	↔	↓11 (↓23 to ↑3)	↓19 (↓32 to ↓4)
Phenytoin 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	13	↔	↑20 (↑8 to ↑34)	↑19 (↑6 to ↑33)
Raltegravir 400 mg b.i.d. for 14 days	1,400 mg b.i.d. for 14 days (fasted)	14	↓27 (↓46 to ↔)	↓36 (↓53 to ↓13)	↓43[g] (↓59 to ↓21)
	1,400 mg b.i.d. for 14 days[h]	14	↓15 (↓27 to ↓1)	↓17 (↓27 to ↓6)	↓32[g] (↓53 to ↓1)
	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 14 days (fasted)	14	↓14 (↓39 to ↑20)	↓17 (↓38 to ↑12)	↓20[g] (↓45 to ↑17)
	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 14 days[h]	12	↓25 (↓42 to ↓2)	↓25 (↓44 to ↔)	↓33[g] (↓52 to ↓7)
Raltegravir 400 mg b.i.d. for 14 days	1,400 mg q.d. plus ritonavir 100 mg q.d. for 14 days (fasted)	13	↓18 (↓34 to ↔)	↓24 (↓41 to ↔)	↓50[g] (↓64 to ↓31)
	1,400 mg q.d. plus ritonavir 100 mg q.d. for 14 days[h]	14	↑27 (↓1 to ↑62)	↑13 (↓7 to ↑38)	↓17[g] (↓45 to ↑26)
Ranitidine 300 mg single dose (administered 1 hour before fosamprenavir)	1,400 mg single dose	30	↓51 (↓43 to ↓58)	↓30 (↓22 to ↓37)	↔ (↓19 to ↑21)
Rifabutin 150 mg q.o.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	15	↑36[c] (↑18 to ↑55)	↑35[c] (↑17 to ↑56)	↑17[c] (↓1 to ↑39)
Tenofovir 300 mg q.d. for 4 to 48 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 to 48 weeks	45	NA	NA	↔[i]

Reference ID: 3986026

Tenofovir 300 mg q.d. for 4 to 48 weeks	1,400 mg q.d. plus ritonavir 200 mg q.d. for 4 to 48 weeks	60	NA	NA	↔[i]

[a] Concomitant medication is also shown in this column where appropriate.
[b] Ritonavir C_{max}, AUC, and C_{min} increased by 63%, 45%, and 13%, respectively, compared with historical control.
[c] Compared with historical control.
[d] Subjects were receiving LEXIVA/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and LEXIVA/ritonavir.
[e] Compared with LEXIVA 700 mg/ritonavir 100 mg b.i.d. for 2 weeks.
[f] Subjects were receiving nevirapine for at least 12 weeks prior to trial.
[g] C_{last} ($C_{12\,h}$ or $C_{24\,h}$).
[h] Doses of LEXIVA and raltegravir were given with food on pharmacokinetic sampling days and without regard to food all other days.
[i] Compared with parallel control group.
↑= Increase; ↓= Decrease; ↔ = No change (↑or ↓ less than or equal to 10%), NA = Not applicable.

Table 11. Drug Interactions: Pharmacokinetic Parameters for Amprenavir after Administration of AGENERASE in the Presence of the Coadministered Drug(s)

Coadministered Drug(s) and Dose(s)	**Dose of AGENERASE[a]**	**n**	**% Change in Amprenavir Pharmacokinetic Parameters (90% CI)**		
			C_{max}	**AUC**	**C_{min}**
Abacavir 300 mg b.i.d. for 2 to 3 weeks	900 mg b.i.d. for 2 to 3 weeks	4	↔[a]	↔[a]	↔[a]
Clarithromycin 500 mg b.i.d. for 4 days	1,200 mg b.i.d. for 4 days	12	↑15 (↑1 to ↑31)	↑18 (↑8 to ↑29)	↑39 (↑31 to ↑47)
Delavirdine 600 mg b.i.d. for 10 days	600 mg b.i.d. for 10 days	9	↑40[b]	↑130[b]	↑125[b]
Ethinyl estradiol/norethindrone 0.035 mg/1 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↓22 (↓35 to ↓8)	↓20 (↓41 to ↑8)
Indinavir 800 mg t.i.d. for 2 weeks (fasted)	750 or 800 mg t.i.d. for 2 weeks (fasted)	9	↑18 (↑13 to ↑58)	↑33 (↑2 to ↑73)	↑25 (↓27 to ↑116)
Ketoconazole 400 mg single dose	1,200 mg single dose	12	↓16 (↓25 to ↓6)	↑31 (↑20 to ↑42)	NA

Reference ID: 3986026

Lamivudine 150 mg single dose	600 mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg q.d. for >30 days	1,200 mg b.i.d. for 10 days	16	↓27[c]	↓30[c]	↓25[c]
Nelfinavir 750 mg t.i.d. for 2 weeks (fed)	750 or 800 mg t.i.d. for 2 weeks (fed)	6	↓14 (↓38 to ↑20)	↔	↑189 (↑52 to ↑448)
Rifabutin 300 mg q.d. for 10 days	1,200 mg b.i.d. for 10 days	5	↔	↓15 (↓28 to 0)	↓15 (↓38 to ↑17)
Rifampin 300 mg q.d. for 4 days	1,200 mg b.i.d. for 4 days	11	↓70 (↓76 to ↓62)	↓82 (↓84 to ↓78)	↓92 (↓95 to ↓89)
Saquinavir 800 mg t.i.d. for 2 weeks (fed)	750 or 800 mg t.i.d. for 2 weeks (fed)	7	↓37 (↓54 to ↓14)	↓32 (↓49 to ↓9)	↓14 (↓52 to ↑54)
Zidovudine 300 mg single dose	600 mg single dose	12	↔	↑13 (↓2 to ↑31)	NA

[a] Compared with parallel control group.
[b] Median percent change; confidence interval not reported.
[c] Compared with historical data.
↑ = Increase; ↓ = Decrease; ↔ = No change (↑or ↓ less than 10%); NA = C_{min} not calculated for single-dose trial.

Table 12. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the Presence of Amprenavir after Administration of LEXIVA

Coadministered Drug(s) and Dose(s)	Dose of LEXIVA[a]	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C_{max}	AUC	C_{min}
Atazanavir 300 mg q.d. for 10 days[b]	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	21	↓24 (↓39 to ↓6)	↓22 (↓34 to ↓9)	↔
Atorvastatin 10 mg q.d. for 4 days	1,400 mg b.i.d. for 2 weeks	16	↑304 (↑205 to ↑437)	↑130 (↑100 to ↑164)	↓10 (↓27 to ↑12)
Atorvastatin 10 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	16	↑184 (↑126 to ↑257)	↑153 (↑115 to ↑199)	↑73 (↑45 to ↑108)
Esomeprazole 20 mg q.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	25	↔	↑55 (↑39 to ↑73)	ND

Reference ID: 3986026

Esomeprazole 20 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	23	↔	↔	ND
Ethinyl estradiol[c] 0.035 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 21 days	25	↓28 (↓21 to ↓35)	↓37 (↓30 to ↓42)	ND
Dolutegravir 50 mg q.d.	700 mg b.i.d. plus ritonavir 100 mg b.i.d.	12	↓24 (↓8 to ↓37)	↓35 (↓22 to ↓46)	↓49 (↓37 to ↓59)
Ketoconazole[d] 200 mg q.d. for 4 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 4 days	15	↑25 (↑0 to ↑56)	↑169 (↑108 to ↑248)	ND
Lopinavir/ritonavir[e] 533 mg/133 mg b.i.d. for 2 weeks	1,400 mg b.i.d. for 2 weeks	18	↔[f]	↔[f]	↔[f]
Lopinavir/ritonavir[e] 400 mg/100 mg b.i.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	18	↑30 (↓15 to ↑47)	↑37 (↓20 to ↑55)	↑52 (↓28 to ↑82)
Maraviroc 300 mg b.i.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 20 days	14	↑52 (↑27 to ↑82)	↑149 (↑119 to ↑182)	↑374 (↑303 to ↑457)
Maraviroc 300 mg q.d. for 10 days	1,400 mg q.d. plus ritonavir 100 mg q.d. for 20 days	14	↑45 (↑20 to ↑74)	↑126 (↑99 to ↑158)	↑80 (↑53 to ↑113)
Methadone 70 to 120 mg q.d. for 2 weeks	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	19	R-Methadone (active)		
			↓21[g] (↓30 to ↓12)	↓18[g] (↓27 to ↓8)	↓11[g] (↓21 to ↑1)
			S-Methadone (inactive)		
			↓43[g] (↓49 to ↓37)	↓43[g] (↓50 to ↓36)	↓41[g] (↓49 to ↓31)
Nevirapine 200 mg b.i.d. for 2 weeks[h]	1,400 mg b.i.d. for 2 weeks	17	↑25 (↑14 to ↑37)	↑29 (↑19 to ↑40)	↑34 (↑20 to ↑49)

Reference ID: 3986026

Nevirapine 200 mg b.i.d. for 2 weeks[h]	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	17	↑13 (↑3 to ↑24)	↑14 (↑5 to ↑24)	↑22 (↑9 to ↑35)
Norethindrone[c] 0.5 mg q.d. for 21 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 21 days	25	↓38 (↓32 to ↓44)	↓34 (↓30 to ↓37)	↓26 (↓20 to ↓32)
Phenytoin 300 mg q.d. for 10 days	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 10 days	14	↓20 (↓12 to ↓27)	↓22 (↓17 to ↓27)	↓29 (↓23 to ↓34)
Rifabutin 150 mg every other day for 2 weeks[i] (25-O-desacetylrifabutin metabolite) Rifabutin + 25-O-desacetylrifabutin metabolite	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 2 weeks	15	↓14 (↓28 to ↑4) ↑579 (↑479 to ↑698) NA	↔ ↑1,120 (↑965 to ↑1,300) ↑64 (↑46 to ↑84)	↑28 (↑12 to ↑46) ↑2,510 (↑1,910 to ↑3,300) NA
Rosuvastatin 10 mg single dose	700 mg b.i.d. plus ritonavir 100 mg b.i.d. for 7 days		(↑45)	(↑8)	NA

[a] Concomitant medication is also shown in this column where appropriate.
[b] Comparison arm of atazanavir 300 mg q.d. plus ritonavir 100 mg q.d. for 10 days.
[c] Administered as a combination oral contraceptive tablet: ethinyl estradiol 0.035 mg/norethindrone 0.5 mg.
[d] Subjects were receiving LEXIVA/ritonavir for 10 days prior to the 4-day treatment period with both ketoconazole and LEXIVA/ritonavir.
[e] Data represent lopinavir concentrations.
[f] Compared with lopinavir 400 mg/ritonavir 100 mg b.i.d. for 2 weeks.
[g] Dose normalized to methadone 100 mg. The unbound concentration of the active moiety, R-methadone, was unchanged.
[h] Subjects were receiving nevirapine for at least 12 weeks prior to trial.
[i] Comparison arm of rifabutin 300 mg q.d. for 2 weeks. AUC is $AUC_{(0\text{-}48\ h)}$.
↑ = Increase; ↓= Decrease; ↔ = No change (↑or ↓less than 10%); ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

Reference ID: 3986026

Table 13. Drug Interactions: Pharmacokinetic Parameters for Coadministered Drug in the Presence of Amprenavir after Administration of AGENERASE

Coadministered Drug(s) and Dose(s)	Dose of AGENERASE	n	% Change in Pharmacokinetic Parameters of Coadministered Drug (90% CI)		
			C_{max}	AUC	C_{min}
Abacavir 300 mg b.i.d. for 2 to 3 weeks	900 mg b.i.d. for 2 to 3 weeks	4	↔[a]	↔[a]	↔[a]
Clarithromycin 500 mg b.i.d. for 4 days	1,200 mg b.i.d. for 4 days	12	↓10 (↓24 to ↑7)	↔	↔
Delavirdine 600 mg b.i.d. for 10 days	600 mg b.i.d. for 10 days	9	↓47[b]	↓61[b]	↓88[b]
Ethinyl estradiol 0.035 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↔	↑32 (↓3 to ↑79)
Indinavir 800 mg t.i.d. for 2 weeks (fasted)	750 mg or 800 mg t.i.d. for 2 weeks (fasted)	9	↓22[a]	↓38[a]	↓27[a]
Ketoconazole 400 mg single dose	1,200 mg single dose	12	↑19 (↑8 to ↑33)	↑44 (↑31 to ↑59)	NA
Lamivudine 150 mg single dose	600 mg single dose	11	↔	↔	NA
Methadone 44 to 100 mg q.d. for >30 days	1,200 mg b.i.d. for 10 days	16	R-Methadone (active)		
			↓25 (↓32 to ↓18)	↓13 (↓21 to ↓5)	↓21 (↓32 to ↓9)
			S-Methadone (inactive)		
			↓48 (↓55 to ↓40)	↓40 (↓46 to ↓32)	↓53 (↓60 to ↓43)
Nelfinavir 750 mg t.i.d. for 2 weeks (fed)	750 mg or 800 mg t.i.d. for 2 weeks (fed)	6	↑12[a]	↑15[a]	↑14[a]
Norethindrone 1 mg for 1 cycle	1,200 mg b.i.d. for 28 days	10	↔	↑18 (↑1 to ↑38)	↑45 (↑13 to ↑88)
Rifabutin 300 mg q.d. for 10 days	1,200 mg b.i.d. for 10 days	5	↑119 (↑82 to ↑164)	↑193 (↑156 to ↑235)	↑271 (↑171 to ↑409)
Rifampin 300 mg q.d. for 4 days	1,200 mg b.i.d. for 4 days	11	↔	↔	ND
Saquinavir 800 mg t.i.d. for 2 weeks (fed)	750 mg or 800 mg t.i.d. for 2 weeks (fed)	7	↑21[a]	↓19[a]	↓48[a]

Reference ID: 3986026

Zidovudine 300 mg single dose	600 mg single dose	12	↑40 (↑14 to ↑71)	↑31 (↑19 to ↑45)	NA

[a] Compared with historical data.

[b] Median percent change; confidence interval not reported.

↑ = Increase; ↓ = Decrease; ↔= No change (↑or ↓ less than 10%); NA = C_{min} not calculated for single-dose trial; ND = Interaction cannot be determined as C_{min} was below the lower limit of quantitation.

12.4 Microbiology

Mechanism of Action

Fosamprenavir is a prodrug that is rapidly hydrolyzed to amprenavir by cellular phosphatases in the gut epithelium as it is absorbed. Amprenavir is an inhibitor of HIV-1 protease. Amprenavir binds to the active site of HIV-1 protease and thereby prevents the processing of viral Gag and Gag-Pol polyprotein precursors, resulting in the formation of immature non-infectious viral particles.

Antiviral Activity

Fosamprenavir has little or no antiviral activity in cell culture. The antiviral activity of amprenavir was evaluated against HIV-1 IIIB in both acutely and chronically infected lymphoblastic cell lines (MT-4, CEM-CCRF, H9) and in peripheral blood lymphocytes in cell culture. The 50% effective concentration (EC_{50}) of amprenavir ranged from 0.012 to 0.08 microM in acutely infected cells and was 0.41 microM in chronically infected cells (1 microM = 0.50 mcg per mL). The median EC_{50} value of amprenavir against HIV-1 isolates from clades A to G was 0.00095 microM in peripheral blood mononuclear cells (PBMCs). Similarly, the EC_{50} values for amprenavir against monocytes/macrophage tropic HIV-1 isolates (clade B) ranged from 0.003 to 0.075 microM in monocyte/macrophage cultures. The EC_{50} values of amprenavir against HIV-2 isolates grown in PBMCs were higher than those for HIV-1 isolates, and ranged from 0.003 to 0.11 microM. Amprenavir exhibited synergistic anti–HIV–1 activity in combination with the nucleoside reverse transcriptase inhibitors (NRTIs) abacavir, didanosine, lamivudine, stavudine, tenofovir, and zidovudine; the non-nucleoside reverse transcriptase inhibitors (NNRTIs) delavirdine and efavirenz; and the protease inhibitors atazanavir and saquinavir. Amprenavir exhibited additive anti–HIV–1 activity in combination with the NNRTI nevirapine, the protease inhibitors indinavir, lopinavir, nelfinavir, and ritonavir; and the fusion inhibitor enfuvirtide. These drug combinations have not been adequately studied in humans.

Resistance

HIV-1 isolates with decreased susceptibility to amprenavir have been selected in cell culture and obtained from subjects treated with fosamprenavir. Genotypic analysis of isolates from treatment-naive subjects failing amprenavir-containing regimens showed substitutions in the

Reference ID: 3986026

HIV-1 protease gene resulting in amino acid substitutions primarily at positions V32I, M46I/L, I47V, I50V, I54L/M, and I84V, as well as substitutions in the p7/p1 and p1/p6 Gag and Gag-Pol polyprotein precursor cleavage sites. Some of these amprenavir resistance-associated substitutions have also been detected in HIV-1 isolates from antiretroviral-naive subjects treated with LEXIVA. Of the 488 antiretroviral-naive subjects treated with LEXIVA 1,400 mg twice daily or LEXIVA 1,400 mg plus ritonavir 200 mg once daily in Trials APV30001 and APV30002, respectively, 61 subjects (29 receiving LEXIVA and 32 receiving LEXIVA/ritonavir) with virologic failure (plasma HIV-1 RNA greater than 1,000 copies per mL on 2 occasions on or after Week 12) were genotyped. Five of the 29 antiretroviral-naive subjects (17%) receiving LEXIVA without ritonavir in Trial APV30001 had evidence of genotypic resistance to amprenavir: I54L/M (n = 2), I54L + L33F (n = 1), V32I + I47V (n = 1), and M46I + I47V (n = 1). No amprenavir resistance-associated substitutions were detected in antiretroviral-naive subjects treated with LEXIVA/ritonavir for 48 weeks in Trial APV30002. However, the M46I and I50V substitutions were detected in isolates from 1 virologic failure subject receiving LEXIVA/ritonavir once daily at Week 160 (HIV-1 RNA greater than 500 copies per mL). Upon retrospective analysis of stored samples using an ultrasensitive assay, these resistant substitutions were traced back to Week 84 (76 weeks prior to clinical virologic failure).

Cross-Resistance

Varying degrees of cross-resistance among HIV-1 protease inhibitors have been observed. An association between virologic response at 48 weeks (HIV-1 RNA level less than 400 copies per mL) and protease inhibitor-resistance substitutions detected in baseline HIV-1 isolates from protease inhibitor-experienced subjects receiving LEXIVA/ritonavir twice daily (n = 88), or lopinavir/ritonavir twice daily (n = 85) in Trial APV30003 is shown in Table 14. The majority of subjects had previously received either one (47%) or 2 protease inhibitors (36%), most commonly nelfinavir (57%) and indinavir (53%). Out of 102 subjects with baseline phenotypes receiving twice-daily LEXIVA/ritonavir, 54% (n = 55) had resistance to at least one protease inhibitor, with 98% (n = 54) of those having resistance to nelfinavir. Out of 97 subjects with baseline phenotypes in the lopinavir/ritonavir arm, 60% (n = 58) had resistance to at least one protease inhibitor, with 97% (n = 56) of those having resistance to nelfinavir.

Table 14. Responders at Trial Week 48 by Presence of Baseline Protease Inhibitor Resistance-Associated Substitutions[a]

Protease Inhibitor Resistance-Associated Substitutions[b]	LEXIVA/Ritonavir b.i.d. (n = 88)		Lopinavir/Ritonavir b.i.d. (n = 85)	
D30N	21/22	95%	17/19	89%
N88D/S	20/22	91%	12/12	100%

Reference ID: 3986026

L90M	16/31	52%	17/29	59%
M46I/L	11/22	50%	12/24	50%
V82A/F/T/S	2/9	22%	6/17	35%
I54V	2/11	18%	6/11	55%
I84V	1/6	17%	2/5	40%

[a] Results should be interpreted with caution because the subgroups were small.
[b] Most subjects had greater than 1 protease inhibitor resistance-associated substitution at baseline.

The virologic response based upon baseline phenotype was assessed. Baseline isolates from protease inhibitor-experienced subjects responding to LEXIVA/ritonavir twice daily had a median shift in susceptibility to amprenavir relative to a standard wild-type reference strain of 0.7 (range: 0.1 to 5.4, n = 62), and baseline isolates from individuals failing therapy had a median shift in susceptibility of 1.9 (range: 0.2 to 14, n = 29). Because this was a select patient population, these data do not constitute definitive clinical susceptibility break points. Additional data are needed to determine clinically relevant break points for LEXIVA.

Isolates from 15 of the 20 subjects receiving twice-daily LEXIVA/ritonavir up to Week 48 and experiencing virologic failure/ongoing replication were subjected to genotypic analysis. The following amprenavir resistance-associated substitutions were found either alone or in combination: V32I, M46I/L, I47V, I50V, I54L/M, and I84V. Isolates from 4 of the 16 subjects continuing to receive twice-daily LEXIVA/ritonavir up to Week 96 who experienced virologic failure underwent genotypic analysis. Isolates from 2 subjects contained amprenavir resistance-associated substitutions: V32I, M46I, and I47V in 1 isolate and I84V in the other.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In long-term carcinogenicity studies, fosamprenavir was administered orally for up to 104 weeks at doses of 250, 400, or 600 mg per kg per day in mice and at doses of 300, 825, or 2,250 mg per kg per day in rats. Exposures at these doses were 0.3- to 0.7-fold (mice) and 0.7- to 1.4-fold (rats) those in humans given 1,400 mg twice daily of fosamprenavir alone, and 0.2- to 0.3-fold (mice) and 0.3- to 0.7-fold (rats) those in humans given 1,400 mg once daily of fosamprenavir plus 200 mg ritonavir once daily. Exposures in the carcinogenicity studies were 0.1- to 0.3-fold (mice) and 0.3- to 0.6-fold (rats) those in humans given 700 mg of fosamprenavir plus 100 mg ritonavir twice daily. There was an increase in hepatocellular adenomas and hepatocellular carcinomas at all doses in male mice and at 600 mg per kg per day in female mice, and in hepatocellular adenomas and thyroid follicular cell adenomas at all doses in male rats, and at 835 mg per kg per day and 2,250 mg per kg per day in female rats. The relevance of the hepatocellular findings in the rodents for humans is uncertain. Repeat-dose studies with

Reference ID: 3986026

fosamprenavir in rats produced effects consistent with enzyme induction, which predisposes rats, but not humans, to thyroid neoplasms. In addition, in rats only there was an increase in interstitial cell hyperplasia at 825 mg per kg per day and 2,250 mg per kg per day, and an increase in uterine endometrial adenocarcinoma at 2,250 mg per kg per day. The incidence of endometrial findings was slightly increased over concurrent controls, but was within background range for female rats. The relevance of the uterine endometrial adenocarcinoma findings in rats for humans is uncertain.

Fosamprenavir was not mutagenic or genotoxic in a battery of in vitro and in vivo assays. These assays included bacterial reverse mutation (Ames), mouse lymphoma, rat micronucleus, and chromosome aberrations in human lymphocytes.

The effects of fosamprenavir on fertility and general reproductive performance were investigated in male (treated for 4 weeks before mating) and female rats (treated for 2 weeks before mating through postpartum day 6). Systemic exposures ($AUC_{0\text{-}24\,h}$) to amprenavir in these studies were 3 (males) to 4 (females) times higher than exposures in humans following administration of the MRHD of fosamprenavir alone or similar to those seen in humans following administration of fosamprenavir in combination with ritonavir. Fosamprenavir did not impair mating or fertility of male or female rats and did not affect the development and maturation of sperm from treated rats.

14 CLINICAL STUDIES

14.1 Therapy-Naive Adult Trials

APV30001

A randomized, open-label trial evaluated treatment with LEXIVA tablets (1,400 mg twice daily) versus nelfinavir (1,250 mg twice daily) in 249 antiretroviral treatment-naive subjects. Both groups of subjects also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the subjects in this trial was 37 years (range: 17 to 70 years); 69% of the subjects were male, 20% were CDC Class C (AIDS), 24% were white, 32% were black, and 44% were Hispanic. At baseline, the median CD4+ cell count was 212 cells per mm^3 (range: 2 to 1,136 cells per mm^3; 18% of subjects had a CD4+ cell count of less than 50 cells per mm^3 and 30% were in the range of 50 to less than 200 cells per mm^3). Baseline median HIV-1 RNA was 4.83 $\log_{10}$ copies per mL (range: 1.69 to 7.41 $\log_{10}$ copies per mL; 45% of subjects had greater than 100,000 copies per mL).

The outcomes of randomized treatment are provided in Table 15.

Reference ID: 3986026

Table 15. Outcomes of Randomized Treatment through Week 48 (APV30001)

Outcome (Rebound or discontinuation = failure)	LEXIVA 1,400 mg b.i.d. (n = 166)	Nelfinavir 1,250 mg b.i.d. (n = 83)
Responder[a]	66% (57%)	52% (42%)
Virologic failure	19%	32%
Rebound	16%	19%
Never suppressed through Week 48	3%	13%
Clinical progression	1%	1%
Death	0%	1%
Discontinued due to adverse reactions	4%	2%
Discontinued due to other reasons[b]	10%	10%

[a] Subjects achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (less than 50 copies per mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).

[b] Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 16.

Table 16. Proportions of Responders through Week 48 by Screening Viral Load (APV30001)

Screening Viral Load HIV-1 RNA (copies/mL)	LEXIVA 1,400 mg b.i.d.		Nelfinavir 1,250 mg b.i.d.	
	<400 copies/mL	n	<400 copies/mL	n
≤100,000	65%	93	65%	46
>100,000	67%	73	36%	37

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 201 cells per mm^3 in the group receiving LEXIVA and 216 cells per mm^3 in the nelfinavir group.

APV30002

A randomized, open-label trial evaluated treatment with LEXIVA tablets (1,400 mg once daily) plus ritonavir (200 mg once daily) versus nelfinavir (1,250 mg twice daily) in 649 treatment-naive subjects. Both treatment groups also received abacavir (300 mg twice daily) and lamivudine (150 mg twice daily).

The mean age of the subjects in this trial was 37 years (range: 18 to 69 years); 73% of the subjects were male, 22% were CDC Class C, 53% were white, 36% were black, and 8% were Hispanic. At baseline, the median CD4+ cell count was 170 cells per mm^3 (range: 1 to 1,055 cells per mm^3; 20% of subjects had a CD4+ cell count of less than 50 cells per mm^3 and

Reference ID: 3986026

35% were in the range of 50 to less than 200 cells per mm^3). Baseline median HIV-1 RNA was 4.81 log_{10} copies per mL (range: 2.65 to 7.29 log_{10} copies per mL; 43% of subjects had greater than 100,000 copies per mL).

The outcomes of randomized treatment are provided in Table 17.

Table 17. Outcomes of Randomized Treatment through Week 48 (APV30002)

Outcome (Rebound or discontinuation = failure)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d. (n = 322)	Nelfinavir 1,250 mg b.i.d. (n = 327)
Responder[a]	69% (58%)	68% (55%)
Virologic failure	6%	16%
Rebound	5%	8%
Never suppressed through Week 48	1%	8%
Death	1%	0%
Discontinued due to adverse reactions	9%	6%
Discontinued due to other reasons[b]	15%	10%

[a] Subjects achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (less than 50 copies per mL) through Week 48 (Roche AMPLICOR HIV-1 MONITOR Assay Version 1.5).

[b] Includes consent withdrawn, lost to follow up, protocol violations, those with missing data, and other reasons.

Treatment response by viral load strata is shown in Table 18.

Table 18. Proportions of Responders through Week 48 by Screening Viral Load (APV30002)

Screening Viral Load HIV-1 RNA (copies/mL)	LEXIVA 1,400 mg q.d./ Ritonavir 200 mg q.d.		Nelfinavir 1,250 mg b.i.d.	
	<400 copies/mL	n	<400 copies/mL	n
≤100,000	72%	197	73%	194
>100,000	66%	125	64%	133

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 203 cells per mm^3 in the group receiving LEXIVA and 207 cells per mm^3 in the nelfinavir group.

14.2 Protease Inhibitor-Experienced Adult Trials

APV30003

A randomized, open-label, multicenter trial evaluated 2 different regimens of LEXIVA plus ritonavir (LEXIVA tablets 700 mg twice daily plus ritonavir 100 mg twice daily or LEXIVA tablets 1,400 mg once daily plus ritonavir 200 mg once daily) versus lopinavir/ritonavir

Reference ID: 3986026

(400 mg/100 mg twice daily) in 315 subjects who had experienced virologic failure to 1 or 2 prior protease inhibitor-containing regimens.

The mean age of the subjects in this trial was 42 years (range: 24 to 72 years); 85% were male, 33% were CDC Class C, 67% were white, 24% were black, and 9% were Hispanic. The median CD4+ cell count at baseline was 263 cells per mm^3 (range: 2 to 1,171 cells per mm^3). Baseline median plasma HIV-1 RNA level was 4.14 log_{10} copies per mL (range: 1.69 to 6.41 log_{10} copies per mL).

The median durations of prior exposure to NRTIs were 257 weeks for subjects receiving LEXIVA/ritonavir twice daily (79% had greater than or equal to 3 prior NRTIs) and 210 weeks for subjects receiving lopinavir/ritonavir (64% had greater than or equal to 3 prior NRTIs). The median durations of prior exposure to protease inhibitors were 149 weeks for subjects receiving LEXIVA/ritonavir twice daily (49% received greater than or equal to 2 prior protease inhibitors) and 130 weeks for subjects receiving lopinavir/ritonavir (40% received greater than or equal to 2 prior protease inhibitors).

The time-averaged changes in plasma HIV-1 RNA from baseline (AAUCMB) at 48 weeks (the endpoint on which the trial was powered) were -1.4 log_{10} copies per mL for twice-daily LEXIVA/ritonavir and -1.67 log_{10} copies per mL for the lopinavir/ritonavir group.

The proportions of subjects who achieved and maintained confirmed HIV-1 RNA less than 400 copies per mL (secondary efficacy endpoint) were 58% with twice-daily LEXIVA/ritonavir and 61% with lopinavir/ritonavir (95% CI for the difference: -16.6, 10.1). The proportions of subjects with HIV-1 RNA less than 50 copies per mL with twice-daily LEXIVA/ritonavir and with lopinavir/ritonavir were 46% and 50%, respectively (95% CI for the difference: -18.3, 8.9). The proportions of subjects who were virologic failures were 29% with twice-daily LEXIVA/ritonavir and 27% with lopinavir/ritonavir.

The frequency of discontinuations due to adverse events and other reasons, and deaths were similar between treatment arms.

Through 48 weeks of therapy, the median increases from baseline in CD4+ cell counts were 81 cells per mm^3 with twice-daily LEXIVA/ritonavir and 91 cells per mm^3 with lopinavir/ritonavir.

This trial was not large enough to reach a definitive conclusion that LEXIVA/ritonavir and lopinavir/ritonavir are clinically equivalent.

Once-daily administration of LEXIVA plus ritonavir is not recommended for protease inhibitor-experienced patients. Through Week 48, 50% and 37% of subjects receiving LEXIVA 1,400 mg plus ritonavir 200 mg once daily had plasma HIV-1 RNA less than 400 copies per mL and less than 50 copies per mL, respectively.

Reference ID: 3986026

14.3 Pediatric Trials

Three open-label trials in pediatric subjects aged at least 4 weeks to 18 years were conducted. In one trial (APV29005), twice-daily dosing regimens (LEXIVA with or without ritonavir) were evaluated in combination with other antiretroviral agents in pediatric subjects aged 2 to 18 years. In a second trial (APV20002), twice-daily dosing regimens (LEXIVA with ritonavir) were evaluated in combination with other antiretroviral agents in pediatric subjects aged at least 4 weeks to younger than 2 years. A third trial (APV20003) evaluated once-daily dosing of LEXIVA with ritonavir; the pharmacokinetic data from this trial did not support a once-daily dosing regimen in any pediatric patient population.

APV29005

LEXIVA: Twenty (18 therapy-naive and 2 therapy-experienced) pediatric subjects received LEXIVA oral suspension without ritonavir twice daily. At Week 24, 65% (13 of 20) achieved HIV-1 RNA less than 400 copies per mL, and the median increase from baseline in CD4+ cell count was 350 cells per mm^3.

LEXIVA plus Ritonavir: Forty-nine protease inhibitor-naive and 40 protease inhibitor-experienced pediatric subjects received LEXIVA oral suspension or tablets with ritonavir twice daily. At Week 24, 71% of protease inhibitor-naive (35 of 49) and 55% of protease inhibitor-experienced (22 of 40) subjects achieved HIV-1 RNA less than 400 copies per mL; median increases from baseline in CD4+ cell counts were 184 cells per mm^3 and 150 cells per mm^3 in protease inhibitor-naive and experienced subjects, respectively.

APV20002

Fifty-four pediatric subjects (49 protease inhibitor-naive and 5 protease inhibitor-experienced) received LEXIVA oral suspension with ritonavir twice daily. At Week 24, 72% of subjects achieved HIV-1 RNA less than 400 copies per mL. The median increases from baseline in CD4+ cell counts were 400 cells per mm^3 in subjects aged at least 4 weeks to younger than 6 months and 278 cells per mm^3 in subjects aged 6 months to 2 years.

16 HOW SUPPLIED/STORAGE AND HANDLING

LEXIVA tablets, 700 mg, are pink, film-coated, capsule-shaped, biconvex tablets, with "GX LL7" debossed on one face.

Bottle of 60 with child-resistant closure (NDC 49702-207-18).

Store at controlled room temperature of 25°C (77°F); excursions permitted to 15° to 30°C (59° to 86°F) (see USP Controlled Room Temperature). Keep container tightly closed.

LEXIVA oral suspension, a white to off-white grape-bubblegum-peppermint–flavored suspension, contains 50 mg of fosamprenavir as fosamprenavir calcium equivalent to approximately 43 mg of amprenavir in each 1 mL.

Reference ID: 3986026

Bottle of 225 mL with child-resistant closure (NDC 49702-208-53).

This product does not require reconstitution.

Store in refrigerator or at room temperature (5° to 30°C; 41° to 86°F). Shake vigorously before using. Do not freeze.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Drug Interactions

A statement to patients and healthcare providers is included on the product's bottle label: ALERT: Find out about medicines that should NOT be taken with LEXIVA.

LEXIVA may interact with many drugs; therefore, advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products, particularly St. John's wort.

Advise patients receiving PDE5 inhibitors that they may be at an increased risk of PDE5 inhibitor-associated adverse events, including hypotension, visual changes, and priapism, and should promptly report any symptoms to their healthcare provider.

Instruct patients receiving hormonal contraceptives to use alternate contraceptive measures during therapy with LEXIVA because hormonal levels may be altered, and if used in combination with LEXIVA and ritonavir, liver enzyme elevations may occur.

Sulfa Allergy

Advise patients to inform their healthcare provider if they have a sulfa allergy. The potential for cross-sensitivity between drugs in the sulfonamide class and fosamprenavir is unknown.

Redistribution/Accumulation of Body Fat

Inform patients that redistribution or accumulation of body fat may occur in patients receiving antiretroviral therapy, including LEXIVA, and that the cause and long-term health effects of these conditions are not known at this time.

Information about HIV-1 Infection

LEXIVA is not a cure for HIV-1 infection and patients may continue to experience illnesses associated with HIV-1 infection, including opportunistic infections. Patients must remain on continuous HIV therapy to control HIV-1 infection and decrease HIV-1-related illness. Patients should be told that sustained decreases in plasma HIV-1 RNA have been associated with a reduced risk of progression to AIDS and death.

Advise patients to remain under the care of a physician when using LEXIVA.

Advise patients to take all HIV medications exactly as prescribed.

Reference ID: 3986026

Advise patients to avoid doing things that can spread HIV-1 infection to others.

Advise patients not to re-use or share needles or other injection equipment.

Advise patients not to share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.

Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with semen, vaginal secretions, or blood.

Female patients should be advised not to breastfeed because it is not known if LEXIVA can be passed to your baby in your breast milk and whether it could harm your baby. Mothers with HIV-1 should not breastfeed because HIV-1 can be passed to the baby in the breast milk.

LEXIVA must always be used in combination with other antiretroviral drugs. Inform patients not to alter the dose or discontinue therapy without consulting their physician. Physicians should instruct their patients that if they miss a dose, they should take it as soon as possible and then return to their normal schedule. Patients should not double their next dose or take more than the prescribed dose.

Oral Suspension

Instruct patients to shake the bottle vigorously before each use and inform them that refrigeration of the oral suspension may improve the taste for some patients.

LEXIVA and AGENERASE are registered trademarks of the ViiV Healthcare group of companies.

The other brands listed are trademarks of their respective owners and are not trademarks of the ViiV Healthcare group of companies. The makers of these brands are not affiliated with and do not endorse the ViiV Healthcare group of companies or its products.

Manufactured for:

ViiV Healthcare
Research Triangle Park, NC 27709

Vertex Pharmaceuticals Incorporated
Cambridge, MA 02139

by:

GlaxoSmithKline
Research Triangle Park, NC 27709

Reference ID: 3986026

©2016 the ViiV Healthcare group of companies. All rights reserved.

LXV:xxPI

PHARMACIST-DETACH HERE AND GIVE INSTRUCTIONS TO PATIENT

PATIENT INFORMATION

LEXIVA® (lex-EE-vah)
(fosamprenavir calcium)
tablets
and
oral suspension

Important: LEXIVA can interact with other medicines and cause serious side effects. It is important to know the medicines that should not be taken with LEXIVA. See the section "Who should not take LEXIVA?"

Read this Patient Information before you start taking LEXIVA and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment.

What is LEXIVA?

LEXIVA is a prescription anti-HIV medicine used with other anti-HIV medicines to treat human immunodeficiency (HIV-1) infections in adults and children 4 weeks of age and older. LEXIVA is a type of anti-HIV medicine called a protease inhibitor. HIV-1 is the virus that causes AIDS (Acquired Immune Deficiency Syndrome).

When used with other anti-HIV medicines, LEXIVA may help:

1. Reduce the amount of HIV-1 in your blood. This is called "viral load".
2. Increase the number of white blood cells called CD4 (T) cells, which help fight off other infections. Reducing the amount of HIV-1 and increasing the CD4 (T) cell count may improve your immune system. This may reduce your risk of death or infections that can happen when your immune system is weak (opportunistic infections).

It is not known if LEXIVA is safe and effective in children younger than 4 weeks of age.

LEXIVA does not cure HIV-1 infection or AIDS. People taking LEXIVA may develop infections or other conditions associated with HIV-1 infection, including opportunistic infections (for example, pneumonia and herpes virus infections).

You should remain under the care of your healthcare provider when using LEXIVA.

Avoid doing things that can spread HIV-1 infection to others.

- Do not re-use or share needles or other injection equipment.
- Do not share personal items that can have blood or body fluids on them, like toothbrushes and razor blades.
- Do not have any kind of sex without protection. Always practice safer sex by using a latex or polyurethane condom to lower the chance of sexual contact with any body fluids such as semen, vaginal secretions, or blood.

Ask your healthcare provider if you have any questions on how to prevent passing HIV to other people.

Who should not take LEXIVA?

Do not take LEXIVA if you take any of the following medicines:

- alfuzosin (UROXATRAL®)
- flecainide
- propafenone (RYTHMOL SR®)
- rifampin (RIFADIN®, RIFAMATE®, RIFATER®, RIMACTANE®)
- ergot including:
  - dihydroergotamine mesylate (D.H.E. 45®, MIGRANAL®)
  - ergotamine tartrate (CAFERGOT®, MIGERGOT®, ERGOMAR®, MEDIHALER ERGOTAMINE®)
  - methylergonovine (METHERGINE®)
- St. John's wort (*Hypericum perforatum*)
- lovastatin (ADVICOR®, ALTOPREV®)
- simvastatin (ZOCOR®, VYTORIN®, SIMCOR®)
- pimozide (ORAP®)
- delavirdine mesylate (RESCRIPTOR®)
- sildenafil (REVATIO®), for treatment of pulmonary arterial hypertension
- triazolam (HALCION®)
- lurasidone (LATUDA®)

Serious problems can happen if you or your child take any of the medicines listed above with LEXIVA.

Do not take LEXIVA if you are allergic to AGENERASE® (amprenavir), fosamprenavir calcium, or any of the ingredients in LEXIVA. See the end of this leaflet for a complete list of ingredients in LEXIVA.

What should I tell my healthcare provider before taking LEXIVA?

Before taking LEXIVA, tell your healthcare provider if you:

- are allergic to medicines that contain sulfa
- have liver problems, including hepatitis B or C
- have kidney problems
- have high blood sugar (diabetes)
- have hemophilia
- have any other medical condition
- are pregnant or plan to become pregnant. It is not known if LEXIVA will harm your unborn baby.

  Pregnancy Registry. There is a pregnancy registry for women who take antiviral medicines during pregnancy. The purpose of the registry is to collect information about the health of you and your baby. Talk to your healthcare provider about how you can take part in this registry.
- **Do not breastfeed.** We do not know if LEXIVA can be passed to your baby in your breast milk and whether it could harm your baby. Also, mothers with HIV-1 should not breastfeed because HIV-1 can be passed to the baby in the breast milk.

Tell your healthcare provider about all prescription and over-the-counter medicines you take. Also tell your healthcare provider about any vitamins, herbal supplements, and dietary supplements you are taking.

Taking LEXIVA with certain other medicines may cause serious side effects. LEXIVA may affect the way other medicines work, and other medicines may affect how LEXIVA works.

Especially tell your healthcare provider if you take:

- quetiapine (SEROQUEL®)
- estrogen-based contraceptives (birth control pills). LEXIVA may reduce effectiveness of estrogen-based contraceptives. During treatment with LEXIVA, you should use a different contraceptive method.
- medicines to treat liver problems, including hepatitis C infection.

Reference ID: 3986026

Know all the medicines that you take. Keep a list of them with you to show healthcare providers and pharmacists when you get a new medicine.

How should I take LEXIVA?

- **Stay under the care of a healthcare provider while taking LEXIVA.**
- Take LEXIVA exactly as prescribed by your healthcare provider.
- Do not change your dose or stop taking LEXIVA without talking with your healthcare provider.
- If your child is taking LEXIVA, your child's healthcare provider will decide the right dose based on your child's weight.
- You can take LEXIVA tablets with or without food.
- **Adults should take LEXIVA oral suspension without food.**
- **Children should take LEXIVA oral suspension with food.** If your child vomits within 30 minutes after taking a dose of LEXIVA, the dose should be repeated.
- Shake LEXIVA oral suspension well before each use.
- If you miss a dose of LEXIVA, take the next dose as soon as possible and then take your next dose at the regular time. Do not double the next dose. If you take too much LEXIVA, call your healthcare provider or go to the nearest hospital emergency room right away.

What are the possible side effects of LEXIVA?

LEXIVA may cause serious side effects including:

- **Severe skin rash.** LEXIVA may cause severe or life-threatening skin reactions or rash.

  If you get a rash with any of the following symptoms, stop taking LEXIVA and call your healthcare provider or get medical help right away:

  - hives or sores in your mouth, or your skin blisters and peels
  - trouble swallowing or breathing
  - swelling of your face, eyes, lips, tongue, or throat

- **Liver problems.** Your healthcare provider should do blood tests before and during your treatment with LEXIVA to check your liver function. Some people with liver problems, including hepatitis B or C, may have an increased risk of developing worsening liver problem during treatment with LEXIVA.

Reference ID: 3986026

- **Diabetes and high blood sugar (hyperglycemia).** Some people who take protease inhibitors, including LEXIVA, can get high blood sugar, develop diabetes, or your diabetes can get worse. Tell your healthcare provider if you notice an increase in thirst or urinate often while taking LEXIVA.
- **Changes in your immune system (Immune Reconstitution Syndrome)** can happen when you start taking HIV medicines. Your immune system may get stronger and begin to fight infections that have been hidden in your body for a long time. Call your healthcare provider right away if you start having new symptoms after starting your HIV medicine.
- **Changes in body fat.** These changes can happen in people who take antiretroviral therapy. The changes may include an increased amount of fat in the upper back and neck ("buffalo hump"), breast, and around the back, chest, and stomach area. Loss of fat from the legs, arms, and face may also happen. The exact cause and long-term health effects of these conditions are not known.
- **Changes in blood tests**. Some people have changes in blood tests while taking LEXIVA. These include increases seen in liver function tests, blood fat levels, and decreases in white blood cells. Your healthcare provider should do regular blood tests before and during your treatment with LEXIVA.
- **Increased bleeding problems in some people with hemophilia.** Some people with hemophilia have increased bleeding with protease inhibitors, including LEXIVA.
- **Kidney stones.** Some people have developed kidney stones while taking LEXIVA. Tell your healthcare provider right away if you develop signs or symptoms of kidney stones:
  - pain in your side
  - blood in your urine
  - pain when you urinate

The most common side effects of LEXIVA in adults include:

- nausea
- vomiting
- diarrhea
- headache

Vomiting is the most common side effect in children when taking LEXIVA.

Reference ID: 3986026

Tell your healthcare provider about any side effect that bothers you or that does not go away.

These are not all the possible side effects of LEXIVA. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store LEXIVA?

- Store LEXIVA tablets at room temperature between 68°F to 77°F (20°C to 25°C).
- Keep the bottle of LEXIVA tablets tightly closed.
- Store LEXIVA oral suspension between 41°F to 86°F (5°C to 30°C). Refrigeration of LEXIVA oral suspension may improve taste for some people.
- Do not freeze.

Keep LEXIVA and all medicines out of the reach of children.

General information about LEXIVA

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use LEXIVA for a condition for which it was not prescribed. Do not give LEXIVA to other people, even if they have the same symptoms you have. It may harm them.

This leaflet summarizes the most important information about LEXIVA. If you would like more information, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about LEXIVA that is written for health professionals.

For more information call 877-844-8872 or go to www.LEXIVA.com.

What are the ingredients in LEXIVA?

Tablets:

Active ingredient: fosamprenavir calcium

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, magnesium stearate, microcrystalline cellulose, and povidone K30. The tablet film-coating contains the inactive ingredients hypromellose, iron oxide red, titanium dioxide, and triacetin.

Oral Suspension:

Active ingredient: fosamprenavir calcium

Inactive ingredients: artificial grape-bubblegum flavor, calcium chloride dihydrate, hypromellose, methylparaben, natural peppermint flavor, polysorbate 80, propylene glycol, propylparaben, purified water, and sucralose.

This Patient Information has been approved by the U.S. Food and Drug Administration.

LEXIVA and AGENERASE are registered trademarks of the ViiV Healthcare group of companies.

The other brands listed are trademarks of their respective owners and are not trademarks of the ViiV Healthcare group of companies. The makers of these brands are not affiliated with and do not endorse the ViiV Healthcare group of companies or its products.

Manufactured for:

ViiV Healthcare
Research Triangle Park, NC 27709

Vertex Pharmaceuticals Incorporated
Cambridge, MA 02139

by:

GlaxoSmithKline
Research Triangle Park, NC 27709

©2016 the ViiV Healthcare group of companies. All rights reserved.

September 2016

LXV:xxPIL