

Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date. Store between 20° to 25°C (68° to 77°F). Questions? 1-866-360-3266. Bayer and Lotrimin Ultra are registered trademarks of Bayer.
© 2021 Bayer. Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981. Product of Japan.

Bayer



Active ingredient	Purpose
Butenafine hydrochloride 1%	Antifungal

Uses ■ cures most athlete's foot between the toes, jock itch and ringworm
■ relieves itching, burning, cracking, and scaling which accompany these conditions

Warnings For external use only

Do not use ■ on nails or scalp ■ in or near the mouth or the eyes ■ for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if too much irritation occurs or irritation gets worse

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube ■ wash the affected skin with soap and water and dry completely before applying ■ for athlete's foot between the toes: apply to affected skin between and around the toes twice a day for 1 week (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventilated shoes. Change shoes and socks at least once daily. ■ for jock itch and ringworm: apply once a day to affected skin for 2 weeks or as directed by a doctor ■ wash hands after each use ■ children under 12 years: ask a doctor



00000000

Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date. Store between 20° to 25°C (68° to 77°F). Questions? 1-866-360-3266. Bayer and Lotrimin Ultra are registered trademarks of Bayer.
© 2021 Bayer. Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981. Product of Japan

Bayer

LOTTRIMIN

ULTRA®

butenafine hydrochloride cream 1% ANTIFUNGAL Net Wt 15g (0.53 oz)

Active ingredient Butenafine hydrochloride 1% **Purpose** Antifungal

Uses ■ cures most athlete's foot between the toes, jock itch and ringworm
■ relieves itching, burning, cracking, and scaling which accompany these conditions

Warnings For external use only

Do not use ■ on nails or scalp ■ in or near the mouth or the eyes ■ for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.


Stop use and ask a doctor if too much irritation occurs or irritation gets worse

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube ■ wash the affected skin with soap and water and dry completely before applying ■ **for athlete's foot between the toes:** apply to affected skin between and around the toes twice a day for 1 week (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventilated shoes. Change shoes and socks at least once daily. ■ **for jock itch and ringworm:** apply once a day to affected skin for 2 weeks or as directed by a doctor ■ wash hands after each use ■ children under 12 years: ask a doctor



00000000



Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date.
Store between 20° to 25°C (68° to 77°F). Questions? 1-866-360-3266
Bayer and Lotrimin Ultra are registered trademarks of Bayer.
© 2021 Bayer Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981 Product of Japan **Bayer**



<i>Active ingredient</i>	<i>Purpose</i>
Butenafine hydrochloride 1%	Antifungal

Uses ■ cures most athlete's foot between the toes, jock itch and ringworm
■ relieves itching, burning, cracking, and scaling which accompany these conditions

Warnings For external use only

Do not use ■ on nails or scalp ■ in or near the mouth or the eyes ■ for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if too much irritation occurs or irritation gets worse

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube
■ wash the affected skin with soap and water and dry completely before applying ■ **for athlete's foot between the toes:** apply to affected skin between and around the toes twice a day for 1 week (morning and night), or once a day for 4 weeks, or as directed by a doctor. Wear well-fitting, ventilated shoes. Change shoes and socks at least once daily. ■ **for jock itch and ringworm:** apply once a day to affected skin for 2 weeks or as directed by a doctor ■ wash hands after each use ■ children under 12 years: ask a doctor

00000000

Do not use if seal on tube is broken or not visible. See crimp for lot number and expiration date. Store between 20° to 25°C (68° to 77°F). Questions? 1-866-360-3286 Bayer and Lotrimin Ultra are registered trademarks of Bayer.
© 2021 Bayer Dist. by: Bayer HealthCare LLC, Whippany, NJ 07981 Product of Japan

Bayer

LOTTRIMIN[®] ULTRA[®]

butenafine hydrochloride cream 1% ANTIFUNGAL Net Wt 12g (0.42 oz)

Active ingredient

Butenafine hydrochloride 1% Antifungal

Purpose

Uses

■ cures most jock itch
■ relieves itching, burning, cracking, and scaling which accompany this condition

Warnings For external use only

Do not use ■ on nails or scalp ■ on or near the mouth or the eyes ■ for vaginal yeast infections

When using this product do not get into the eyes. If eye contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if too much irritation occurs or irritation gets worse

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years and older: ■ use the tip of the cap to break the seal and open the tube ■ wash the affected skin with soap and water and dry completely before applying ■ apply once a day to affected skin for 2 weeks or as directed by a doctor ■ wash hands after each use ■ children under 12 years: ask a doctor



00000000

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
10/18/2021 10:15:12 AM