



PAA166901

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Base Label Page 5

Datos sobre el fármaco (continuación)

Cuadro de dosis		
Peso (libras)	Edad (años)	Dosis (mL)
menos de 24 libras	menos de 2 años	consulte a un médico
24 a 35 libras	2 a 3 años	5 mL
36 a 47 libras	4 a 5 años	7.5 mL
48 a 59 libras	6 a 8 años	10 mL
60 a 71 libras	9 a 10 años	12.5 mL
72 a 95 libras	11 años	15 mL

Información adicional

- cada 5 mL contienen: **3 mg de sodio**
- una dosis dura entre 6 y 8 horas
- consérvelo entre 20 y 25 °C (68 a 77 °F)
- consulte el número de lote y la fecha de vencimiento en la base de la caja

Ingredientes inactivos

saborizante artificial, carboximetilcelulosa sódica, ácido cítrico monohidratado, edetato disódico, FD&C rojo n.º 40, glicerina, celulosa microcristalina, polisorbato 80, agua purificada, benzoato de sodio, solución de sorbitol, sacarosa, goma xantana

¿Preguntas o comentarios?

llame gratuitamente al 1-800-88-ADVIL o consulte a su farmacéutico, médico o profesional de atención de la salud

Base Label Page 5

Insert Back Page 2

Datos sobre el fármaco (continuación)

Advertencias

Alerta de alergia: El ibuprofeno puede provocar una reacción alérgica intensa, especialmente en personas alérgicas a la aspirina. Los síntomas pueden incluir:

- urticaria
- hinchazón facial
- asma (sibilancia)
- choque
- enrojecimiento de la piel
- erupción cutánea
- ampollas.

Si se produce una reacción alérgica, interrumpa su uso y busque ayuda médica de inmediato.

Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede provocar sangrado estomacal intenso. Las probabilidades son mayores si el niño:

- ha tenido problemas de úlceras o sangrado en el estómago
- usa un anticoagulante o un corticosteroide
- usa otros fármacos de venta con o sin receta que contienen AINE [aspirina, ibuprofeno, naproxeno u otros]
- toma este producto en dosis mayores o durante un período más prolongado que lo indicado.

Advertencia de ataque cardíaco y accidente cerebrovascular: Los AINE, exceptuando la aspirina, aumentan el riesgo de ataque cardíaco, infarto y accidentes cerebrovasculares. Estos pueden ser fatales. El riesgo es más alto si usted usa más de lo que se receta, o por un período más largo del que se receta.

Advertencia de dolor de garganta: El dolor de garganta intenso o persistente, o dolor de garganta acompañado de fiebre alta, dolor de cabeza, náuseas y vómitos pueden ser graves. Consulte a un médico sin demora. No use este producto durante más de 2 días ni lo administre a niños menores de 3 años a menos que lo indique un médico.

Insert Back Page 2

Datos sobre el fármaco (continuación)

No lo use

- si el niño ha presentado alguna vez una reacción alérgica a cualquier otro analgésico-antifebril
- inmediatamente antes o después de una cirugía de corazón.

Consulte a un médico antes de usarlo si

- la advertencia de sangrado estomacal es aplicable al niño
- el niño tiene problemas o efectos secundarios graves por tomar analgésicos o antifebriles
- el niño tiene antecedentes de problemas estomacales, como acidez;
- el niño tiene presión arterial alta, enfermedad cardíaca, cirrosis hepática, enfermedad renal, asma, o tuvo un accidente cerebrovascular;
- el niño no ha estado tomando líquidos;
- el niño ha perdido gran cantidad de líquido debido a vómitos o diarrea,
- el niño está tomando un diurético.

Consulte a un médico o farmacéutico antes de usarlo si el niño

- recibe cuidados médicos por cualquier afección grave
- está usando algún otro fármaco.

Al usar este producto

- adminístrelo con alimentos o leche si se produce malestar estomacal

Interrumpa su uso y consulte a un médico si

- el niño presenta alguno de los siguientes signos de sangrado estomacal:
 - siente que va a desmayarse,
 - vomita sangre
 - tiene heces con sangre o de color negro
 - tiene dolor estomacal que no mejora

Insert Back Page 3

Insert Front Page 4

Datos sobre el fármaco (continuación)

- niño(a) tiene síntomas de problemas cardíacos o accidente cerebrovascular:
 - dolor de pecho
 - problemas para respirar
 - debilidad en una parte o lado del cuerpo
 - habla ininteligible o afectada
 - hinchazón de piernas
- el niño no presenta ninguna mejoría dentro del primer día (24 horas) de tratamiento;
- la fiebre o el dolor empeoran o duran más de 3 días
- se presenta enrojecimiento o hinchazón en el área dolorida
- aparece algún síntoma nuevo.

Manténgalo fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o comuníquese con un centro de toxicología de inmediato.

Instrucciones

- este producto no contiene instrucciones ni advertencias completas para su uso en adultos
- no administre cantidades mayores de las indicadas
- agite bien el frasco antes de usarlo
- mL = mililitro
- consulte la dosis correcta en el cuadro. Si es posible, use el peso como referencia para la dosis; como alternativa, use la edad.
- si es necesario, repita la dosis cada **6 a 8 horas**
- no lo use más de **4 veces por día**
- mida únicamente con el vaso dosificador suministrado. El vaso dosificador debe utilizarse exclusivamente con Children's Advil Suspension/Suspensión. No lo use con otros productos. Las líneas que marcan las dosis toman en cuenta el producto que queda adherido al vaso debido a la viscosidad de la suspensión.

Insert Front Page 4

Drug Facts (continued)

Other information

- each 5 mL contains: **sodium 3 mg**
- one dose lasts 6-8 hours
- store at 20-25°C (68-77°F)
- see bottom of box for lot number and expiration date

Inactive ingredients

artificial flavor, carboxymethylcellulose sodium, citric acid monohydrate, edetate disodium, FD&C red no. 40, glycerin, microcrystalline cellulose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, sucrose, xanthan gum

Questions or comments?

call toll-free 1-800-88-ADVIL or ask your pharmacist, doctor or health care professional

Importante: Lea toda la información sobre el producto antes de usarlo. Conserve esta caja para información importante. Este producto está pensado para ser administrado a niños de 2 a 11 años.

Datos sobre el fármaco

Ingrediente activo (por cada 5 mL)	Finalidad
Ibuprofeno 100 mg (AINE)*	Antifebril/analgésico
*antiinflamatorio no esteroideo	

Usos: temporalmente: ■ baja la fiebre

- alivia dolores y malestares leves causados por el resfriado común, la gripe, el dolor de garganta, el dolor de cabeza y el dolor de muelas

For Drug Facts in Spanish, LIFT HERE
Para obtener datos sobre el fármaco en español, LEVANTE AQUÍ

PAA166902

Insert Front Page 1

Insert Front Page 1

PAA166902

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

TOP LABEL

Do not use if plastic wrap is broken or missing. No lo use si el empaquetado plástico con la leyenda "SAFETY SEAL" está roto o falta.

Children's Advil

Suspension/Suspensión
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)
Ibuprofeno suspensión oral
100 mg por cada 5 mL
Analgésico/antifebril (AINE)
Fruit-Flavored Liquid
Líquido con sabor a fruta
4 FL OZ (120 mL)

DO NOT USE if plastic wrap is broken or missing. No lo use si el empaquetado plástico con la leyenda "SAFETY SEAL" está roto o falta.

Active ingredient (per each 5 mL): Ibuprofen 100 mg. It temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache.

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor. Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.
Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic. Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug. When using this product take with food or milk if stomach upset occurs.

Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 3 mg. Store at 20-25° C (68-77° F).

PAA166835

BASE LABEL

Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cap provided. Dosing cap to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cap due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Questions or comments? call 1-800-88-ADVIL. Distributed by: GSK Consumer Healthcare, Warren, NJ 07059. For most recent product information, visit www.Advil.com.

Ingredientes (por cada 5 mL): Ibuprofeno 100 mg.

Úsese bajo temporalmente la fiebre y alivia dolores y malestares leves causados por el resfriado común, la gripe, el dolor de garganta, el dolor de cabeza y el dolor de muelas.

Advertencias
Alerta de alergia: El ibuprofeno puede provocar una reacción alérgica intensa, especialmente en personas alérgicas a la aspirina. Los síntomas pueden incluir: urticaria, hinchazón facial, asma (sibilancia), choque anafiláctico, erupción cutánea y ampollas. Si se produce una reacción alérgica, interrumpa su uso y busque ayuda médica de inmediato.
Advertencia de sangrado estomacal: Este producto contiene un AINE, que puede provocar sangrado estomacal intenso. Las probabilidades son mayores si el niño(a) ha tenido problemas de úlcera o sangrado en el estómago, usa un anticoagulante o un corticoesteroide, usa otros fármacos de venta con o sin receta que contienen AINES (aspirina, ibuprofeno, naproxeno u otros), tome este producto en dosis mayores o durante un período más prolongado que lo indicado.
Advertencia de Ataque Cardíaco y Accidente Cerebrovascular: Los AINES, exceptando la aspirina, aumentan el riesgo de ataque cardíaco, infarto y accidentes cerebrovasculares. Estos pueden ser fatales. El riesgo es más alto si usted usa más de lo que se receta, o por un período más largo del que se receta.
Advertencia de dolor de garganta: El dolor de garganta intenso o persistente, o dolor de garganta acompañado de fiebre alta, dolor de cabeza, náuseas y vómitos pueden ser graves. Consulte a un médico sin demora. No use este producto durante más de 2 días si lo administra a niños menores de 3 años a menos que lo indique un médico.
No lo use si el niño ha presentado alguna vez una reacción alérgica a cualquier otro analgésico/antifebril, inmediatamente antes o después de una cirugía de corazón. Consulte a un médico antes de usarlo si la advertencia de sangrado estomacal es aplicable al niño; el niño tiene antecedentes de problemas estomacales, como úlcera; el niño tiene presión arterial alta, enfermedad cardíaca, cirrosis hepática, enfermedad renal, asma, o tuvo un accidente cerebrovascular; el niño no ha estado tomando líquidos; el niño ha perdido gran cantidad de líquido debido a vómitos o diarrea, o el niño está tomando un diurético. Consulte a un médico e farmacéutico antes de usarlo si el niño recibe cuidados médicos por cualquier afección grave; si está usando algún otro fármaco.

INSIDE TOP LABEL

Al usar este producto, adminístrelo con alimentos o leche si se produce malestar estomacal. Interrumpa su uso y consulte a un médico si el niño presenta alguno de los siguientes signos de sangrado estomacal: vómito que sea a demasayase, yomita sangre, heces hechas con sangre o de color negro, o tiene dolor estomacal que no mejora; niño(a) tiene síntomas de problemas cardíacos o accidente cerebrovascular: dolor de pecho, problemas para respirar, debilidad en una parte o lado del cuerpo, habla ininteligible o alterada, hinchazón de piernas; el niño no presenta ninguna mejoría dentro del primer día (24 horas) de tratamiento; la fiebre o el dolor empeoran o duran más de 3 días; se presenta enrojecimiento o hinchazón en el área dolorida, o aparece algún síntoma nuevo. Manténgalo fuera del alcance de los niños. En caso de sobredosis, obtenga ayuda médica o comuníquese con un centro de toxicología de inmediato. Cada 5 mL contienen 3 mg de sodio. Conservarlo entre 20 y 25 °C (68 a 77 °F). Instrucciones: Este producto no contiene instrucciones ni advertencias completas para su uso en adultos. No administre cantidades mayores de las indicadas.

Agite bien el frasco antes de usarlo. mL = mililitro. Consulte la dosis correcta en el cuadro. Si es posible, use el peso como referencia para la dosis; como alternativa, use la edad. Si es necesario, repita la dosis cada 6 a 8 horas. No lo use más de 4 veces por día. Mida únicamente con el vaso dosificador suministrado. El vaso dosificador debe utilizarse exclusivamente con Children's Advil Suspension. No lo use con otros productos. Las líneas que marcan las dosis toman en cuenta el producto que queda adherido al vaso debido a la viscosidad de la suspensión.

¿Preguntas o comentarios? Llame al 1-800-88-ADVIL. Distribuido por: GSK Consumer Healthcare, Warren, NJ 07059. Para ver la información más reciente sobre el producto, visite www.Advil.com.

Cuadro de dosis		
Peso (libras)	Edad (años)	Dosis (mL)
menos de 24 libras	menos de 2 años	consulte a un médico
24 a 35 libras	2 a 3 años	5 mL
36 a 47 libras	4 a 5 años	7.5 mL
48 a 59 libras	6 a 8 años	10 mL
60 a 71 libras	9 a 10 años	12.5 mL
72 a 95 libras	11 años	15 mL

PAA166835

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.75 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166837

NON MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's
Advil
Suspension

Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Grape-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL) Ibuprofen 100 mg

Use temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Some throat warnings: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor. Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever, right before or after heart surgery.

Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn, child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic. Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.

When using this product take with food or milk to upset stomach occurs.

Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 2 mg. Store at 20-25°C (68-77°F).

Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)
under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

LOT: PAA166830
EXP:

© Questions or comments? call 1-800-88-ADVIL.
Dist. by: GSK, CN, Warren, NJ 07059-62021 GSK or licensee.
For most recent product information, visit www.AAI.com

UNVARNISHED UNPRINTED AREA
FOR LOT AND EXP. DATE.
(PRE-PRINTED LOT INFORMATION AS EXAMPLE,
SHOULD MATCH MARKET REQUIREMENTS)

PAA166830

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.75 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166841

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

White Grape-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg
Use temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches
Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness/itch, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery
Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition, taking any other drug.

When using this product take with food or milk. If stomach upset occurs
Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomit to blood, has bloody or black stools, or has stomach pain that does not get better; **child has symptoms of heart problems or stroke:** chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; **fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear**
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 3 mg. Store at 20-25° C (68-77° F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using, mL = mL for. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

Overdose or comments? call 1-800-86-ADWIL
Date: GSK CH Warner, NJ 07059 ©2011 GSK or licensee
 For most recent product information, visit www.Achil.com

LOT: PAA166834
EXP:

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166834

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.57 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166839

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Bubble Gum-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg
 Use temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches.

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach Bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
 Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.
Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug.

When using this product take with food or milk if stomach upset occurs.
 Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: pale faint, vomit blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
 Each 5 mL contains sodium 3 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using, mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

LOT: PAA166832
EXP:

*For questions or comments call 1-800-98-AD-IVIL.
 Dist. by: GSK CH, Warren, NJ 07059 ©2011 GSK or its licensor.
 For most recent product information, visit www.Advil.com

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166832

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.6 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



DATA MATRIX CODE MUST HAVE 12 CELLS MINIMUM (SIZE AT LEAST 0.25 MM EACH) WITH A WHITE QUIET ZONE OF MINIMUM 1 MM

CODING AREA TO BE PRINTED IN A DARK COLOR AND UNVARNISHED

PAA166840

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Sugar-Free Dye-Free Berry-Flavored Liquid
4 FL OZ (120 mL)

Active ingredient (in each 5 mL) ibuprofen 100 mg

Use temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches.

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include hives, facial swelling, asthma (wheezing), shock, skin redness, rash, blisters. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.
Do not use if the child has ever had an allergic reaction to any other pain reliever/fever reducer, right before or after heart surgery.
Ask a doctor before use if stomach bleeding warning applies to the child, child has a history of stomach problems, such as heartburn, child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition, taking any other drug.

When using this product take with food or milk if stomach upset occurs.
Stop use and ask a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling, the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains sodium 10 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using. mL = mL for. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	Under 2 yr	Ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-95 lb	11 yr	15 mL

LOT: PAA166833
 EXP: [REDACTED]

*Questions or comments? call 1-800-38-40MIL.
 Dist. by: SSK CH, Warren, NJ 07069 ©2021 SSK or licensee
 For most recent product information, visit www.Advil.com.*

UNVARNISHED UNPRINTED AREA FOR LOT AND EXP. DATE. (PRE-PRINTED LOT INFORMATION AS EXAMPLE, SHOULD MATCH MARKET REQUIREMENTS)

PAA166833

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.57 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt



PAA166838

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

Do Not Use if plastic wrap imprinted with "SAFETY SEAL" is broken or missing.

Children's Advil Suspension
Ibuprofen Oral Suspension
100 mg per 5 mL
Pain Reliever/Fever Reducer (NSAID)

Dye-Free / Alcohol-Free
 Blue Raspberry-Flavored Liquid
 4 FL OZ (120 mL)

Active ingredient (in each 5 mL): Ibuprofen 100 mg
Use temporarily reduces fever, relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

Warnings:
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin redness, rash, itches. If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if the child has had stomach ulcers or bleeding problems, takes a blood thinning (anticoagulant) or steroid drug, takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others), takes more or for a longer time than directed.
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult a doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.
Do not use if the child has ever had an allergic reaction to any other fever reducer/pain reliever, right before or after heart surgery
Ask a doctor before use if stomach bleeding warning applies to the child; child has a history of stomach problems, such as heartburn; child has high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke; child has not been drinking fluids; child has lost a lot of fluid due to vomiting or diarrhea; child is taking a diuretic.
Ask a doctor or pharmacist before use if the child is under a doctor's care for any serious condition; taking any other drug

When using this product take with food or milk if upset stomach occurs
Stop use and seek a doctor if child experiences any of the following signs of stomach bleeding: feels faint, vomits blood, has bloody or black stools, or has stomach pain that does not get better; child has symptoms of heart problems or stroke: chest pain, trouble breathing, weakness in one part or side of body, slurred speech, leg swelling; the child does not get any relief within first day (24 hours) of treatment; fever or pain gets worse or lasts more than 3 days; redness or swelling is present in the painful area; any new symptoms appear
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Each 5 mL contains: sodium 10 mg. Store at 20-25°C (68-77°F).
Directions: This product does not contain directions or complete warnings for adult use. Do not give more than directed. Shake well before using, mL = milliliter. Find right dose on chart. If possible, use weight to dose; otherwise use age. Repeat dose every 6-8 hours, if needed. Do not use more than 4 times a day. Measure only with the dosing cup provided. Dosing cup to be used with Children's Advil Suspension only. Do not use with other products. Dose lines account for product remaining in cup due to thickness of suspension.

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)
Under 24 lb	under 2 yr	ask a doctor
24-35 lb	2-3 yr	5 mL
36-47 lb	4-5 yr	7.5 mL
48-59 lb	6-8 yr	10 mL
60-71 lb	9-10 yr	12.5 mL
72-85 lb	11 yr	15 mL

Questions or comments? call 1-800-88-ADVIL.
 Dist. by: CSM, Inc., Warren, NJ 07059-6202 © 2011 CSM or licensee
 For most recent product information, visit www.Adi.com

PAA166831

LOT:
EXP:

UNVARNISHED UNPRINTED AREA
 FOR LOT AND EXP. DATE.
 (PRE-PRINTED LOT INFORMATION AS EXAMPLE,
 SHOULD MATCH MARKET REQUIREMENTS)

PAA166831

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	4.5 pt
• SUBHEADINGS/BODY TEXT	4.5 pt
• LEADING	4.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	.5 pt

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

NUSHIN F TODD
09/20/2022 09:12:00 AM