

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older a have had stomach ulcers or bleeding problems attake a blood thinhing attake about and stomach thinhing attake about or nonprescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) a have 3 or more adoubted crinks every day while using this product. It is the more of for a longer time than directed. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children, in case of overfose, get medical help or contact a Poison Control Directions.

Directions

Adults and children it gears and over: take 4 to 8 capsules every 4 hours while symptoms persist. Do not exceed 48 capsules in 24 hours unless directed by a doctor.

Questions or comments? Call 1-888-842-3567 9:00 AM - 7:00 PM ET, Monday - Friday



Distributed by: PLx Pharma Inc. 9 Fishers Lane, Sparta, NJ 07871 Visit us at: www.vazalore.com

092622

Drug Facts Label Size **Drug Facts Title** NA NA **Drug Facts Continued** Headings 8 pt Subheadings 6 pt **Body Text** 6 pt Leading 7 pt Characters Per Inch <39 **Bullets** 5 pt Space Before Bullet 1 em NA **Barlines** Hairlines NA

| Drug Facts Label | Size |
|----------------------|---------|
| Drug Facts Title | 12.8 pt |
| Drug Facts Continued | 9.3 pt |
| Headings | 9.3 pt |
| Subheadings | 7 pt |
| Body Text | 7 pt |
| Leading | 7.6 pt |
| Characters Per Inch | <39 |
| Bullets | 5.8 pt |
| Space Before Bullet | 2 ems |
| Barlines | 1.5 pt |
| Hairlines | .33 pt |

BB89001 Patents www.xazalore.com/patents MICO. 97016 SEV.WW W 16 2U TIZI V Distrib uted by: PLx Pharma Inc. 9 Fishers Lane, Sparta Au 10 X871 Exp. Date 065 622 Lot Code PLX. CONTROL & POISON CONTROL CENTER TIGHT AWAY. OB BROKEN. Keep out of reach of children. In case of overdose, get medical help or IS MISSING coplems in the unborn child or complications during delivery. nuless definitely directed to do so by a doctor because it may cause PROTECTION" it pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy FOR YOUR ■ sult new symptoms occur. These could be signs of a serious condition. ■ ringing in the ears or loss of hearing occurs ■ redness or swelling is present in the painful area **GEPEATED** ■ fever gets worse or lasts for more than 3 days SEAL WITH ■ pain gets worse or lasts more than 10 days SAFETY ■ have stomach pain that does not get better **MPRINTED** ■ feel faint ■ vormit blood ■ have ebloody or black atools ■ vou experience any of the following signs of spomach bleeding: NZE IŁ Stop use and ask a doctor if ■ anallergic reaction occurs. Seek medical help right away. DONOT EVIDENT: ■ taking a prescription drug for diabetes, gout or arthritis Ask a doctor or pharmacist before useifyou are ■ you aretaking adjuretic ■ you have asthma qızesse ■ you have a history of stomach problems, such as heartburn stomach bleeding warning applies to you Ask a doctor before use if ann reliever/Tever reducer. **po not n**ae ityou have ever had an allergic reaction to aspirin or any other ■ bake more or for a longer time than directed (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product Remove Bottle. Dotted Line to ■ fake other drugs confaining prescription or nonprescription NSAIDs gnolA tuO ■ ÆKE a blood thinning (anticoagulant) or særold drug a sreage 60 or older 👚 have had stomach ulœrs or bleeding problems cause severe stormach diseding. The chance is higher if you: Stomach bleeding warming: This product contains an N SAID, which may Lhis product contains soy. Call 1-888-842-3567 9:00 AM - 7:00 PM ET, Monday - Friday ■ lives ■ facial swelling ■ shock ■assthma (wheezing) Questions or comments? All ergy alert : Aspirin may cause a severe allergic reaction which may include DUSTUUSCEN DOSI IUK' DOES SINUD CUIOLIGE' SOA DESU OII' ELESUINUD GIOXIGE yndrome, a rare but serious illness. illcon dioxide, FD&C Blue #1, hypromellose, lecithin, oleic acid, a doctor because these symptoms could be an early sign of Reye's this product, if changes in behavior with nausea and vorniting occur, con Inactive ingredients anhydrous citric acid, carrageeran, colloidal cylicken box or fin-likes ymptoms should not use this product. When using do not use if the blue band around the capsule is missing or broken ■ avoid excessive heat above 40°C (104°F) Reye's syndrome: Children and teenagers who have or are recovering from ■ store at 15-30°C (59-86°F) ■ save this card for full drections and warnings ■ ask your doctor about other uses for aspirin capsules noitsmrotni rədt0 ■ temporarily reduces fever ■ b.eu.suzunaj and menatruaj cramps ethrifts to nisq nonim 🔳 children under 12 years: consult a doctor Uses ■ for the temporary relief of minor aches and pains associated with:

■ headache ■ backache ■ muscular aches ■ acold ■ toditache
■ headache = muscular aches ■ acold ■ toditache ■ do not exceed 48 capsules in 24 hours unless drected by a doctor ■ take 4 to 8 capsules every 4 hours while symptoms perslet ■ drink a full glass of water with each dose Directions səsodınd Active ingredient (in each capsule) Drug Facts (continued) Drug Facts



| This is a representation of an electronic record that was signed |
|--|
| electronically. Following this are manifestations of any and all |
| electronic signatures for this electronic record. |

/s/

NUSHIN F TODD 10/03/2022 01:38:29 PM