


000079729

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	N/A



000079731

DRUG FACTS TEXT DEFINED	TYPE SIZE
• HEADINGS	N/A
• SUBHEADINGS/BODY TEXT	7 pt
• LEADING	7.5 pt
• BULLETS	N/A
• SPACE BEFORE BULLET	N/A
• WARNING BOX LINE, HAIRLINES	N/A



000079728

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8.0 pt
• DRUG FACTS CONTINUED	7.0 pt
• HEADINGS	7.0 pt
• SUBHEADINGS/BODY TEXT	6.0 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5.0 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



000079730

DRUG FACTS MODIFIED TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces



Stop use and ask a doctor if:
 

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or pink pain
- if pregnant or breast-feeding, ask a health professional before use.

 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect.

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (if needed)**

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use.

Heartburn in children may sometimes be caused by a serious condition.

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

Overdose or comments? call 1-free weekdays 9 AM to 5 PM EST at 1-866-226-1900

Active ingredient (in each capsule)	Purpose
Esomeprazole 20 mg	Acid reducer

(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

**Use**

- treats frequent heartburn (occurs 2 or more days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect.

**Warnings**

**Allergy alert:**

- Do not use if you are allergic to esomeprazole.
- Esomeprazole may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If an allergic reaction occurs, stop use and seek medical help right away.

Do not use if you have:
 

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness
- frequent chest pain. These may be signs of a serious condition. See your doctor.

Ask a doctor before use if you have:
 

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Ask a doctor or pharmacist before use if you are:
 

- taking a prescription drug. Acid reducers may interact with certain prescription drugs.



Stop use and ask a doctor if:

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:** Adults 18 years of age and older: this product is to be used once a day (every 24 hours), every day for 14 days. You may take 1 to 4 days for full effect.

**14-Day Course of Treatment:** swallow 1 capsule with a glass of water before eating in the morning. Take every day for 14 days. Do not take more than 1 capsule a day. Swallow whole. Do not crush or chew capsules. Do not use for more than 14 days unless directed by your doctor.

**Repeated 14-Day Courses (if needed):** you may repeat a 14-day course every 4 months. Do not take for more than 14 days or more often than every 4 months unless directed by a doctor. Children under 18 years of age: ask a doctor before use.

Heartburn in children may sometimes be caused by a serious condition.

**Other information:** read the directions and warnings before use. Keep the carton. It contains important information.

Store at 20-25°C (68-77°F).

Questions or comments? call toll-free weekdays 9 AM to 5 PM EST at 1-800-226-1800

Active ingredient (in each capsule)	Purpose
Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer
<b>Uses:</b> treats frequent heartburn (occurs 2 or more days a week)	not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect
<b>Warnings:</b>	
<b>Allergy alert:</b> Do not use if you are allergic to esomeprazole. Esomeprazole may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If an allergic reaction occurs, stop use and seek medical help right away.	
Do not use if you have: trouble or pain swallowing food, vomiting with blood, or bloody or black stools; heartburn with lightheadedness, swelling or dizziness; chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness or frequent chest pain. These may be signs of a serious condition. See your doctor.	
Ask a doctor before use if you have: had heartburn over 3 months. This may be a sign of a more serious condition.	
frequent wheezing, particularly with heartburn; unexplained weight loss; nausea or vomiting; stomach pain.	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.	



Stop use and ask a doctor if:
 

- your heartburn continues or worsens
- you need to take this product for more than 14 days
- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain
- if pregnant or breast-feeding, ask a health professional before use.

 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.  
**Directions** ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ may take 1 to 4 days for full effect  
**14-Day Course of Treatment** ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days  
 ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor  
**Repeated 14-Day Courses (if needed)** ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use.  
 Heartburn in children may sometimes be caused by a serious condition.  
**Other Information** ■ read the directions and warnings before use ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F)  
**Overdose or comments?** call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1800

**Active ingredient (in each capsule)** ..... Purpose  
 Esomeprazole 20 mg ..... Acid reducer  
 (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

**Uses** ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:** ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash. If an allergic reaction occurs, stop use and seek medical help right away.  
 Do not use if you have: ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain. These may be signs of a serious condition. See your doctor.  
 Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain  
 Ask a doctor or pharmacist before use if you are ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.



KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

**Drug Facts**

**Active ingredient (in each capsule)** Purpose  
Esomeprazole 20 mg Acid reducer  
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

**Uses**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:**  
■ Do not use if you are allergic to esomeprazole.  
■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have:**  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are:**  
■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if:**  
■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days  
■ you need to take more than 1 course of treatment every 4 months

**Drug Facts (continued)**

■ you get diarrhea ■ you develop a rash or joint pain  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ may take 1 to 4 days for full effect

**14-Day Course of Treatment**  
■ swallow 1 capsule with a glass of water before eating in the morning  
■ take every day for 14 days  
■ do not take more than 1 capsule a day  
■ swallow whole. Do not crush or chew capsules.  
■ do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor  
■ children under 18 years of age: ask a doctor before use.  
Heartburn in children may sometimes be caused by a serious condition.

**Other Information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F)

**Inactive ingredients**  
cvm starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

**Questions or comments?** call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.



Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK group of companies or its licensor  
Made in France  
For most recent product information, visit [www.Nexium24HR.com](http://www.Nexium24HR.com)  
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

**Nexium<sup>®</sup> 24HR**  
Clinically Proven to Treat Frequent Heartburn  
Capsules

See new warning information NDC 0573-2450-14

Treats Frequent Heartburn



Capsules

**14 CAPSULES**  
One 14-day course of treatment

May take 1 to 4 days for full effect

**1 BOTTLE INSIDE**



ACTUAL SIZE

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.**

000078381



FPO

3 0573 2450 14 8





KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

**Drug Facts**  
**Active ingredient (in each capsule)** Purpose  
 Esomeprazole 20 mg Acid reducer  
 Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate

**Uses**  
 ■ treats frequent heartburn (occurs 2 or more days a week)  
 ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:**  
 ■ Do not use if you are allergic to esomeprazole.  
 ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
 ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
 ■ heartburn with lightheadedness, sweating or dizziness  
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness  
 ■ frequent chest pain  
 These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**  
 ■ had heartburn over 3 months. This may be a sign of a more serious condition.  
 ■ frequent wheezing, particularly with heartburn  
 ■ unexplained weight loss  
 ■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are**  
 ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**  
 ■ your heartburn continues or worsens  
 ■ you need to take this product for more than 14 days  
 ■ you need to take more than 1 course of treatment every 4 months  
 ■ you get diarrhea ■ you develop a rash or joint pain

**Drug Facts (continued)**  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ adults 18 years of age and older  
 ■ this product is to be used once a day (every 24 hours), every day for 14 days  
 ■ may take 1 to 4 days for full effect

**14-Day Course of Treatment**  
 ■ swallow 1 capsule with a glass of water before eating in the morning  
 ■ take every day for 14 days  
 ■ do not take more than 1 capsule a day  
 ■ swallow whole. Do not crush or chew capsules.  
 ■ do not use for more than 14 days unless directed by your doctor

**Repeated 14-Day Courses (if needed)**  
 ■ you may repeat a 14-day course every 4 months  
 ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor  
 ■ children under 18 years of age: ask a doctor before use.  
 Heartburn in children may sometimes be caused by a serious condition.

**Other Information**  
 ■ read the directions and warnings before use  
 ■ keep the carton. It contains important information.  
 ■ store at 20-25°C (68-77°F)

**Inactive ingredients**  
 camille, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polyisobutyl 80, sucrose, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
 call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.



Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
 ©2022 GSK group of companies or its licensor  
 Made in France  
 For most recent product information, visit www.Nexium24HR.com  
 Nexium is a registered trademark of AstraZeneca AB and is used under license.

**Nexium<sup>®</sup> 24HR**  
 esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

**Clear Minis<sup>™</sup>**

See new warning information

Treats Frequent Heartburn

**14 CAPSULES**  
 One 14-day course of treatment

May take 1 to 4 days for full effect

**1 BOTTLE INSIDE**

**Nexium<sup>®</sup> 24HR**  
 esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

ACTUAL SIZE

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

000078392

FPO

3 0573 2452 14 2

**KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.**

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each capsule)</b> Purpose Esomeprazole 20 mg Acid reducer (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)</p> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>treats frequent heartburn (occurs 2 or more days a week)</li> <li>not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</li> </ul> <p><b>Warnings</b></p> <p><b>Allergy alert:</b></p> <ul style="list-style-type: none"> <li>Do not use if you are allergic to esomeprazole.</li> <li>Esomeprazole may cause severe skin reactions. Symptoms may include: skin reddening, blisters, rash</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use if you have:</b></p> <ul style="list-style-type: none"> <li>trouble or pain swallowing food, vomiting with blood, or bloody or black stools</li> <li>heartburn with lightheadedness, sweating or dizziness</li> <li>chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>frequent chest pain</li> </ul> <p>These may be signs of a serious condition. See your doctor.</p> <p><b>Ask a doctor before use if you have</b></p> <ul style="list-style-type: none"> <li>had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>frequent wheezing, particularly with heartburn</li> <li>unexplained weight loss</li> <li>nausea or vomiting</li> <li>stomach pain</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are</b></p> <ul style="list-style-type: none"> <li>taking a prescription drug. Acid reducers may interact with certain prescription drugs.</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>your heartburn continues or worsens</li> <li>you need to take this product for more than 14 days</li> </ul>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>you need to take more than 1 course of treatment every 4 months</li> <li>you get diarrhea</li> <li>you develop a rash or joint pain</li> </ul> <p><b>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>adults 18 years of age and older</li> <li>this product is to be used once a day (every 24 hours), every day for 14 days</li> <li>may take 1 to 4 days for full effect</li> </ul> <p><b>14-Day Course of Treatment</b></p> <ul style="list-style-type: none"> <li>swallow 1 capsule with a glass of water before eating in the morning</li> <li>take every day for 14 days</li> <li>do not take more than 1 capsule a day</li> <li>swallow whole. Do not crush or chew capsules.</li> <li>do not use for more than 14 days unless directed by your doctor</li> </ul> <p><b>Repeated 14-Day Courses (if needed)</b></p> <ul style="list-style-type: none"> <li>you may repeat a 14-day course every 4 months</li> <li>do not take for more than 14 days or more often than every 4 months unless directed by a doctor</li> <li>children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</li> </ul> <p><b>Other Information</b></p> <ul style="list-style-type: none"> <li>read the directions and warnings before use</li> <li>keep the carton. It contains important information.</li> <li>store at 20-25°C (68-77°F)</li> </ul> <p><b>Inactive ingredients</b></p> <p>corn starch, D&amp;C red no. 28, FD&amp;C blue no. 1, FD&amp;C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p><b>Questions or comments?</b></p> <p>call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600</p>
---	--

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK group of companies or its licensor  
Made in France  
For most recent product information, visit [www.Nexium24HR.com](http://www.Nexium24HR.com)  
Nexium and the colors purple and gold as applied to the AstraZeneca AB and are used under license.

Easy Open Cap!

**Nexium<sup>®</sup> 24HR**  
Capsules

See new warning information  
**Treats Frequent Heartburn**

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer

**28 CAPSULES**

Two 14-day courses of treatment

May take 1 to 4 days for full effect

**THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

000078385

**FPO**

3 0573 2450 28 5


**2 BOTTLES INSIDE**

**EASY OPEN CAP!**

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules 20 mg/acid reducer  
Capsules

ACTUAL SIZE

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.**





KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

**Drug Facts**

**Active ingredient (in each capsule)** Purpose  
Esomeprazole 20 mg Acid reducer  
(Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)

**Uses**  
■ treats frequent heartburn (occurs 2 or more days a week)  
■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

**Warnings**  
**Allergy alert:**  
■ Do not use if you are allergic to esomeprazole.  
■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use if you have:**  
■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools  
■ heartburn with lightheadedness, sweating or dizziness  
■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness  
■ frequent chest pain  
These may be signs of a serious condition. See your doctor.

**Ask a doctor before use if you have**  
■ had heartburn over 3 months. This may be a sign of a more serious condition.  
■ frequent wheezing, particularly with heartburn  
■ unexplained weight loss  
■ nausea or vomiting ■ stomach pain

**Ask a doctor or pharmacist before use if you are**  
■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**  
■ your heartburn continues or worsens  
■ you need to take this product for more than 14 days

**Drug Facts (continued)**

■ you need to take more than 1 course of treatment every 4 months  
■ you get diarrhea ■ you develop a rash or joint pain  
If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**  
■ adults 18 years of age and older  
■ this product is to be used once a day (every 24 hours), every day for 14 days  
■ may take 1 to 4 days for full effect  
**14-Day Course of Treatment**  
■ swallow 1 capsule with a glass of water before eating in the morning  
■ take every day for 14 days  
■ do not take more than 1 capsule a day  
■ swallow whole. Do not crush or chew capsules.  
■ do not use for more than 14 days unless directed by your doctor  
**Repeated 14-Day Courses (if needed)**  
■ you may repeat a 14-day course every 4 months  
■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor  
■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other Information**  
■ read the directions and warnings before use  
■ keep the carton. It contains important information.  
■ store at 20-25°C (68-77°F)

**Inactive ingredients**  
corn starch, D&C red no. 28, FD&C blue no. 1, FD&C red no. 40, ferrous oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polyorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

**Questions or comments?**  
call toll-free weekdays 9 AM to 5 PM EST at 1-866-226-1600


**Tips for Managing Heartburn**  
• Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.  
• Eat slowly and do not eat big meals.  
• Do not eat late at night or just before bedtime.  
• Do not lie flat or bend over soon after eating.  
• Raise the head of your bed.  
• Wear loose-fitting clothing around your stomach.  
• If you are overweight, lose weight.  
• If you smoke, quit smoking.



Distributed by:  
GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK group of companies or its licensor  
Made in France  
For most recent product information, visit [www.Nexium24HR.com](http://www.Nexium24HR.com)  
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

**Nexium<sup>®</sup> 24HR** Capsules  
Clinically Proven to Treat Frequent Heartburn

See new warning information  
Treats Frequent Heartburn



**42 CAPSULES**  
Three 14-day courses of treatment  
May take 1 to 4 days for full effect



Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

NDC 68750-010-01

See new warning information

Treats Frequent Heartburn

# Nexium<sup>®</sup>

esomeprazole magnesium  
delayed-release capsules  
20 mg/acid reducer

## 24HR

May take 1 to 4 days for full effect

3 Pack

Capsules

Flange of Blister

Area covered by  
PRODUCT

42  
CAPSULES

Three 14-day courses  
of treatment

Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.

# Nexium<sup>®</sup>

esomeprazole magnesium  
delayed-release capsules  
20 mg/acid reducer

## 24HR

### Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

### KEEP CARD FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.

<b>Drug Facts</b>		<b>Drug Facts (continued)</b>	
<b>Active ingredient (in each capsule)</b> Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	<b>Purpose</b> Acid reducer	<ul style="list-style-type: none"> <li>■ you need to take more than 1 course of treatment every 4 months</li> <li>■ you get diarrhea</li> <li>■ you develop a rash or joint pain</li> </ul>	
<b>Uses</b> ■ treats frequent heartburn (occurs 2 or more days a week) ■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect	<b>Warnings</b> <b>Allergy alert:</b> ■ Do not use if you are allergic to esomeprazole. ■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If an allergic reaction occurs, stop use and seek medical help right away. <b>Do not use if you have:</b> ■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath, sweating, pain spreading to arms, neck or shoulders, or lightheadedness ■ frequent chest pain These may be signs of a serious condition. See your doctor. <b>Ask a doctor before use if you have</b> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain <b>Ask a doctor or pharmacist before use if you are</b> ■ taking a prescription drug. Acid reducers may interact with certain prescription drugs. <b>Stop use and ask a doctor if</b> ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days	<p><b>if pregnant or breast-feeding</b>, ask a health professional before use. <b>Keep out of reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b> ■ adults 18 years of age and older ■ this product is to be used once a day (every 24 hours), every day for 14 days ■ may take 1 to 4 days for full effect</p> <p><b>14-Day Course of Treatment</b> ■ swallow 1 capsule with a glass of water before eating in the morning ■ take every day for 14 days ■ do not take more than 1 capsule a day ■ swallow whole. Do not crush or chew capsules. ■ do not use for more than 14 days unless directed by your doctor</p> <p><b>Repeated 14-Day Courses (if needed)</b> ■ you may repeat a 14-day course every 4 months ■ do not take for more than 14 days or more often than every 4 months unless directed by a doctor ■ children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.</p> <p><b>Other Information</b> ■ read the directions and warnings before use ■ keep the card. It contains important information. ■ store at 20-25°C (68-77°F)</p> <p><b>Inactive ingredients</b> corn starch, D&amp;C red no. 28, FD&amp;C blue no. 1, FD&amp;C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate</p> <p><b>Questions or comments?</b> call toll-free weekdays 9 AM to 5 PM EST at 1-800-245-1040</p>	



Distributed by: GSK Consumer Healthcare, Warren, NJ 07059  
©2022 GSK group of companies or its licensor  
Made in France  
For most recent product information, visit [www.Nexium24HR.com](http://www.Nexium24HR.com)  
Nexium and the colors purple and gold as applied to the capsule are registered trademarks of AstraZeneca AB and are used under license.

LOT/EXP  
WHITE HOLDOUT  
REMOVE VARNISH

000078555



**KEEP CARTON FOR COMPLETE WARNINGS AND IMPORTANT INFORMATION.**

<b>Drug Facts</b>	<b>Purpose</b>
<b>Active ingredient (in each capsule)</b> Esomeprazole 20 mg (Each delayed-release capsule corresponds to 22.3 mg esomeprazole magnesium trihydrate)	Acid reducer
<b>Uses</b>	
<ul style="list-style-type: none"> <li>■ treats frequent heartburn (occurs 2 or more days a week)</li> <li>■ not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect</li> </ul>	
<b>Warnings</b>	
<b>Allergy alert:</b>	
<ul style="list-style-type: none"> <li>■ Do not use if you are allergic to esomeprazole.</li> <li>■ Esomeprazole may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash</li> <li>■ If an allergic reaction occurs, stop use and seek medical help right away.</li> </ul>	
<b>Do not use if you have:</b>	
<ul style="list-style-type: none"> <li>■ trouble or pain swallowing food, vomiting with blood, or bloody or black stools</li> <li>■ heartburn with lightheadedness, sweating or dizziness</li> <li>■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</li> <li>■ frequent chest pain</li> </ul>	
These may be signs of a serious condition. See your doctor.	
<b>Ask a doctor before use if you have</b>	
<ul style="list-style-type: none"> <li>■ had heartburn over 3 months. This may be a sign of a more serious condition.</li> <li>■ frequent wheezing, particularly with heartburn</li> <li>■ unexplained weight loss</li> <li>■ nausea or vomiting ■ stomach pain</li> </ul>	
<b>Ask a doctor or pharmacist before use if you are</b>	
<ul style="list-style-type: none"> <li>■ taking a prescription drug. Acid reducers may interact with certain prescription drugs.</li> </ul>	
<b>Stop use and ask a doctor if</b>	
<ul style="list-style-type: none"> <li>■ your heartburn continues or worsens</li> <li>■ you need to take this product for more than 14 days</li> <li>■ you need to take more than 1 course of treatment every 4 months</li> </ul>	

**Drug Facts (continued)**

- you get diarrhea
- you develop a rash or joint pain

**Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

**14-Day Course of Treatment**

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor
- Repeated 14-Day Courses (if needed)
- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

**Other Information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20-25°C (68-77°F)

**Inactive ingredients**  
carmine, corn starch, FD&C blue no. 2, FD&C blue no. 2 aluminum lake, FD&C red no. 3, FD&C red no. 40, ferric oxide, gelatin, glyceryl monostearate, hydroxypropyl cellulose, hypromellose, magnesium stearate, methacrylic acid copolymer, pharmaceutical ink, polysorbate 80, sucrose, talc, titanium dioxide, triethyl citrate

**Questions or comments?** call toll-free weekdays 9 AM to 5 PM EST at 1 866-226-1600

**Tips for Managing Heartburn**

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.
- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules  
20 mg/acid reducer

Distributed by: GSK Consumer Healthcare, Warren, NJ 07059 ©2022 GSK group of companies or its licensors  
Made in France  
For most recent product information, visit [www.Nexium24HR.com](http://www.Nexium24HR.com)  
Nexium is a registered trademark of AstraZeneca AB and is used under license.

**Nexium<sup>®</sup> 24HR** | Clinically Proven to Treat Frequent Heartburn

**Clear Minis<sup>™</sup>**

See new warning information

**Treats Frequent Heartburn**

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules  
20 mg/acid reducer

**Clear Minis<sup>™</sup>**

May take 1 to 4 days for full effect

**42 CAPSULES**

Three 14-day courses of treatment

000078397

**FPO**


3 0573 2452 42 5

**3 BOTTLES INSIDE**

**Nexium<sup>®</sup> 24HR**  
esomeprazole magnesium delayed-release capsules  
20 mg/acid reducer

ACTUAL SIZE

**Do Not Use if seal under bottle cap imprinted with "SEALED for YOUR PROTECTION" or yellow band around the center of each capsule is broken or missing.**





-----  
**This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.**  
-----

/s/  
-----

NUSHIN F TODD  
09/06/2022 11:09:55 AM