



EXP MM / YYYY

40110065057200 NITa

Tracé contour/ Profile Rainage/

Fold
Mi-chair/
Half cutting

Pré-découpe/ Perforation

7106173

WARNING: Accidental overdose of iron-containing products is a leading cause of fatal poisoning in children under 6.

Keep this product out of reach of children.

In case of accidental overdose, call a doctor or poison control center immediately.



Usual dosage: 1 capsule twice a day on an empty stomach. See package insert for full prescribing information. Read the Patient Insert before

use.
For information about this

product call 1-888-963-6267 Patents 7459569, 9248148, 9802973, 10179120

Product of the UK

*Each capsule contains 30 mg iron (as ferric maltol) NDC 73059-001-10

Inactive ingredients: lactose monohydrate, sodium laurilsulfate, magnesium stearate, colloidal anhydrous silica, crospovidone, hypromellose, titanium dioxide, FD&C Blue No. 1, FD&C Red No. 40, FD&C Yellow No. 6.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F). (See USP controlled room temperature).

Distributed by Shield Therapeutics Inc NDC 73059-001-10



30 mg

For oral administration only

10 capsules

Professional Sample -Not For Sale









30 mg

For oral administration only


10 capsules




cx 616 a2i



PLEASE NOTE:
WHILST ARK PHARMA GRAPHICS WILL MAKE
EVERY EFFORT TO ENSURE THE ACCURACY OF TEXT,
BRAILLE AND BARCODES ARE CAPTURED, IT IS THE
ULTIMATE RESPONSIBILITY OF THE CUSTOMER'S
REGULATORY DEPARTMENT AND PRODUCTION
SITES TO ENSURE THAT TEXT (INCLUDING BRAILLE)
AND BARCODES ARE CORRECT IN BOTH CONTENT
AND ACCURACY.





Digitally signed by Ramesh Raghavachari

Date: 6/01/2022 02:31:34PM

GUID: 502d0913000029f375128b0de8c50020

(