

Drug Facts		000079243
Active ingredient (in each capsule) Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*Fever reducer (present as the free acid and potassium salt) *nonsteroidal anti-inflammatory drug	Purpose Pain reliever/fever reducer	
Uses		Drug Facts (continued) Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug
<ul style="list-style-type: none"> ■ Temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ toothache ■ backache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis ■ temporarily reduces fever 		

PANEL 1

Drug Facts (continued)	Drug Facts (continued)
<ul style="list-style-type: none"> ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed <p>Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery 	<p>Ask a doctor before use if</p> <ul style="list-style-type: none"> ■ stomach bleeding warning applies to you ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ taking any other drug

PANEL 2

Drug Facts (continued)	Drug Facts (continued)
<p>When using this product</p> <ul style="list-style-type: none"> ■ take with food or milk if stomach upset occurs <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: <ul style="list-style-type: none"> ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling 	<ul style="list-style-type: none"> ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear <p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>

BASE PANEL

000079243

DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	9 pt
• DRUG FACTS CONTINUED	8 pt
• HEADINGS	8 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.5 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

UNPRINTED AREA
FOR DATE & LOT
(UNVARNISHED)

PRINTED AREA FOR DATE & LOT
LASER CODING (VARNISHED)

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts
Active ingredient (in each capsule): Solubilized Ibuprofen (NSAID) 200 mg (equivalent to 200 mg of ibuprofen) (NSAID) (present as the free acid and potassium salt) (nonsteroidal anti-inflammatory drug)

Purpose: Pain reliever/fever reducer

Uses:
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache ■ menstrual cramps ■ the common cold ■ muscular aches ■ minor pain of arthritis
 ■ temporarily relieves fever

Warnings:
 Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing) ■ stomach pain ■ skin redness/itching ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
 Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing aspirin or naproxen (NSAIDs, aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
 Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery
 Ask a doctor before use if:
 ■ stomach bleeding warning applies to you ■ you have problems or surgery, side effects from taking pain relievers or fever reducers ■ you have a history of stomach problems, such as heartburn, ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke. ■ you are taking a diuretic.
 Ask a doctor or pharmacist before use if you are:
 ■ older than 65 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin. ■ taking any other drug.
 When using this product:
 ■ take with food or milk if stomach upset occurs

Drug Facts (continued)
 Stop use and ask a doctor if:
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ chest pain ■ trouble breathing ■ weakness in one part of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area. ■ any new symptoms appear
 If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:
 ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist. ■ If pain or fever does not respond to 1 capsule, 2 capsules may be used. ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information:
 ■ each capsule contains 200 mg of ibuprofen (NSAID) ■ each capsule contains 200 mg of potassium bitartrate (KCl) ■ each capsule contains 200 mg of hydroxypropyl methylcellulose (HPMC) ■ each capsule contains 200 mg of hydroxypropyl methylcellulose (HPMC) ■ each capsule contains 200 mg of hydroxypropyl methylcellulose (HPMC) ■ each capsule contains 200 mg of hydroxypropyl methylcellulose (HPMC)

Questions or comments? Call 1-800-89-4ADVIL
 Do not use if seal under bottle cap is cracked or missing. **SEAL FOR YOUR PROTECTION** is broken or missing.

For most retail products, visit www.advil.com or call 1-800-89-4ADVIL. For more information, visit www.advil.com.
 ©2022 GSK. All rights reserved. GSK and the GSK logo are trademarks owned or licensed by GSK.
 Facts: www.pfudat.com

LIQUI-GELS® is a trademark or registered trademark of Cabot Pharmaceuticals.

000078887

FPO

3 0573 1769 31 9

128 CODE AREA

128 CODE AREA

128 CODE AREA

128 CODE AREA

10 FREE Liquid Filled Capsules

Smaller Capsule Same Strength*

Advil LIQUI-GELS®

Solubilized Ibuprofen Capsules, 200 mg Pain Reliever/Fever Reducer (NSAID)

*Compared to Advil® Liqui-Gels®

20 30 Liquid Filled Capsules

000078887

MODIFIED DRUG FACTS TEXT DEFINED	TYPE SIZE
• DRUG FACTS TITLE	8 pt
• DRUG FACTS CONTINUED	7 pt
• HEADINGS	7 pt
• SUBHEADINGS/BODY TEXT	6 pt
• LEADING	6.25 pt
• # OF CHARACTERS PER INCH	<39
• BULLETS	5 pt
• SPACE BEFORE BULLET	2 ems
• BARLINES, HAIRLINES	1.5 pt, .5 pt
• SPACE BETWEEN HAIRLINES AND BOX END	2 spaces

This is a representation of an electronic record that was signed electronically. Following this are manifestations of any and all electronic signatures for this electronic record.

/s/

JODY E GREEN
02/23/2023 10:12:47 AM