

TAKE TASIGNA® ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

**Tasigna**® (nilotinib) capsules 200 mg per capsule Dosage: See Prescribing Information.

TAKE TASIGNA® ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.




46300682 US


@Novartis

**b** novartis





# Tasigna® (nilotinib) capsules 200 mg per capsule





U NOVARTIS

NDC 0078-0526-87

**500 mg** 

\*\*EngiseTasigna\*\*
(nilotinib) capsules

or call 1-866-411-Tasigna For more information, visit www.US.TASIGNA.com Keep this and all drugs out of the reach of children. J capsule contains 200 mg nilotinib [see USP Controlled Room Temperature].

Permitted between 15°C and 30°C (59°F and 86°F) Store at 20°C to 25°C (68°F to 77°F); excursions Dosage: See Prescribing Information.

For oral use.

78 cabenjes each. Contents:
4 individual blister packs containing

46300691 US

Contents:
4 individual blister packs containing
28 capsules each.



**200** mg per capsule

Manufactured by: Novartis Pharma Stein AG, Stein, Switzerland Distributed by:

Novartis Pharmaceuticals Corp., East Hanover, NJ 07936 Product of Ireland

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna



DOSE IS TAKEN. 1 HOUR AFTER THE

**TSA3J TA GNA** 

**DOSE OF TASIGNA** 

**BEFORE TAKING A** 

AT LEAST 2 HOURS AVOID FOOD FOR

EMPTY STOMACH.

TAKE TASIGNA ON

Rx Only





200 mg per capsule

**b** Novartis

# Contents: 4 individual blister packs containing 28 capsules each.

#### **Rx Only**

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS **BEFORE TAKING A** DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.

Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 200 mg nilotinib

Keep this and all drugs out of the reach of children. For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna



NDC 0078-0526-87

**200** mg

**DISPENSE WITH MEDICATION GUIDE** ATTACHED OR PROVIDED SEPARATELY.



**200** mg





205x82x89\_E





**U** NOVARTIS

#### PHYSICIAN SAMPLE - NOT FOR SALE

#### 28 capsules

TAKE TASIGNA ON **EMPTY STOMACH. AVOID FOOD FOR** AT LEAST 2 HOURS **BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST** 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.

reach of children.

Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 200 mg nilotinib Keep this and all drugs out of the

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna



**DISPENSE WITH MEDICATION GUIDE** ATTACHED OR PROVIDED SEPARATELY.

#### INSTRUCTIONS FOR USE



## **200 mg**

#### To open kit:

1. Firmly press and hold button 🕕 below. While holding this button, pull tab 😢 below.

**OPEN HERE** 

- 2. Once the card is fully out, unfold it completely.
- 3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
- 4. Remove any loose paper backing from the back of the card.

# **(** Tasigna® (nilotinib) capsules

200

**D**m

#### To open kit:



While pressing firmly on 1, pull on tab 2.



Pull

#### **Press** firmly

#### To close kit:

- 1. The card should not be damaged, it should fit flat in the box.
- 2. Fold the card back. Push the folded card back into the box.
- 3. Fold the cover-flap over to complete closure.
- 4. Store in a safe place out of the reach of children.

# **Tasigna®** (nilotinib) capsules 200 mg per capsule

@Novartis

0078-9526-90



#### PHYSICIAN SAMPLE - NOT FOR SALE



28 capsules

For oral use.

Dosage: See Prescribing Information.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

Manufactured by: Novartis Pharma Stein AG Stein, Switzerland Distributed by: Novartis Pharmaceuticals Corp. East Hanover, NJ 07936

Product of Ireland

1 capsule contains 200 mg nilotinib Keep this and all drugs out of the reach of children.

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna



**200** mg

**U** NOVARTIS



**Tasigna®** (nilotinib) capsules 200 mg per capsule













46300683 US



TAKE TASIGNA® ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

Tasigna® (nilotinib) capsules

Dosage: See Prescribing Information.

46300680 US


@Novartis

**b** novartis


TAKE TASIGNA® ON EMPTY STOMACH.
PLEASE SEE PATIENT MEDICATION GUIDE FOR FULL INFORMATION.

2





**© Tasigna**<sup>®</sup> (nilotinib) capsules 150 mg per capsule



#### PHYSICIAN SAMPLE - NOT FOR SALE

#### 28 capsules

TAKE TASIGNA ON **EMPTY STOMACH. AVOID FOOD FOR** AT LEAST 2 HOURS **BEFORE TAKING A DOSE OF TASIGNA** AND AT LEAST **1 HOUR AFTER THE** DOSE IS TAKEN.

For oral use.

Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children. For more information,

visit www.US.TASIGNA.com or call 1-866-411-Tasigna







**DISPENSE WITH MEDICATION GUIDE** ATTACHED OR PROVIDED SEPARATELY.

#### **INSTRUCTIONS FOR USE**



#### To open kit:

- 1. Firmly press and hold button 1 below. While holding this button, pull tab 2 below.
- 2. Once the card is fully out, unfold it completely.
- 3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
- 4. Remove any loose paper backing from the back of the card.



150 mg

#### To open kit:



While pressing firmly on 1, pull on tab 2.





**Press** firmly

#### To close kit:

- 1. The card should not be damaged, it should fit flat in the box.
- 2. Fold the card back. Push the folded card back into the box.
- 3. Fold the cover-flap over to complete closure.
- 4. Store in a safe place out of the reach of children.





@Novartis

0078-9592-90



#### PHYSICIAN SAMPLE - NOT FOR SALE



46300687 US

Manufactured by: Novartis Pharma Stein AG Stein, Switzerland Distributed by: Novartis Pharmaceuticals Corp. East Hanover, NJ 07936 Product of Ireland

For oral use.

Dosage: See Prescribing Information.

Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children.

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna





**U** NOVARTIS





#### 28 capsules

**TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST** 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.

Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children. For more information,

visit www.US.TASIGNA.com or call 1-866-411-Tasigna



**DISPENSE WITH MEDICATION GUIDE** ATTACHED OR PROVIDED SEPARATELY.

#### **INSTRUCTIONS FOR USE**



# 150 mg

#### To open kit:

1. Firmly press and hold button 1 below. While holding this button, pull tab 2 below.

**OPEN HERE** 

- 2. Once the card is fully out, unfold it completely.
- 3. To release the capsule, press gently on the capsule from the clear side of the card and push it through the paper backing.
- 4. Remove any loose paper backing from the back of the card.

## To open kit:



While pressing firmly on 1, pull on tab 2.





Pull

**Press** firmly

#### To close kit:

- 1. The card should not be damaged, it should fit flat in the box.
- 2. Fold the card back. Push the folded card back into the box.
- 3. Fold the cover-flap over to complete closure.
- 4. Store in a safe place out of the reach of children.

# **Tasigna** (nilotinib) capsules



**ONovartis** 

NDC 0078-0592-51



0078-0592-

#### 28 capsules

Rx Only

Manufactured by: Novartis Pharma Stein AG Stein, Switzerland

Novartis Pharmaceuticals Corp. East Hanover, NJ 07936 46300686x US

For oral use.

Dosage: See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between

15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 150 mg nilotinib

Keep this and all drugs out of the reach of children.

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna





**U** NOVARTIS















(nilotinib) capsules



[see USP Controlled Room Temperature]. permitted between 15°C and 30°C (59°F and 86°F) Store at 20°C to 25°C (68°F to 77°F); excursions **Dosage:** See Prescribing Information. For oral use.

**3 HOUR AFTER THE** For more information, visit www.US.TASIGNA.com or call Keep this and all drugs out of the reach of children. DOSE OF TASIGNA 1 capsule contains 150 mg nilotinib

**BEFORE TAKING A SAUOH 2 TRABL TA AVOID FOOD FOR** ЕМРТУ STOMACH. **TAKE TASIGNA ON** 

Rx Only

DOSE IS TAKEN.

TSA3J TA QNA

28 capsules each. 4 individual blister packs containing contents:

46300690 US

**Contents:** 

4 individual blister packs containing 28 capsules each.



Distributed by: Novartis Pharmaceuticals Corp., East Hanover, NJ 07936

Novartis Pharma Stein AG, Stein, Switzerland

Product of Ireland

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna

ONovartis









**b** NOVARTIS

**Contents:** 

4 individual blister packs containing 28 capsules each.

#### Rx Only

TAKE TASIGNA ON EMPTY STOMACH. AVOID FOOD FOR AT LEAST 2 HOURS BEFORE TAKING A DOSE OF TASIGNA AND AT LEAST 1 HOUR AFTER THE DOSE IS TAKEN.

For oral use.

**Dosage:** See Prescribing Information. Store at 20°C to 25°C (68°F to 77°F); excursions permitted between 15°C and 30°C (59°F and 86°F) [see **USP Controlled Room Temperature].** 

1 capsule contains 150 mg nilotinib Keep this and all drugs out of the reach

For more information, visit www.US.TASIGNA.com or call 1-866-411-Tasigna







NDC 0078-0592-87

**DISPENSE WITH MEDICATION GUIDE** ATTACHED OR PROVIDED SEPARATELY.



205x82x89\_E





