

Reference ID: 5347109

RB Artwork and Print Specification

RB514682 TR2479418 RBL2203003

G

Delsym Adult Cough 12 Hour Grape 15 ml USA 21/04/2023

RB514683 TR2479395 RBL2202988

Delsym Adult Cough

15 ml

USA

21/04/2023

NDC 63824-171-75_ **Delsym**⁶ 15/16" 2-13/16"

Base





TAMPER EVIDENT: Do not use if the neckband printed with of is broken or missing.

Drug Facts

Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide....

Purpose . Cough suppressar

Uses temporarily relieves ■ cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants

the impulse to cough to help you get to sleep

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

■ chronic cough that lasts such as occurs with smoking, asthma, or emphysema cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ shake bottle well before use

- measure only with dosing cup provided
 do not use dosing cup with other products
 dose as follows or as directed by a doctor

■ mL = milliliter		
adults and children	10 mL every 12 hours,	
12 years of age and over	not to exceed 20 mL in 24 hours	
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours	
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours	
children under 4 years of age	do not use	

Other information

■ each 5 mL contains: sodium 7 mg ■ store at 20-25°C (68-77° F) ■ dosing cup provided



Drug Facts (continued)

Inactive ingredients citric acid anhydrous, D&C red no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1, flavor, high fructose corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639







| Dist. by:
| R8 | Health (US) | C | | A | 000000 |
| reckitt | N | Parisippany, | PT# C | | A | 60329 |
| O | 2022 RB | Health | RB000000 | 042123










89 mL (3 fl oz)

ALCOHOL FREE



DELSYM® DOSING

Measure only with dosing cup provided. Do not use dosing cup with other products		
Age (yr)	Dose	
12 years	10 mL	
to adu l t	EVERY 12 HOURS	
6 to	5 mL	
under 12	EVERY 12 HOURS	
4 to	2.5 mL	
under 6	EVERY 12 HOURS	
Under 4	Do not use	





Lot No.:

Exp. Date:

RB Artwork and Print Specification


> TR2478306 RBL2202999

Adult

Cough 12 Hour Grape

21/04/2023

 $7^{1/2}$



BACK SIDE OF TOP LAYER

Drug Facts
Active ingredient (in each 5 mL)
Destromethorphan polistrex equivalent to
30 mg dextromethorphan hydrobromide....Cough suppressant
Uses: temporarily relevers cough due to minor throat and
bronchial irritation as may occur with the common cold or
inhaled irritants; the impulse to cough to help you get to sleep.
Warnings
Do not use if you are now taking a prescription monoamine
oxidase inhibitor (MAOI) (certain drugs for depression,
psychiatric, or emotional conditions, or Parkinson's disease),
or for 2 weeks after stopping the MAOI drug. If you do not
know if your prescription drug contains an MAOI, ask a
doctor before use if you have chronic cough that lasts such
as occurs with smoking, asthma, or emphysema; osugh—bet
occurs with too much phlegm (mucus). Stop use and ask a
doctor if cough lasts more than 7 days, comes back, or

BASE LABEL

be signs of a serious breast-feeding, ask a health out of reach of children. In help or contact a Poison Contu Directions: Shake bottle	well before use. Measure only o not use dosing cup with other
adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use
Other Information, F. J.	Elisabeth and the second second

Other information: Each 5 mL contains: sodium 7 mg. Store at 20-25°C (68-77°F). Dosing cup provided.

RB Artwork and Print Specification

RB515328 TR2462502

Delsym Adult Cough 12 Hour Grape 3 fl oz USA 26/04/2022

RB514094 TR2476938 RBL2202984 H Delsym Cough 12 Hour Grape 3 fl oz USA 16/03/2023





children 4 to under 6 years of age

Other information

children under 4 years of age do not use

■ each 5 mL contains: sodium 7 mg
■ store at 20-25°C (68-77° F) ■ dosing cup provided

2.5 mL every 12 hours, not to exceed 5 mL in 24 hours



Drug Facts (continued)

Inactive ingredients citric acid anhydrous, edetate disodium, ethylcellulose, FD&C yellow no. 6, flavor, high fructose corn syrup, methylparaben, partially hydrogenate vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639











RB Artwork and Print Specification

TR2478311 RBL2203000

G

Adult Cough 12 Hour Orange USA

21/04/2023



Delsym

Lot No.:


Exp. Date:





Α

Delsym Adult Cough 12 Hour Orange 3 fl oz USA 26/04/2022



BACK SIDE OF TOP LAYER

Drug Facts
Active ingredient (in each 5 mL)
Dexiromethorphan polistrex equivalent to
30 mg dextromethorphan hydrobromide....Cough suppressant
Uses: temporarily relieves: cough due to minor throat and
pronchial irritation as may occur with the common cold or inhaled irritants; the impulse to cough to help you get to sleep.
Warnings
Do not use if you are now taking a prescription monoamineoxidase inhibitor (MAOI) (certain drugs for depression,
psychlatric, or emotional conditions, or Parkinson's disease),
por for 2 weeks after stopping the MAOI drug. If you do not
know if your prescription drug contains an MAOI, ask a
doctor before use if you have chronic cough that lasts such
as occurs with smoking, asthma, or emphysema; ough; that
occurs with too much phlegm (mucus). Stop use and ask a
doctor if cough lasts more than 7 days, comes back, or

BASE LABEL

occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:* Shake bottle well before use. Measure only with dosing oup with other products. Dose as follows or as directed by a doctor.

1	adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
	children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
	of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
1	children under 4 years of age	do not use

Other information: Each 5 mL contains: sodium 7 mg. Store at 20-25°C (68-\(\mathbb{W}\)°F). Dosing cup provided.

RB514098 TR2494983 RBL2202985

Н

Delsym Adult Cough 12 Hour Orange 3 fl oz USA 16/03/2023





RB514681 TR2478315 RBL2203001


G Delsym Adult

Cough
12 Hour Grape
5 fl oz
USA

5¹¹/₁₆

USA 21/04/2023

87/8





adults and children 12 years of age and over children 6 to under 12 years

children under 4 years of age do not use Other information:

Questions? 1-866-682-4639

Store at 20-25°C (68-77°F). Dosing cup provided.

Inactive ingredients citric acid anhydrous, D&C red
no. 33, edetate disodium, ethylcellulose, FD&C blue no. 1,
flavor, high fructose corn syrup, methylparaben, partially

glycol 3350, polysorbate 80, propylene glycol, propylenaben, purified water, sucrose, tragacanth, xanthan gum

You may also report side effects to this phone number.

10 mL every 12 hours, not to exceed 20 mL in 24 hour

5 mL every 12 hours, not to exceed 10 mL in 24 hours 2.5 mL every 12 hours, not to exceed 5 mL in 24 hours



TR2462522

Α

Delsym

Adult Cough

12 Hour Grape

5 fl oz

20/04/2022

Reference ID: 5347109

NDC 63824-171-65 Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant) 2-7/8" COUGH ALCOHOL FREE HOUR RELIEF grape Flavored Liquid 148 mL (5 fl oz)

RB Artwork and Print Specification

RB514117 TR2476952 RBL2202986

Delsym Adult

Cough 12 Hour Grape

5 fl oz USA 14/03/2023



915/32

RB Artwork and Print Specification

RB514668 TR2893441 RBL2304188

B Delsym Adult

5¹¹/₁₆

Adult Cough 12 Hour Orange 5 fl oz

13/06/2023

Reference ID: 5347109

87/8

RB515329

TR2462517

В


Delsym Adult

Cough

12 Hour Orange

5 fl oz USA

13/05/2022



Drug Facts Active ingredient (in each 5 mL) Purpose Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide....Cough suppressant Uses: temporarily relieves: cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants; the impulse to cough to help you get to Do not use if you are now taking a prescription monoamine loxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have chronic cough that lasts such as occurs with smoking, asthma, or emphysema; cough that occurs with too much phlegm (mucus). Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Shake bottle well before use. Measure only with dosing cup provided. Do not use dosing cup with other products. Dose as follows or as directed by a doctor.

adults and children 12 years | 10 mL every 12 hours, not to exceed 20 mL in 24 hours or age and over

finite exceed 2 mil. in 24 hours,
of age

children 4 to under 6 years

for age

children 4 to under 6 years

children under 4 years of age

do not use Other information: Uther Information:
Each 5 mL contains: sodium 7 mg.
Store at 20-25°C (68-77°F). Dosing cup provided.
Inactive ingredients citic acid anhydrous, edetate disodium, ethylecllulose, FD&C yellow no. 6, flavor, high fructuse corn syrup, methylparaben, partially hydrogenated vegetable oil (soybean, cottonseed), polyethylene glycol 3350, polysorbate 80, propylene glycol, propylparaben, purified water, sucrose, tragacanth, xanthan gum

Questions? 1-866-682-4639

ou may also report side effects to this phone number

Delsym Destromethorphan Polistirex Extended-Release Suspension (Cough Suppressant) Cough Alcohol FREE Payored Liquid Alcohol Flavored Liquid 148 mL (5 fl oz)

RB Artwork and Print Specification

RB514116 TR2496484 RBL2202987

G

Delsym Adult Cough 12 Hour Orange 5 fl oz USA 14/03/2023





RB537209 TR2616547 RBL2209707 F Delsym Adult

Delsym Adult Cough 12 Hour Grape 2 x 5 fl oz USA 24/04/2023

15¹⁵/₃₂

963/64





RB537039
TR2610668
RBL2209706
F
Delsym
Adult
Cough

Delsym Adult Cough 12 Hour Orange 2 x 5 fl oz USA 24/04/2023

963/64



RB Artwork and Print Specification

TR2490427 RBL2203004


Delsym

4⁵/₁₆

Children's Cough 12 Hour Grape

3 fl oz USA 21/04/2023

Reference ID: 5347109



BACK SIDE OF TOP LAYER

Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide....Cough suppressant Uses: temporarily relieves: cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants; the impulse to cough to help you get to sleep.

Warnings

Do not use if you are now taking a prescription monoamine-loxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emboroal conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have chronic cough that lasts such as occurs with smoking, asthma, or emphysema; cough that accurs with too much phlegm (mucus). Stop use and ask a doctor if cough lasts more than 7 days, comes back, or

BASE LABEL

occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Shake bottle well before use. Measure only with dosing cup provided. Do not use dosing cup with other products. Dose as follows or as directed by a doctor.

		10 mL every 12 hours, not to exceed 20 mL in 24 hours
	children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
	children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
4	children under 4 years of age	do not use

Other information: Each 5 mL contains: sodium 7 mg. Store at 20-25°C (68-77°F). Dosing cup provided.

RB Artwork and Print Specification

RB517704

TR2476874

Delsym Children's Cough 12 Hour Grape 3 fl oz USA 26/04/2022

RB514120 TR2489256 RBL2202993

Delsym Children's Cough 12 Hour Grape 3 fl oz USA 16/03/2023






RB514509 TR2490410 RBL2203005

Delsym Children's Cough 12 Hour Orange 3 fl oz

USA 21/04/2023

4⁵/₁₆



BACK SIDE OF TOP LAYER

Drug Facts
Active ingredient (in each 5 mL)
Dextromethorphan polis lirex equivalent to 30 mg dextromethorphan hydrobromide.....Cough suppressant USes: temporarily relieves: cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants; the impulse to cough to help you get to sleep. Warnings
Do not use if you are now taking a prescription monoamine-oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor before use if you have chronic cough that lasts such as occurs with smoking, asthma, or emphysema; coeyal-bet occurs with too much phlegm (mucus). Stop use and ask a doctor if cough lasts more than 7 days, comes back, or

BASE LABEL

occurs with fever, rash, or headache that lasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Directions: Shake bottle well before use. Measure only with dosing cup provided. Do not use dosing cup with other products. Dose as follows or as directed by a doctor.

	adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
	children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
1	children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
1	children under 4 years of age	do not use

Other information: Each 5 mL contains: sodium 7 mg. Store at 20-25°C (68- 77°F). Dosing cup provided.

RB Artwork and Print Specification

RB517697

TR2476856



Delsym Children's Cough 12 Hour Orange 3 fl oz USA 26/04/2022

RB514119
TR2489236
RBL2202994
H
Delsym
Children's Cough 12 Hour Orange 3 fl oz USA 16/03/2023







TR2490449 RBL2203006

G

Children's

12 Hour Grape

5 fl oz


21/04/2023


RB517708 TR2476894


A Delsym

Children's Cough

12 Hour Grape 5 fl oz USA 20/04/2022







2-1/4"_ children's Dextromethorphan Polistirex Extended-Release Suspension (Cough Suppressant) AGES 4+ 2-7/8" COUGH ALCOHOL FREE HOUR RELIEF DAY or NIGHT For Children & Adults grape Flavored Liquid 148 mL (5 fl oz)

RB Artwork and Print Specification

RB514123 TR2489261 RBL2202995

G

Delsym Children's Cough 12 Hour Grape 5 fl oz USA 14/03/2023

This is a representation of an electronic record that was signed
electronically. Following this are manifestations of any and all
electronic signatures for this electronic record.

/s/

NUSHIN F TODD 03/15/2024 09:57:45 AM